

Averages of EPSONDE Microstructure Profiles from the Shallow Site, R/V Seward Johnson cruise SJ9506 in the Gulf of Maine and Georges Bank in 1995 as part of the U.S. GLOBEC program (GB project)

Website: <https://www.bco-dmo.org/dataset/2425>

Data Type: Cruise Results

Version: 1

Version Date: 2004-09-10

Project

» [U.S. GLOBEC Georges Bank](#) (GB)

Program

» [U.S. GLOBal ocean ECosystems dynamics](#) (U.S. GLOBEC)

| Contributors | Affiliation | Role |
|--------------------------------|---|------------------------|
| Hebert, Dave | University of Rhode Island (URI-GSO) | Principal Investigator |
| Allison, Dicky | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

Averages of EPSONDE Microstructure Profiles from the Shallow Site, R/V Seward Johnson cruise SJ9506 in the Gulf of Maine and Georges Bank in 1995 as part of the U.S. GLOBEC program.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: Lat:41.13 Lon:-67.75

Temporal Extent: 1995-04 - 1995-04

Dataset Description

Averages of EPSONDE Microstructure Profiles

Shallow Site, Seward Johnson 95-06

Typically, an EPSONDE microstructure profile is the average of 10 deployments of the EPSONDE instrument. One microstructure profile per profile number. During Seward Johnson cruise SJ9508, 20 deployments were made during each sampling event resulting in 2 microstructure profiles per profile number.

DMO Note: This data set as submitted contained the parameter "station". In the context of our definition, station was used incorrectly. We have changed this parameter to "profile" with the definition of: consecutively

numbered averaged EPSONDE profile. Users consulting the GSO Tech. Report should equate profile number with station number.

Prepared by: Russ Burgett, University of Rhode Island, GSO
Reference: (S06 - GL1AVG)

Details of data processing are described in: U.S. GLOBEC Georges Bank microstructure data. GSO Univ. Rhode Island Tech. report 96-6.

Contributor:

Dave Hebert
Graduate School of Oceanography
Univ. of Rhode Island
Narragansett, RI 02882-1197

phone: 401 874 6610

fax: 401 874 6728

e-mail: david.hebert@gso.uri.edu Updated: September 10, 2004; gfh

Methods & Sampling

Typically, an EPSONDE microstructure profile is the average of 10 deployments of the EPSONDE instrument. One microstructure profile per profile number. During Seward Johnson cruise SJ9508, 20 deployments were made during each sampling event resulting in 2 microstructure profiles per profile number.

Data Processing Description

This data set as submitted contained the parameter "station". In the context of our definition, station was used incorrectly. We have changed this parameter to "profile" with the definition of: consecutively numbered averaged EPSONDE profile. Users consulting the GSO Tech. Report should equate profile number with station number.

[[table of contents](#) | [back to top](#)]

Data Files

| File |
|--|
| S06_rs.csv (Comma Separated Values (.csv), 55.89 KB) MD5:094dc9334f083367b253d357795b1cad Primary data file for dataset ID 2425 |

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|-----------|---|--------------------------------|
| cruiseid | cruise identification | |
| lat | latitude, negative = South | decimal degrees |
| lon | longitude, negative = West | decimal degrees |
| profile | consecutively numbered EPSONDE profile | |
| yrday_gmt | year day, Julian calender | decimal day, GMT |
| press | depth of sample, reported as pressure | decibars |
| temp | temperature | degrees C |
| sal | salinity | PSU |
| sigma_t | density | kilograms/meter ³ |
| eps | epsilon, turbulent kinetic energy dissipation | watts/kilogram |
| chi_t | chi_theta, temperature variance dissipation | degrees C ² /second |
| k_t | K_t, vertical diffusivity for heat | meters ² /second |
| k_rho | K_rho, vertical diffusivity for density | meters ² /second |

[[table of contents](#) | [back to top](#)]

Instruments

| | |
|---|--|
| Dataset-specific Instrument Name | EPSONDE |
| Generic Instrument Name | EPSONDE |
| Dataset-specific Description | An EPSONDE microstructure profile is the average of 10 deployments of the EPSONDE instrument. One microstructure profile per profile number. During Seward Johnson cruise SJ9508, 20 deployments were made during each sampling event resulting in 2 microstructure profiles per profile number. |
| Generic Instrument Description | An EPSONDE is a tethered free-fall profiling system used to obtain temperature microstructure and velocity turbulence data in the water column. The EPSONDE profiler carries a variety of slow and fast sensors for measuring temperature microstructure, velocity microstructure, conductivity and depth. These data yield turbulent kinetic energy dissipation rates and temperature variance dissipation rates as well as derived quantities such as turbulent diffusivity. |

[[table of contents](#) | [back to top](#)]

Deployments

SJ9506

| | |
|--------------------|--|
| Website | https://www.bco-dmo.org/deployment/57485 |
| Platform | R/V Seward Johnson |
| Report | http://globec.who.edu/globec-dir/reports/sj9506.html |
| Start Date | 1995-04-25 |
| End Date | 1995-05-03 |
| Description | <p>this was a process cruise. Process turbulence.</p> <p>Methods & Sampling Typically, an EPSONDE microstructure profile is the average of 10 deployments of the EPSONDE instrument. One microstructure profile per profile number. During Seward Johnson cruise SJ9508, 20 deployments were made during each sampling event resulting in 2 microstructure profiles per profile number.</p> <p>Processing Description This data set as submitted contained the parameter "station". In the context of our definition, station was used incorrectly. We have changed this parameter to "profile" with the definition of: consecutively numbered averaged EPSONDE profile. Users consulting the GSO Tech. Report should equate profile number with station number.</p> |

[[table of contents](#) | [back to top](#)]

Project Information

U.S. GLOBEC Georges Bank (GB)

Website: http://globec.who.edu/globec_program.html

Coverage: Georges Bank, Gulf of Maine, Northwest Atlantic Ocean

The U.S. GLOBEC [Georges Bank](#) Program is a large multi-disciplinary multi-year oceanographic effort. The proximate goal is to understand the population dynamics of key species on the Bank - Cod, [Haddock](#), and two species of zooplankton ([Calanus finmarchicus](#) and [Pseudocalanus](#)) - in terms of their coupling to the physical environment and in terms of their [predators and prey](#). The ultimate goal is to be able to predict changes in the distribution and abundance of these species as a result of changes in their physical and biotic environment as well as to anticipate how their populations might respond to climate change.

The effort is substantial, requiring broad-scale surveys of the entire Bank, and process studies which focus both on the links between the target species and their physical environment, and the determination of fundamental aspects of these species' life history (birth rates, growth rates, death rates, etc).

Equally important are the modelling efforts that are ongoing which seek to provide realistic predictions of the flow field and which utilize the life history information to produce an integrated view of the dynamics of the populations.

The U.S. GLOBEC Georges Bank [Executive Committee \(EXCO\)](#) provides program leadership and effective communication with the funding agencies.

[[table of contents](#) | [back to top](#)]

Program Information

U.S. GLOBAL ocean ECosystems dynamics (U.S. GLOBEC)

Website: <http://www.usglobec.org/>

Coverage: Global

U.S. GLOBEC (GLOBAL ocean ECosystems dynamics) is a research program organized by oceanographers and fisheries scientists to address the question of how global climate change may affect the abundance and production of animals in the sea.

The U.S. GLOBEC Program currently had major research efforts underway in the Georges Bank / Northwest Atlantic Region, and the Northeast Pacific (with components in the California Current and in the Coastal Gulf of Alaska). U.S. GLOBEC was a major contributor to International GLOBEC efforts in the Southern Ocean and Western Antarctic Peninsula (WAP).

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|---------------------------------|
| National Science Foundation (NSF) | unknown GB NSF |
| National Oceanic and Atmospheric Administration (NOAA) | unknown GB NOAA |

[[table of contents](#) | [back to top](#)]