

# Phytoplankton growth rates and microzooplankton grazing from R/V Alpha Helix HX242, HX244, HX247, HX271, HX275 in the Northeast Pacific from 2001-2003

**Website:** <https://www.bco-dmo.org/dataset/2475>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2007-03-28

## Project

» [U.S. GLOBEC Northeast Pacific](#) (NEP)

## Program

» [U.S. GLOBal ocean ECosystems dynamics](#) (U.S. GLOBEC)

Contributors	Affiliation	Role
<a href="#">Strom, Suzanne</a>	Western Washington University (WWU)	Principal Investigator
<a href="#">Batchelder, Hal</a>	Oregon State University (OSU-CEOAS)	Data Manager
<a href="#">Allison, Dicky</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Phytoplankton growth rates and microzooplankton grazing from R/V Alpha Helix HX242, HX244, HX247, HX271, HX275 in the Northeast Pacific from 2001-2003

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Program Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:60.535 E:-146.607 S:58.097 W:-150.419

**Temporal Extent:** 2001-04-17 - 2003-08-09

---

## Dataset Description

### Phytoplankton Growth Rates in the coastal Gulf of Alaska

Phytoplankton growth rates, the phytoplankton growth rate response to nutrient enrichment, and microzooplankton grazing rates on phytoplankton in three different size classes were measured during three cruises to the coastal Gulf of Alaska in 2001.

## Methods & Sampling

[excerpted from Strom, et al. (2006) Microzooplankton grazing in the coastal Gulf of Alaska: Variations in top-down control of phytoplankton. *Limnol Oceanogr* in press.]

Water drawn from multiple Niskin bottles closed at a single depth was pooled into two 25-liter polycarbonate carboys. Most often, water was collected from the depth corresponding to 50% of surface irradiance (50% I<sub>0</sub>, 3 to 10 m). Once during April, and once per station during July, water was collected from the depth of the subsurface chlorophyll maximum (SCM, 12 to 25 m). The contents of one carboy were gravity-filtered (0.2 μm) to generate particle-free filtered seawater (FSW, the diluent for the dilution series). The contents of the other were gently pre-screened through 200 μm Nitex mesh to exclude macrozooplankton (WSW, the whole seawater for the dilution series). Using gentle siphoning and mixing techniques, FSW and WSW were combined in known proportions in 2.35-liter polycarbonate bottles to generate a dilution series consisting of 9, 16, 24, 41, 61, and 100% WSW (each in duplicate). An additional pair of bottles diluted to 4% was added during the May and July cruises, as well as an additional pair of 100% WSW bottles to control for the effects of nutrient enrichment on phytoplankton growth rate. Clean techniques and inert materials (silicone, polycarbonate) were used throughout.

## Data Processing Description

Initial samples for size-fractionated chlorophyll (<5, 5 to 20, and >20 μm, in quadruplicate), nutrients (nitrate, nitrite, silicic acid, phosphate), and microzooplankton abundance and composition (in duplicate, see below) were taken from the WSW carboy at intervals during experiment set-up. Initial chlorophyll levels in diluted bottles were calculated from these measured WSW values and known dilution factors. Coefficients of variation for quadruplicate initial chlorophyll samples averaged 7.9%, 13.5%, and 8.9% for the <5, 5 to 20, and >20 μm size fractions, respectively. During May (all but outer shelf experiments) and July cruises, all diluted bottles and two 100% WSW bottles were enriched with nitrate (4.7 μmol L<sup>-1</sup> as NaNO<sub>3</sub>) and phosphate (0.27 μmol L<sup>-1</sup> as Na<sub>2</sub>HPO<sub>4</sub>). The other two 100% WSW bottles were left unenriched. Bottles were screened to collection-depth light levels with neutral density screening and incubated on deck in seawater-cooled incubators for 24 hr. All bottles were then sampled in duplicate for size-fractionated chlorophyll (filtration volumes ranged from 0.15 to 1.08 liter depending on WSW chlorophyll and dilution levels); 100% WSW bottles were additionally sampled for microzooplankton abundance and composition.

Net growth rates ( $k$ , d<sup>-1</sup>) for total chlorophyll and individual chlorophyll size fractions were calculated as  $(1/t) \ln[P_t/P_o]$ , where  $P_t$  = final chlorophyll concentration,  $P_o$  = initial chlorophyll concentration, and  $t$  = incubation time in d. Intrinsic growth rates ( $\mu$ , d<sup>-1</sup>) of phytoplankton were estimated from the y-intercept of net growth rates regressed upon fraction WSW. For experiments exhibiting saturated grazing (i.e. a leveling of net growth rate across the least-dilute bottles) (Gallegos 1989), intrinsic growth rate estimates were based on regression of net growth rates in only the most dilute bottles (generally those with <40% WSW). Microzooplankton grazing rates ( $g$ , d<sup>-1</sup>) were estimated from the slope of the regression for experiments with linear relationships between net growth and fraction WSW, and as  $g = \mu_n - k_n$  (where  $k_n$  = net growth rate of phytoplankton in enriched, 100% WSW bottles) for experiments with saturated grazing. In experiments with nutrient enrichment, unenriched phytoplankton growth rates ( $\mu_o$ ) were calculated as  $\mu_o = k_o + g$ , where  $k_o$  = net growth rate of phytoplankton in unenriched, 100% WSW bottles. Estimates of  $\mu_o$  were used to compare microzooplankton grazing to phytoplankton growth in situ ( $g : \mu_o$ ). These ratios represent the fraction of primary production consumed each day by microzooplankton grazing. Ratios were arctan transformed for estimation of means and standard deviations.

More detailed methods reference in Strom, *et al.* (2006).

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

**File**

**phyto\_growth.csv**(Comma Separated Values (.csv), 24.11 KB)  
 MD5:bc87c1da92a74895c867a67a0c30acc3

Primary data file for dataset ID 2475

[ [table of contents](#) | [back to top](#) ]

## Related Publications

Strom, S., Olson, M., Macri, E., & Mord, C. (2006). Cross-shelf gradients in phytoplankton community structure, nutrient utilization, and growth rate in the coastal Gulf of Alaska. *Marine Ecology Progress Series*, 328, 75–92. doi:[10.3354/meps328075](https://doi.org/10.3354/meps328075)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
cruiseid	Cruise ID.	text
expt	Experiment number.	dimensionless
date_local	Day-month-year.	dd-mon-yy
station_std	standard station number	dimensionless
lat	Latitude	decimal degrees (North is positive)
lon	Longitude	decimal degrees (East is positive)
time_start_local	experiment start time	24-hour clock
depth	water collection depth	meters
chla_start_gt20um	initial chlorophyll concentration for dilutions experiments	micrograms/liter
chla_start_5_to_20um	initial chlorophyll concentration for dilutions experiments	micrograms/liter
chla_start_lt5um	initial chlorophyll concentration for dilutions experiments	micrograms/liter
chla_start_total	sum of three size fractions	micrograms/liter
fed_phyto_growth_gt20um	enriched (addition of nitrate and phosphate) phytoplankton growth rate	per day
fed_phyto_growth_5_to_20um	enriched (addition of nitrate and phosphate) phytoplankton growth rate	per day
fed_phyto_growth_lt5um	enriched (addition of nitrate and phosphate) phytoplankton growth rate	per day
fed_phyto_growth_total	sum of three size fractions	per day
fed_phyto_growth_std_err_gt20um	enriched phytoplankton growth rate standard error	per day
fed_phyto_growth_std_err_5_to_20um	enriched phytoplankton growth rate standard error	per day
fed_phyto_growth_std_err_lt5um	enriched phytoplankton growth rate standard error	per day

fed_phyto_growth_std_err_total	total standard error	per day
unfed_phyto_growth_gt20um	unenriched phytoplankton growth rate	per day
unfed_phyto_growth_5_to_20um	unenriched phytoplankton growth rate	per day
unfed_phyto_growth_lt5um	unenriched phytoplankton growth rate	per day
unfed_phyto_growth_total	sum of three size fractions	per day
unfed_phyto_growth_std_err_gt20um	unenriched phytoplankton growth rate standard error	per day
unfed_phyto_growth_std_err_5_to_20um	unenriched phytoplankton growth rate standard error	per day
unfed_phyto_growth_std_err_lt5um	unenriched phytoplankton growth rate standard error	per day
unfed_phyto_growth_std_err_total	total standard error	per day
microzoo_graz_gt20um	microzooplankton grazing rate for >20μ size fraction	per day
microzoo_graz_5_to_20um	microzooplankton grazing rate for >5 and	per day
microzoo_graz_lt5um	microzooplankton grazing rate for	per day
microzoo_graz_total	total microzooplankton grazing rate for all size fractions	per day
microzoo_graz_std_err_gt20um	standard error	per day
microzoo_graz_std_err_5_to_20um	standard error	per day
microzoo_graz_std_err_lt5um	standard error	per day
microzoo_graz_std_err_total	total standard error	per day
temp	temperature at sample depth	degrees Celsius
sal	dalinity at sample depth	Practical Salinity Units (PSU)
irradiance_mol_per_m2	irradiance at sample depth	mol photons/meter <sup>2</sup> (per second?)

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Niskin Bottle
<b>Generic Instrument Name</b>	Niskin bottle
<b>Dataset-specific Description</b>	Niskin bottle cast, use Bottle_Niskin
<b>Generic Instrument Description</b>	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### HX242

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57523">https://www.bco-dmo.org/deployment/57523</a>
<b>Platform</b>	R/V Alpha Helix
<b>Report</b>	<a href="http://globec.whoi.edu/nep/reports/cgoa_cruises/hx242cr.pdf">http://globec.whoi.edu/nep/reports/cgoa_cruises/hx242cr.pdf</a>
<b>Start Date</b>	2001-04-17
<b>End Date</b>	2001-05-01

<b>Description</b>	<p>Original cruise data are available from the NSF R2R data catalog</p> <p><b>Methods &amp; Sampling</b>  [excerpted from Strom, et al. (2006) Microzooplankton grazing in the coastal Gulf of Alaska: Variations in top-down control of phytoplankton. <i>Limnol Oceanogr</i> in press.] Water drawn from multiple Niskin bottles closed at a single depth was pooled into two 25-liter polycarbonate carboys. Most often, water was collected from the depth corresponding to 50% of surface irradiance (50% I<sub>0</sub>, 3 to 10 m). Once during April, and once per station during July, water was collected from the depth of the subsurface chlorophyll maximum (SCM, 12 to 25 m). The contents of one carboy were gravity-filtered (0.2 μm) to generate particle-free filtered seawater (FSW, the diluent for the dilution series). The contents of the other were gently pre-screened through 200 μm Nitex mesh to exclude macrozooplankton (WSW, the whole seawater for the dilution series). Using gentle siphoning and mixing techniques, FSW and WSW were combined in known proportions in 2.35-liter polycarbonate bottles to generate a dilution series consisting of 9, 16, 24, 41, 61, and 100% WSW (each in duplicate). An additional pair of bottles diluted to 4% was added during the May and July cruises, as well as an additional pair of 100% WSW bottles to control for the effects of nutrient enrichment on phytoplankton growth rate. Clean techniques and inert materials (silicone, polycarbonate) were used throughout.</p> <p><b>Processing Description</b>  Initial samples for size-fractionated chlorophyll (20 μm, in quadruplicate), nutrients (nitrate, nitrite, silicic acid, phosphate), and microzooplankton abundance and composition (in duplicate, see below) were taken from the WSW carboy at intervals during experiment set-up. Initial chlorophyll levels in diluted bottles were calculated from these measured WSW values and known dilution factors. Coefficients of variation for quadruplicate initial chlorophyll samples averaged 7.9%, 13.5%, and 8.9% for the 20 μm size fractions, respectively. During May (all but outer shelf experiments) and July cruises, all diluted bottles and two 100% WSW bottles were enriched with nitrate (4.7 μmol L<sup>-1</sup> as NaNO<sub>3</sub>) and phosphate (0.27 μmol L<sup>-1</sup> as Na<sub>2</sub>HPO<sub>4</sub>). The other two 100% WSW bottles were left unenriched. Bottles were screened to collection-depth light levels with neutral density screening and incubated on deck in seawater-cooled incubators for 24 hr. All bottles were then sampled in duplicate for size-fractionated chlorophyll (filtration volumes ranged from 0.15 to 1.08 liter depending on WSW chlorophyll and dilution levels); 100% WSW bottles were additionally sampled for microzooplankton abundance and composition. Net growth rates (k, d<sup>-1</sup>) for total chlorophyll and individual chlorophyll size fractions were calculated as <math>(1/t)(\ln[P_t/P_o])</math>, where P<sub>t</sub> = final chlorophyll concentration, P<sub>o</sub> = initial chlorophyll concentration, and t = incubation time in d. Intrinsic growth rates (μ, d<sup>-1</sup>) of phytoplankton were estimated from the y-intercept of net growth rates regressed upon fraction WSW. For experiments exhibiting saturated grazing (i.e. a leveling of net growth rate across the least-dilute bottles) (Gallegos 1989), intrinsic growth rate estimates were based on regression of net growth rates in only the most dilute bottles (generally those with &lt;40% WSW). Microzooplankton grazing rates (g, d<sup>-1</sup>) were estimated from the slope of the regression for experiments with linear relationships between net growth and fraction WSW, and as <math>g = \mu_n - k_n</math> (where k<sub>n</sub> = net growth rate of phytoplankton in enriched, 100% WSW bottles) for experiments with saturated grazing. In experiments with nutrient enrichment, unenriched phytoplankton growth rates (μ<sub>o</sub>) were calculated as <math>\mu_o = k_o + g</math>, where k<sub>o</sub> = net growth rate of phytoplankton in unenriched, 100% WSW bottles. Estimates of μ<sub>o</sub> were used to compare microzooplankton grazing to phytoplankton growth in situ (g : μ<sub>o</sub>). These ratios represent the fraction of primary production consumed each day by microzooplankton grazing. Ratios were arctan transformed for estimation of means and standard deviations. More detailed methods reference: Strom, et al. (2006) Cross-shelf gradients in phytoplankton community structure, nutrient utilization, and growth rate in the coastal Gulf of Alaska. <i>Marine Ecology Progress Series</i> (in press)</p>
--------------------	--

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57525">https://www.bco-dmo.org/deployment/57525</a>
<b>Platform</b>	R/V Alpha Helix
<b>Report</b>	<a href="http://globec.whoi.edu/nep/reports/cgoa_cruises/hx244cr.pdf">http://globec.whoi.edu/nep/reports/cgoa_cruises/hx244cr.pdf</a>
<b>Start Date</b>	2001-05-17
<b>End Date</b>	2001-05-31
<b>Description</b>	<p>Original cruise data are available from the NSF R2R data catalog</p> <p><b>Methods &amp; Sampling</b>  [excerpted from Strom, et al. (2006) Microzooplankton grazing in the coastal Gulf of Alaska: Variations in top-down control of phytoplankton. <i>Limnol Oceanogr</i> in press.] Water drawn from multiple Niskin bottles closed at a single depth was pooled into two 25-liter polycarbonate carboys. Most often, water was collected from the depth corresponding to 50% of surface irradiance (50% I<sub>0</sub>, 3 to 10 m). Once during April, and once per station during July, water was collected from the depth of the subsurface chlorophyll maximum (SCM, 12 to 25 m). The contents of one carboy were gravity-filtered (0.2 μm) to generate particle-free filtered seawater (FSW, the diluent for the dilution series). The contents of the other were gently pre-screened through 200 μm Nitex mesh to exclude macrozooplankton (WSW, the whole seawater for the dilution series). Using gentle siphoning and mixing techniques, FSW and WSW were combined in known proportions in 2.35-liter polycarbonate bottles to generate a dilution series consisting of 9, 16, 24, 41, 61, and 100% WSW (each in duplicate). An additional pair of bottles diluted to 4% was added during the May and July cruises, as well as an additional pair of 100% WSW bottles to control for the effects of nutrient enrichment on phytoplankton growth rate. Clean techniques and inert materials (silicone, polycarbonate) were used throughout.</p> <p><b>Processing Description</b>  Initial samples for size-fractionated chlorophyll (20 μm, in quadruplicate), nutrients (nitrate, nitrite, silicic acid, phosphate), and microzooplankton abundance and composition (in duplicate, see below) were taken from the WSW carboy at intervals during experiment set-up. Initial chlorophyll levels in diluted bottles were calculated from these measured WSW values and known dilution factors. Coefficients of variation for quadruplicate initial chlorophyll samples averaged 7.9%, 13.5%, and 8.9% for the 20 μm size fractions, respectively. During May (all but outer shelf experiments) and July cruises, all diluted bottles and two 100% WSW bottles were enriched with nitrate (4.7 μmol L<sup>-1</sup> as NaNO<sub>3</sub>) and phosphate (0.27 μmol L<sup>-1</sup> as Na<sub>2</sub>HPO<sub>4</sub>). The other two 100% WSW bottles were left unenriched. Bottles were screened to collection-depth light levels with neutral density screening and incubated on deck in seawater-cooled incubators for 24 hr. All bottles were then sampled in duplicate for size-fractionated chlorophyll (filtration volumes ranged from 0.15 to 1.08 liter depending on WSW chlorophyll and dilution levels); 100% WSW bottles were additionally sampled for microzooplankton abundance and composition. Net growth rates (k, d<sup>-1</sup>) for total chlorophyll and individual chlorophyll size fractions were calculated as <math>(1/t)(\ln[P_t/P_o])</math>, where P<sub>t</sub> = final chlorophyll concentration, P<sub>o</sub> = initial chlorophyll concentration, and t = incubation time in d. Intrinsic growth rates (μ, d<sup>-1</sup>) of phytoplankton were estimated from the y-intercept of net growth rates regressed upon fraction WSW. For experiments exhibiting saturated grazing (i.e. a leveling of net growth rate across the least-dilute bottles) (Gallegos 1989), intrinsic growth rate estimates were based on regression of net growth rates in only the most dilute bottles (generally those with ~40% WSW). Microzooplankton grazing rates (g, d<sup>-1</sup>) were estimated from the slope of the regression for experiments with linear relationships between net growth and fraction WSW, and as <math>g = \mu_n - k_n</math> (where k<sub>n</sub> = net growth rate of phytoplankton in enriched, 100% WSW bottles) for experiments with saturated grazing. In experiments with nutrient enrichment, unenriched phytoplankton growth rates (μ<sub>o</sub>) were calculated as <math>\mu_o = k_o + g</math>, where k<sub>o</sub> = net growth rate of phytoplankton in unenriched, 100% WSW bottles. Estimates of μ<sub>o</sub> were used to compare microzooplankton grazing to phytoplankton growth in situ (g : μ<sub>o</sub>). These ratios represent the fraction of primary production consumed each day by microzooplankton grazing. Ratios were arctan transformed for estimation of means and standard deviations. More detailed methods reference: Strom, et al. (2006) Cross-shelf gradients in phytoplankton community structure, nutrient utilization, and growth rate in the coastal Gulf of Alaska. <i>Marine Ecology Progress Series</i> (in press)</p>

**HX247**

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57527">https://www.bco-dmo.org/deployment/57527</a>
<b>Platform</b>	R/V Alpha Helix
<b>Report</b>	<a href="http://globec.whoi.edu/nep/reports/cgoa_cruises/hx247cr.pdf">http://globec.whoi.edu/nep/reports/cgoa_cruises/hx247cr.pdf</a>
<b>Start Date</b>	2001-07-12
<b>End Date</b>	2001-07-26
<b>Description</b>	<p>Original cruise data are available from the NSF R2R data catalog</p> <p><b>Methods &amp; Sampling</b>  [excerpted from Strom, et al. (2006) Microzooplankton grazing in the coastal Gulf of Alaska: Variations in top-down control of phytoplankton. <i>Limnol Oceanogr</i> in press.] Water drawn from multiple Niskin bottles closed at a single depth was pooled into two 25-liter polycarbonate carboys. Most often, water was collected from the depth corresponding to 50% of surface irradiance (50% I<sub>0</sub>, 3 to 10 m). Once during April, and once per station during July, water was collected from the depth of the subsurface chlorophyll maximum (SCM, 12 to 25 m). The contents of one carboy were gravity-filtered (0.2 μm) to generate particle-free filtered seawater (FSW, the diluent for the dilution series). The contents of the other were gently pre-screened through 200 μm Nitex mesh to exclude macrozooplankton (WSW, the whole seawater for the dilution series). Using gentle siphoning and mixing techniques, FSW and WSW were combined in known proportions in 2.35-liter polycarbonate bottles to generate a dilution series consisting of 9, 16, 24, 41, 61, and 100% WSW (each in duplicate). An additional pair of bottles diluted to 4% was added during the May and July cruises, as well as an additional pair of 100% WSW bottles to control for the effects of nutrient enrichment on phytoplankton growth rate. Clean techniques and inert materials (silicone, polycarbonate) were used throughout.</p> <p><b>Processing Description</b>  Initial samples for size-fractionated chlorophyll (20 μm, in quadruplicate), nutrients (nitrate, nitrite, silicic acid, phosphate), and microzooplankton abundance and composition (in duplicate, see below) were taken from the WSW carboy at intervals during experiment set-up. Initial chlorophyll levels in diluted bottles were calculated from these measured WSW values and known dilution factors. Coefficients of variation for quadruplicate initial chlorophyll samples averaged 7.9%, 13.5%, and 8.9% for the 20 μm size fractions, respectively. During May (all but outer shelf experiments) and July cruises, all diluted bottles and two 100% WSW bottles were enriched with nitrate (4.7 μmol L<sup>-1</sup> as NaNO<sub>3</sub>) and phosphate (0.27 μmol L<sup>-1</sup> as Na<sub>2</sub>HPO<sub>4</sub>). The other two 100% WSW bottles were left unenriched. Bottles were screened to collection-depth light levels with neutral density screening and incubated on deck in seawater-cooled incubators for 24 hr. All bottles were then sampled in duplicate for size-fractionated chlorophyll (filtration volumes ranged from 0.15 to 1.08 liter depending on WSW chlorophyll and dilution levels); 100% WSW bottles were additionally sampled for microzooplankton abundance and composition. Net growth rates (k, d<sup>-1</sup>) for total chlorophyll and individual chlorophyll size fractions were calculated as <math>(1/t)(\ln[Pt/Po])</math>, where Pt = final chlorophyll concentration, Po = initial chlorophyll concentration, and t = incubation time in d. Intrinsic growth rates (μ, d<sup>-1</sup>) of phytoplankton were estimated from the y-intercept of net growth rates regressed upon fraction WSW. For experiments exhibiting saturated grazing (i.e. a leveling of net growth rate across the least-dilute bottles) (Gallegos 1989), intrinsic growth rate estimates were based on regression of net growth rates in only the most dilute bottles (generally those with ~40% WSW). Microzooplankton grazing rates (g, d<sup>-1</sup>) were estimated from the slope of the regression for experiments with linear relationships between net growth and fraction WSW, and as <math>g = \mu_n - k_n</math> (where <math>k_n</math> = net growth rate of phytoplankton in enriched, 100% WSW bottles) for experiments with saturated grazing. In experiments with nutrient enrichment, unenriched phytoplankton growth rates (μ<sub>0</sub>) were calculated as <math>\mu_0 = k_0 + g</math>, where <math>k_0</math> = net growth rate of phytoplankton in unenriched, 100% WSW bottles. Estimates of μ<sub>0</sub> were used to compare microzooplankton grazing to phytoplankton growth in situ (<math>g : \mu_0</math>). These ratios represent the fraction of primary production consumed each day by microzooplankton grazing. Ratios were arctan transformed for estimation of means and standard deviations. More detailed methods reference: Strom, et al. (2006) Cross-shelf gradients in phytoplankton community structure, nutrient utilization, and growth rate in the coastal Gulf of Alaska. <i>Marine Ecology Progress Series</i> (in press)</p>



## HX271

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57540">https://www.bco-dmo.org/deployment/57540</a>
<b>Platform</b>	R/V Alpha Helix
<b>Report</b>	<a href="http://globec.whoi.edu/nep/reports/cgoa_cruises/hx271cr.pdf">http://globec.whoi.edu/nep/reports/cgoa_cruises/hx271cr.pdf</a>
<b>Start Date</b>	2003-04-24
<b>End Date</b>	2003-05-15
<b>Description</b>	Original cruise data are available from the NSF R2R data catalog

## HX275

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57542">https://www.bco-dmo.org/deployment/57542</a>
<b>Platform</b>	R/V Alpha Helix
<b>Report</b>	<a href="http://globec.whoi.edu/nep/reports/cgoa_cruises/hx275cr.pdf">http://globec.whoi.edu/nep/reports/cgoa_cruises/hx275cr.pdf</a>
<b>Start Date</b>	2003-07-20
<b>End Date</b>	2003-08-12
<b>Description</b>	Original cruise data are available from the NSF R2R data catalog

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### U.S. GLOBEC Northeast Pacific (NEP)

**Website:** <http://nepglobec.bco-dmo.org>

**Coverage:** Northeast Pacific Ocean, Gulf of Alaska

### Program in a Nutshell

**Goal:** To understand the effects of climate variability and climate change on the distribution, abundance and production of marine animals (including commercially important living marine resources) in the eastern North Pacific. To embody this understanding in diagnostic and prognostic ecosystem models, capable of capturing the ecosystem response to major climatic fluctuations.

**Approach:** To study the effects of past and present climate variability on the population ecology and population dynamics of marine biota and living marine resources, and to use this information as a proxy for how the ecosystems of the eastern North Pacific may respond to future global climate change. The strong temporal variability in the physical and biological signals of the NEP will be used to examine the biophysical mechanisms through which zooplankton and salmon populations respond to physical forcing and biological interactions in the coastal regions of the two gyres. Annual and interannual variability will be studied directly through **long-term observations** and detailed **process studies**; variability at longer time scales will be examined through **retrospective analysis** of directly measured and proxy data. Coupled **biophysical models** of the ecosystems of these regions will be developed and tested using the process studies and data collected from the long-term observation programs, then further tested and improved by hindcasting selected retrospective data series.

[ [table of contents](#) | [back to top](#) ]

---

## Program Information

## U.S. GLOBAL ocean ECosystems dynamics (U.S. GLOBEC)

**Website:** <http://www.usglobec.org/>

**Coverage:** Global

U.S. GLOBEC (GLOBAL ocean ECosystems dynamics) is a research program organized by oceanographers and fisheries scientists to address the question of how global climate change may affect the abundance and production of animals in the sea.

The U.S. GLOBEC Program currently had major research efforts underway in the Georges Bank / Northwest Atlantic Region, and the Northeast Pacific (with components in the California Current and in the Coastal Gulf of Alaska). U.S. GLOBEC was a major contributor to International GLOBEC efforts in the Southern Ocean and Western Antarctic Peninsula (WAP).

[ [table of contents](#) | [back to top](#) ]

---

### Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0101397</a>
National Oceanic and Atmospheric Administration (NOAA)	<a href="#">unknown NEP NOAA</a>

[ [table of contents](#) | [back to top](#) ]