HPLC measured pigments, TM Primary Production casts from R/V Thomas G. Thompson TT045 cruise in the Arabian Sea in 1995 (U.S. JGOFS Arabian Sea project)

Website: https://www.bco-dmo.org/dataset/2549

Version: May 8, 2001 Version Date: 2001-05-08

Project

» <u>U.S. JGOFS Arabian Sea</u> (Arabian Sea)

» U.S. Joint Global Ocean Flux Study (U.S. JGOFS)

Contributors	Affiliation	Role
Bidigare, Robert R.	University of Hawaii (UH)	Principal Investigator
Chandler, Cynthia L.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- **Parameters**
- **Instruments**
- **Deployments**
- **Project Information**
- **Program Information**
- <u>Funding</u>

Dataset Description

HPLC measured pigments, samples drawn from "TM" Primary Production casts.

Methods & Sampling

Robert R. Bidigare of: University of Hawaii

Pigments, HPLC method, sampled from TM casts dataset:

March 17, 1995 to April 07, 1995 dates:

location: N: 19.883 S: 9.9988 W: 58.0035 E: 68.7474
project/cruise: Arabian Sea/TTN-045; Process cruise 2 (Spring Intermonsoon)

ship: R/V Thomas Thompson

Data Processing Description

HPLC Pigment methods

Method by Wright et al (Mar. Ecol. Prog. Ser. 1991, 77:183-196) CHLA1, CHLA2, CHLB1 and CHLB2 estimated following the method of Latasa et al (Mar. Chem. 1996, 51:315-324)

Pigment data for P2 & P5:

A comparison of the TURNER-determined chlorophyll a concentrations with the HPLC-determined TOTCHLA concentrations (monovinyl chlorophyll a + divinyl chlorophyll a + monovinyl chlorophyllide a; units = ng Chl a equivalents/L) was performed for Process Cruise #2 (TTN-045) and Process Cruise #5 (TTN-050). While good correlations were obtained for both cruises, the slope obtained for Process Cruise #5 was significantly different from 1 (i.e., TURNER > HPLC). This difference was probably caused by the presence of Chl a-related pigments during Process Cruise #5. Thus, we recommend that whenever possible use the HPLC pigment data and not the TURNER pigment data. If HPLC data is not available for a given cast, we further recommend that you use the following equations to transform the TURNER data into HPLC-equivalent concentrations (cf., Babin, M., A. Morel, H. Claustre, A. Bricaud, Z. Kolber and P.G. Falkowski. 1996. Nitrogenand irradiance-dependent variations of the maximum quantum yield of carbon fixation in eutrophic, mesotrophic and oligotrophic marine systems. Deep-Sea Research, in press).

Results of geometric mean regression analyses (reduced major axis):

Y = HPLC TOTCHLA (monovinyl chlorophyll a + divinyl chlorophyll a + monovinyl chlorophyllide a), units = ng Chl a equivalents/L

X = TURNER chlorophyll a (it is necessary to convert the Turner Chl a concentrations in the Arabian Sea data base from mg/m3 to ng/L by multiplying concentrations by 1000)

(1) Process Cruise #2 (TTN-045)

HPLC TOTCHLA = TURNER*(0.975) + 4.833 (r = 0.9822, n = 146)

(2) Process Cruise #5 (TTN-050)

Robert R. Bidigare Department of Oceanography University of Hawaii Honolulu, HI 96822 808-956-6567 (voice mail) 808-956-9516 (fax)

[table of contents | back to top]

Data Files

File

HPLC_pigmentsTM.csv(Comma Separated Values (.csv), 12.73 KB) MD5::6945210a6868147dbe444937171083f

Primary data file for dataset ID 2549

[table of contents | back to top]

Parameters

Parameter	Description	Units
event	event number, from event log	
sta	station number, from event log	
sta_std	Arabian Sea standard station identifier	
cast	cast number, from event log	
bot	rosette bottle number	
depth_n	nominal sample depth	meters
chlide_a	Chlorophyllide a	nanogram/liter
chl_c3	Chlorophyll c3	nanogram/liter
chl_c	Chlorophyll c1 + chlorophyll c2 + Mg 3,8 divinyl pheoporphyrin a5	nanogram/liter
peridinin	Peridinin	nanogram/liter
fucox_but	19'-Butanoyloxyfucoxanthin	nanogram/liter
fucox	Fucoxanthin	nanogram/liter
fucox_hex	19'-Hexanoyloxyfucoxanthin	nanogram/liter
cis_fucox	Cis-fucoxanthin	nanogram/liter
cis_hex	Cis-19'-hexanoyloxyfucoxanthin	nanogram/liter
prasinox	Prasinoxanthin	nanogram/liter
violax	Violaxanthin	nanogram/liter
diadinox	Diadinoxanthin	nanogram/liter
allox	Alloxanthin	nanogram/liter
diatox	Diatoxanthin	nanogram/liter
lutein	Lutein	nanogram/liter
zeax	Zeaxanthin	nanogram/liter
carotene_a	alpha-carotene	nanogram/liter
carotene_b	beta-carotene	nanogram/liter
chl_b2	Divinyl chlorophyll b	nanogram/liter
chl_b1	Monovinyl chlorophyll b	nanogram/liter
chl_a2	Divinyl chlorophyll a	nanogram/liter
chl_a1	Monovinyl chlorophyll a	nanogram/liter
chl_b_tot	Divinyl chlorophyll b plus Monovinyl chlorophyll b	nanogram/liter
chl_a_tot	Divinyl chlorophyll a plus Monovinyl chlorophyll a plus chlorophyllide a	nanogram/liter

[table of contents | back to top]

Instruments

Dataset-specific Instrument Name	Trace Metal Bottle	
Generic Instrument Name	Trace Metal Bottle	
Dataset-specific Description	Trace Metal (TM) Rosette bottles	
Generic Instrument Description	Trace metal (TM) clean rosette bottle used for collecting trace metal clean seawater samples.	

[table of contents | back to top]

Deployments

TT045

Website		
Platform		
Start Date	1995-03-14	
End Date	1995-04-10	

[table of contents | back to top]

Project Information

U.S. JGOFS Arabian Sea (Arabian Sea)

Website: http://usigofs.whoi.edu/research/arabian.html

Coverage: Arabian Sea

The U.S. Arabian Sea Expedition which began in September 1994 and ended in January 1996, had three major components: a U.S. JGOFS Process Study, supported by the National Science Foundation (NSF); Forced Upper Ocean Dynamics, an Office of Naval Research (ONR) initiative; and shipboard and aircraft measurements supported by the National Aeronautics and Space Administration (NASA). The Expedition consisted of 17 cruises aboard the R/V Thomas Thompson, year-long moored deployments of five instrumented surface buoys and five sediment-trap arrays, aircraft overflights and satellite observations. Of the seventeen ship cruises, six were allocated to repeat process survey cruises, four to SeaSoar mapping cruises, six to mooring and benthic work, and a single calibration cruise which was essentially conducted in transit to the Arabian Sea.

[table of contents | back to top]

Program Information

U.S. Joint Global Ocean Flux Study (U.S. JGOFS)

Website: http://usjgofs.whoi.edu/

Coverage: Global

The United States Joint Global Ocean Flux Study was a national component of international JGOFS and an integral part of global climate change research.

The U.S. launched the Joint Global Ocean Flux Study (JGOFS) in the late 1980s to study the ocean carbon cycle. An ambitious goal was set to understand the controls on the concentrations and fluxes of carbon and associated nutrients in the ocean. A new field of ocean biogeochemistry emerged with an emphasis on quality measurements of carbon system parameters and interdisciplinary field studies of the biological, chemical and physical process which control the ocean carbon cycle. As we studied ocean biogeochemistry, we learned that our simple views of carbon uptake and transport were severely limited, and a new "wave" of ocean science was born. U.S. JGOFS has been supported primarily by the U.S. National Science Foundation in collaboration with the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration, the Department of Energy and the Office of Naval Research. U.S. JGOFS, ended in 2005 with the conclusion of the Synthesis and Modeling Project (SMP).

[table of contents | back to top]

Funding

Funding Source	Award	
National Science Foundation (NSF)	unknown Arabian Sea NSF	

[table of contents | back to top]