

# Gross and integrated gross oxygen productivity from R/V Atlantis II cruise AII-119-4 in the North Atlantic in 1989 (U.S. JGOFS NABE project)

Website: <https://www.bco-dmo.org/dataset/2576>

Version: June 15, 1995

Version Date: 1995-06-15

## Project

» [U.S. JGOFS North Atlantic Bloom Experiment](#) (NABE)

## Program

» [U.S. Joint Global Ocean Flux Study](#) (U.S. JGOFS)

Contributors	Affiliation	Role
<a href="#">Bender, Michael L.</a>	University of Rhode Island (URI-GSO)	Principal Investigator
<a href="#">Kiddon, J.</a>	University of Rhode Island (URI-GSO)	Co-Principal Investigator
<a href="#">Chandler, Cynthia L.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Dataset Description

Gross and Integrated Gross Oxygen Productivity

## Methods & Sampling

**PI:** Michael Bender and J. Kiddon  
**of:** University of Rhode Island  
**dataset:** Gross and Integrated Gross Oxygen Productivity  
**dates:** April 29, 1989 to May 7, 1989  
**location:** N: 46.8 S: 46.4 W: -20.2 E: -19  
**project/cruise:** North Atlantic Bloom Experiment/Atlantis II 119, leg 4  
**ship:** Atlantis II

sta=station number, each day cycles a new station number  
gross\_integ\_prod=Integrated O-18 Gross O2 Production,  
units=mmoles o2/m2/day  
depth=sample depth, units=meters  
gross\_o2=O-18 Gross O2 Production, units=umoles O2/L/14hr

## Methodology: J. Kiddon, M.L. Bender; URI O-18 Gross O2 Production

units: umoles O2/L/14 hr

## North Atlantic Bloom Experiment

## Atlantis II Cruise 119 Leg 4

Seawater samples were spiked with H<sub>2</sub>O enriched in O-18, such that the 18/16 oxygen mass ratio in the sample water was about twice the natural level. Gross production during a 14 hour bottle incubation generates O<sub>2</sub> with the same enriched ratio, thereby enhancing the 18/16 ratio in the large original dissolved O<sub>2</sub> pool. The tagged O<sub>2</sub> mixes with the large O<sub>2</sub> pool before being respired, thereby minimizing changes in the isotopic composition of the measured O<sub>2</sub> pool associated with respiration. Thus, the O-18 enrichment of the dissolved O<sub>2</sub> is taken to be a proportional measure of gross production.

Water samples were drawn before dawn, spiked with 0.2 ml of H<sub>2</sub>O(18) and incubated in 100 ml quartz bottles for 14 daylight hrs at the depth of collection. A drifting buoy was used for sample deployment. The incubated samples, as well as unincubated samples from each depth were processed to strip and collect dissolved gases. This stripping process was accomplished by introducing approximately 50 ml. of seawater into an evacuated two chamber container; the degassed water remained in a lower chamber and the stripped gases rose to an upper glass ampoule. The ampoule was flame sealed and returned to the lab where the 18/16 ratio of the dissolved O<sub>2</sub> was measured with an isotope ratio mass spectrometer as the per mil difference relative to a laboratory standard (referred to as 'del' measurements).

The gross O<sub>2</sub> production, denoted [O<sub>2</sub>]<sub>p</sub>, was calculated as:  $[O_2]_p = \frac{(\delta f - (\delta i)/[(\delta p - (\delta f))]}{(\delta i)} * [O_2]_i$  (eqn 1) The parameters (δ)<sub>i</sub> and (δ)<sub>f</sub> are the 'del' values of the dissolved O<sub>2</sub> measured respectively before and after incubation. (δ)<sub>p</sub> is the isotopic composition of the O<sub>2</sub> produced during the incubation, calculated knowing the volumes (V) of the H<sub>2</sub>O(18) spike and the sample, the mole fraction of O-18 in the spike (0.980) and the mole fraction of O-18 in natural seawater (0.002). That is, (δ)<sub>p</sub> =  $\frac{[(X)_{incub}/(X)_{ref}] - 1}{1000}$ , where (X)<sub>incub</sub> is the mole fraction of O-18 in the spiked sample water, itself calculated as:  $(V_{spike}/V_{bottle}) * 0.978 + 0.002$ ; and (X)<sub>ref</sub> is the separately determined O-18 mole fraction in the laboratory standard. [O<sub>2</sub>]<sub>i</sub> in equation 1 is the initial O<sub>2</sub> concentration, determined via Winkler titration by the Oceanographic Data facility. Equation 1 may be derived from a more intuitive equation which expresses the final isotopic composition (δ)<sub>f</sub> as a weighted average of the isotopic compositions of the initial O<sub>2</sub> and the O<sub>2</sub> added during production:  $(\delta f) = \frac{[O_2]_i * (\delta i) + [O_2]_p * (\delta p)}{[O_2]_i + [O_2]_p}$ .

## Integrated gross O<sub>2</sub> production

**units: mmoles O<sub>2</sub>/m<sup>2</sup>/day**

Leg 4

Integrated values of productivity were calculated using the histogram method. The euphotic region (surface to the 1% light level, Knudson et al.) was divided into intervals of uniform productivity associated with O-18 gross production measurements. The summation interval was defined as the depth interval bounded by the two mid points of three adjacent sampling depths. For example, if depths 4, 12, 20 and 30 meters were sampled, the productivity measured at 20m was taken to represent the interval 16 to 25 meters. The productivity of the shallowest sample represents the interval

from the surface to the mid point with the next deepest sample, i.e., from 0 to 8 meters. The deepest sample represents the interval from the last mid point to the 1% light level.

## **Net O<sub>2</sub> Production**

**units: umoles O<sub>2</sub>/L/24 hr**

### **Leg 4**

Net O<sub>2</sub> production was determined as the difference in the measured dissolved O<sub>2</sub> concentrations of sea water, measured before and after a 24 hour light/dark incubation by Winkler titrations. High precision in the Winkler determinations, +/- 0.1% umole O<sub>2</sub>/L, was achieved both by using an automated titrator (Radiometer, model ABU93) and by averaging four replicate measurements for each water sample.

Eight replicate water samples were drawn into quartz bottles from a Go-Flo flask containing water collected from the euphotic region before dawn (sample volumes about 100 ml, known to 0.01ml). Winkler titrations were performed on four of the replicates immediately, and the results averaged to establish the initial O<sub>2</sub> concentration. The remaining four samples were incubated for 14 daylight hours at the depth of collection (attached to a drifting buoy), and further incubated for 10 night hours in a darkened, ship board incubator, maintained at constant temperature by flowing surface water. Winkler titrations were then performed on the incubated samples and the results averaged. Net production was calculated as the difference between the final and initial O<sub>2</sub> concentrations.

## **Integrated net O<sub>2</sub> production**

**units: mmoles O<sub>2</sub>/m<sup>2</sup>/day**

### **Leg 4**

Integrated values of net O<sub>2</sub> productivity were calculated using the histogram method. The summation intervals were defined as described above for the integrated gross production, e.g., with boundaries set midway between sampled depths. Productivity was integrated over the euphotic region, to the 1% light level (Knudson et al.). In cases where data was sparse (stations 19 and 20 and, in general, for depths near the 1% light level), the net productivity was augmented using computed values of respiration rates and extrapolated gross O<sub>2</sub> productivities: net prod = gross prod - 24 \* resp rate.

Reference:

Knudson, C. W.S. Chamberlin and J. Marra (1989)

Primary production and irradiance data for U.S. JGOFS (Leg 4) Atlantis II (Cruise 112.4) Technical report LDGO-89.4. Lamont-Doherty Geological Observatory, Palisades, N.Y.

[ [table of contents](#) | [back to top](#) ]

---

## **Data Files**

<b>File</b>
<b>ox18.csv</b> (Comma Separated Values (.csv), 2.03 KB) MD5:3af8367a934fb56de68d68d28e99f52d
Primary data file for dataset ID 2576

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
year	year, reported as YYYY	YYYY
date	date, reported as MMDD	MMDD
sta	station number, from event log	dimensionless
lat	latitude, minus = south	decimal degrees
lon	longitude, minus = west	decimal degrees
gross_integ_prod	integrated O-18 gross O2 production to the 1% light level	millimoles O2/m2/day
depth	sample depth	meters
gross_o2	O-18 gross O2 production	micromoles O2/liter/14hours

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Drifter Buoy
<b>Generic Instrument Name</b>	Drifter Buoy
<b>Dataset-specific Description</b>	Used to obtain samples.
<b>Generic Instrument Description</b>	<p>Drifting buoys are free drifting platforms with a float or buoy that keep the drifter at the surface and underwater sails or socks that catch the current. These instruments sit at the surface of the ocean and are transported via near-surface ocean currents. They are not fixed to the ocean bottom, therefore they "drift" with the currents. For this reason, these instruments are referred to as drifters, or drifting buoys. The surface float contains sensors that measure different parameters, such as sea surface temperature, barometric pressure, salinity, wave height, etc. Data collected from these sensors are transmitted to satellites passing overhead, which are then relayed to land-based data centers. definition sources:</p> <p><a href="https://mmisw.org/ont/ioos/platform/drifting_buoy">https://mmisw.org/ont/ioos/platform/drifting_buoy</a> and <a href="https://www.aoml.noaa.gov/phod/gdp/faq.php#drifter1">https://www.aoml.noaa.gov/phod/gdp/faq.php#drifter1</a></p>

<b>Dataset-specific Instrument Name</b>	GO-FLO Bottle
<b>Generic Instrument Name</b>	GO-FLO Bottle
<b>Dataset-specific Description</b>	samples were drawn from GO-FLO bottles, fixed and then redeployed on a drifter buoy for incubation
<b>Generic Instrument Description</b>	GO-FLO bottle cast used to collect water samples for pigment, nutrient, plankton, etc. The GO-FLO sampling bottle is specially designed to avoid sample contamination at the surface, internal spring contamination, loss of sample on deck (internal seals), and exchange of water from different depths.

<b>Dataset-specific Instrument Name</b>	Isotope-ratio Mass Spectrometer
<b>Generic Instrument Name</b>	Isotope-ratio Mass Spectrometer
<b>Dataset-specific Description</b>	An isotope ratio mass spectrometer was used to measure the 18/16 ratio of the dissolved O <sub>2</sub> .
<b>Generic Instrument Description</b>	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

<b>Dataset-specific Instrument Name</b>	Quartz Bottle
<b>Generic Instrument Name</b>	Light-Dark Bottle
<b>Dataset-specific Description</b>	100 ml quartz bottles were used to store samples which spiked with 0.2 ml of H <sub>2</sub> O(18) for 14 daylight hrs at the depth of collection.
<b>Generic Instrument Description</b>	The light/dark bottle is a way of measuring primary production by comparing before and after concentrations of dissolved oxygen. Bottles containing seawater samples with phytoplankton are incubated for a predetermined period of time under light and dark conditions. Incubation is preferably carried out in situ, at the depth from which the samples were collected. Alternatively, the light and dark bottles are incubated in a water trough on deck, and neutral density filters are used to approximate the light conditions at the collection depth. Rates of net and gross photosynthesis and respiration can be determined from measurements of dissolved oxygen concentration in the sample bottles.

<b>Dataset-specific Instrument Name</b>	Winkler Oxygen Titrator
<b>Generic Instrument Name</b>	Winkler Oxygen Titrator
<b>Generic Instrument Description</b>	A Winkler Oxygen Titration system is used for determining concentration of dissolved oxygen in seawater.

## Deployments

### All-119-4

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57737">https://www.bco-dmo.org/deployment/57737</a>
<b>Platform</b>	R/V Atlantis II
<b>Start Date</b>	1989-04-17
<b>End Date</b>	1989-05-11
<b>Description</b>	early bloom cruise; 17 locations; 60N 21W to 46N 18W

## Project Information

### U.S. JGOFS North Atlantic Bloom Experiment (NABE)

**Website:** <http://usjgofs.whoi.edu/research/nabe.html>

**Coverage:** North Atlantic

One of the first major activities of JGOFS was a multinational pilot project, North Atlantic Bloom Experiment (NABE), carried out along longitude 20° West in 1989 through 1991. The United States participated in 1989 only, with the April deployment of two sediment trap arrays at 48° and 34° North. Three process-oriented cruises were conducted, April through July 1989, from R/V *Atlantis II* and R/V *Endeavor* focusing on sites at 46° and 59° North. Coordination of the NABE process-study cruises was supported by NSF-OCE award # 8814229. Ancillary sea surface mapping and AXBT profiling data were collected from NASA's P3 aircraft for a series of one day flights, April through June 1989.

A detailed description of NABE and the initial synthesis of the complete program data collection efforts appear in: Topical Studies in Oceanography, JGOFS: The North Atlantic Bloom Experiment (1993), Deep-Sea Research II, Volume 40 No. 1/2.

The U.S. JGOFS Data management office compiled a preliminary NABE data report of U.S. activities: Slagle, R. and G. Heimerdinger, 1991. U.S. Joint Global Ocean Flux Study, North Atlantic Bloom Experiment, Process Study Data Report P-1, April-July 1989. NODC/U.S. JGOFS Data Management Office, Woods Hole Oceanographic Institution, 315 pp. (out of print).

## Program Information

### U.S. Joint Global Ocean Flux Study (U.S. JGOFS)

**Website:** <http://usjgofs.whoi.edu/>

**Coverage:** Global

The United States Joint Global Ocean Flux Study was a national component of international JGOFS and an integral part of global climate change research.

The U.S. launched the Joint Global Ocean Flux Study (JGOFS) in the late 1980s to study the ocean carbon cycle. An ambitious goal was set to understand the controls on the concentrations and fluxes of carbon and associated nutrients in the ocean. A new field of ocean biogeochemistry emerged with an emphasis on quality measurements of carbon system parameters and interdisciplinary field studies of the biological, chemical and physical process which control the ocean carbon cycle. As we studied ocean biogeochemistry, we learned that our simple views of carbon uptake and transport were severely limited, and a new "wave" of ocean science was born. U.S. JGOFS has been supported primarily by the U.S. National Science Foundation in collaboration with the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration, the Department of Energy and the Office of Naval Research. U.S. JGOFS, ended in 2005 with the conclusion of the Synthesis and Modeling Project (SMP).

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
National Science Foundation (NSF)	<a href="#">unknown NABE NSF</a>

[ [table of contents](#) | [back to top](#) ]