

# Mesozooplankton biomass from R/V Atlantis II cruises All-119-4, All-119-5 in the North Atlantic in 1989 (U.S. JGOFS NABE project)

**Website:** <https://www.bco-dmo.org/dataset/2583>

**Version:** December 5, 1994

**Version Date:** 1994-12-05

## Project

» [U.S. JGOFS North Atlantic Bloom Experiment](#) (NABE)

## Program

» [U.S. Joint Global Ocean Flux Study](#) (U.S. JGOFS)

Contributors	Affiliation	Role
<a href="#">Dam, Hans G.</a>	University of Connecticut (UConn)	Principal Investigator
<a href="#">Chandler, Cynthia L.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Dataset Description

Mesozooplankton biomass

## Methods & Sampling

**PI:** Hans Dam  
**of:** University of Connecticut  
**dataset:** Mesozooplankton biomass  
**dates:** April 25, 1989 to May 31, 1989  
**location:** N: 47.0603 S: 46.255 W: -20.2152 E: -17.6567  
**project/cruise:** North Atlantic Bloom Experiment/Atlantis II 119, leg 4 and 5  
**ship:** R/V Atlantis II

### Methodology

See: Dam, Hans G., Carolyn Miller and Sigrun Jonasdottir, 1993. The trophic role of mesozooplankton at 47N and 20W during the North Atlantic Bloom Experiment. Deep-Sea Research II, Vol. 40, NO. 12 pp 197-212.

### DMO Note:

The reporting of plankton tow events was inconsistent, event numbers may reference a single tow or a series of tows. Use caution in a merging of data from different sources.

## Data Files

File
<b>zoomass.csv</b> (Comma Separated Values (.csv), 4.02 KB) MD5:44ee060c50a9786771ff24d3e1489ca7
Primary data file for dataset ID 2583

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
year	year (as YYYY)	dimensionless
event	event number, from event log	dimensionless
sta	station number from event log	dimensionless
cast	cast/operation number, numbered consecutively within station	dimensionless
depth_tow	depth of plankton tow	meters
aliquot	aliquot value, the reciprocal (1/value) of the fraction of the sample volume employed for the CHN measurement	1/value
vol_filt	volume of water filtered by the net	cubic meters
sizefrac	size fraction/interval of organisms analyzed	millimeters
C	carbon content	micrograms
N	nitrogen content	micrograms
C_mass	carbon biomass values are the means of two pseudoreplicates (subsamples from one tow)	milligrams C/cubic meter
N_mass	nitrogen biomass	milligrams N/cubic meter
C_to_N	carbon/nitrogen ratio	

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Mechanical Flowmeter
<b>Generic Instrument Name</b>	Mechanical Flowmeter
<b>Dataset-specific Description</b>	A General Oceanics flowmeter mounted at the mouth of the net and a filtering cod end (mesh size 0.2 mm).
<b>Generic Instrument Description</b>	Manufactured by General Oceanics, a mechanical flow meter is used with plankton tows to determine the volume of water which flows through the net. Flow meters are also used in rivers, estuaries, canals, sewer outfalls, pipes, and harbor entrances to determine water velocity and flow distance information.

<b>Dataset-specific Instrument Name</b>	RingNet
<b>Generic Instrument Name</b>	Ring Net
<b>Dataset-specific Description</b>	a 1 meter ring net fitted with a 0.2 mm mesh
<b>Generic Instrument Description</b>	A Ring Net is a generic plankton net, made by attaching a net of any mesh size to a metal ring of any diameter. There are 1 meter, .75 meter, .25 meter and .5 meter nets that are used regularly. The most common zooplankton ring net is 1 meter in diameter and of mesh size .333mm, also known as a 'meter net' (see Meter Net).

[ [table of contents](#) | [back to top](#) ]

## Deployments

### All-119-5

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57738">https://www.bco-dmo.org/deployment/57738</a>
<b>Platform</b>	R/V Atlantis II
<b>Start Date</b>	1989-05-15
<b>End Date</b>	1989-06-06
<b>Description</b>	late bloom cruise; 31 locations; 61N 22W to 41N 17W  <b>Methods &amp; Sampling</b> PI: Hans Dam of: University of Connecticut dataset: Mesozooplankton biomass dates: April 25, 1989 to May 31, 1989 location: N: 47.0603 S: 46.255 W: -20.2152 E: -17.6567 project/cruise: North Atlantic Bloom Experiment/Atlantis II 119, leg 5 ship: R/V Atlantis II Methodology See: Dam, Hans G., Carolyn Miller and Sigrun Jonasdottir, 1993. The trophic role of mesozooplankton at 47N and 20W during the North Atlantic Bloom Experiment. Deep-Sea Research II, Vol. 40, NO. 12 pp 197-212. DMO Note: The reporting of plankton tow events was inconsistent, event numbers may reference a single tow or a series of tows. Use caution in a merging of data from different sources.

### All-119-4

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57737">https://www.bco-dmo.org/deployment/57737</a>
<b>Platform</b>	R/V Atlantis II
<b>Start Date</b>	1989-04-17
<b>End Date</b>	1989-05-11
<b>Description</b>	early bloom cruise; 17 locations; 60N 21W to 46N 18W  <b>Methods &amp; Sampling</b> PI: Hans Dam of: University of Connecticut dataset: Mesozooplankton biomass dates: April 25, 1989 to May 31, 1989 location: N: 47.0603 S: 46.255 W: -20.2152 E: -17.6567 project/cruise: North Atlantic Bloom Experiment/Atlantis II 119, leg 4 ship: R/V Atlantis II Methodology See: Dam, Hans G., Carolyn Miller and Sigrun Jonasdottir, 1993. The trophic role of mesozooplankton at 47N and 20W during the North Atlantic Bloom Experiment. Deep-Sea Research II, Vol. 40, NO. 12 pp 197-212. DMO Note: The reporting of plankton tow events was inconsistent, event numbers may reference a single tow or a series of tows. Use caution in a merging of data from different sources.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### U.S. JGOFS North Atlantic Bloom Experiment (NABE)

**Website:** <http://usjgofs.whoi.edu/research/nabe.html>

**Coverage:** North Atlantic

One of the first major activities of JGOFS was a multinational pilot project, North Atlantic Bloom Experiment (NABE), carried out along longitude 20° West in 1989 through 1991. The United States participated in 1989 only, with the April deployment of two sediment trap arrays at 48° and 34° North. Three process-oriented cruises were conducted, April through July 1989, from R/V *Atlantis II* and R/V *Endeavor* focusing on sites at 46° and 59° North. Coordination of the NABE process-study cruises was supported by NSF-OCE award # 8814229. Ancillary sea surface mapping and AXBT profiling data were collected from NASA's P3 aircraft for a series of one day flights, April through June 1989.

A detailed description of NABE and the initial synthesis of the complete program data collection efforts appear in: Topical Studies in Oceanography, JGOFS: The North Atlantic Bloom Experiment (1993), Deep-Sea Research II, Volume 40 No. 1/2.

The U.S. JGOFS Data management office compiled a preliminary NABE data report of U.S. activities: Slagle, R. and G. Heimerdinger, 1991. U.S. Joint Global Ocean Flux Study, North Atlantic Bloom Experiment, Process Study Data Report P-1, April-July 1989. NODC/U.S. JGOFS Data Management Office, Woods Hole Oceanographic Institution, 315 pp. (out of print).

[ [table of contents](#) | [back to top](#) ]

---

## Program Information

### U.S. Joint Global Ocean Flux Study (U.S. JGOFS)

**Website:** <http://usjgofs.whoi.edu/>

**Coverage:** Global

The United States Joint Global Ocean Flux Study was a national component of international JGOFS and an integral part of global climate change research.

The U.S. launched the Joint Global Ocean Flux Study (JGOFS) in the late 1980s to study the ocean carbon cycle. An ambitious goal was set to understand the controls on the concentrations and fluxes of carbon and associated nutrients in the ocean. A new field of ocean biogeochemistry emerged with an emphasis on quality measurements of carbon system parameters and interdisciplinary field studies of the biological, chemical and physical process which control the ocean carbon cycle. As we studied ocean biogeochemistry, we learned that our simple views of carbon uptake and transport were severely limited, and a new "wave" of ocean science was born. U.S. JGOFS has been supported primarily by the U.S. National Science Foundation in collaboration with the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration, the Department of Energy and the Office of Naval Research. U.S. JGOFS, ended in 2005 with the conclusion of the Synthesis and Modeling Project (SMP).

[ [table of contents](#) | [back to top](#) ]

---

---

## Funding

Funding Source	Award
National Science Foundation (NSF)	<a href="#">unknown NABE NSF</a>

[ [table of contents](#) | [back to top](#) ]