Fluorometric chlorophyll-a from R/V Roger Revelle KIWI6, KIWI7, KIWI8, KIWI9 cruises in the Southern Ocean, 1998 (U.S. JGOFS AESOPS project)

Website: https://www.bco-dmo.org/dataset/2776

Version: August 23, 2001 Version Date: 2001-08-23

Project

» U.S. JGOFS Antarctic Environment and Southern Ocean Process Study (AESOPS)

Program

» <u>U.S. Joint Global Ocean Flux Study</u> (U.S. JGOFS)

Contributors	Affiliation	Role
Smith, Walker O.	Virginia Institute of Marine Science (VIMS)	Principal Investigator
Chandler, Cynthia L.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- <u>Dataset Description</u>
- Parameters
- Instruments
- <u>Deployments</u>
- Project Information
- <u>Program Information</u>

Dataset Description

Fluorometric chlorophyll-a

[table of contents | back to top]

Parameters

Parameter	Description	Units
event	event number from event log	
sta	station number from event log	
cast	cast number	
cast_type	CTD = CTD rosette TM = trace metal rosette	
bot	rosette bottle number	
depth_n	nominal sample depth	meters
chl_a_fluor	chlorophyll-a, fluorometric method	micrograms/liter

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	Niskin Bottle
Generic Instrument Name	Niskin bottle
Dataset- specific Description	CTD clean rosette (Niskin) bottles were used to collect water samples.
	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset-specific Instrument Name	Trace Metal Bottle
Generic Instrument Name	Trace Metal Bottle
Dataset-specific Description	Trace metal (TM) clean rosette bottles were used to collect water samples.
Generic Instrument Description	Trace metal (TM) clean rosette bottle used for collecting trace metal clean seawater samples.

[table of contents | back to top]

Deployments

KIW16

Website	https://www.bco-dmo.org/deployment/57724
Platform	R/V Roger Revelle
Report	http://usjgofs.whoi.edu/aesops/RRs1.html
Start Date	1997-10-20
End Date	1997-11-24
Description	Polar Front Survey I Methods & Sampling PI: Walker O. Smith of: University of Tennessee dataset: Fluorometric chlorophyll-a dates: October 23, 1997 to November 18, 1997 location: N: -56.9998 S: -62.3658 W: -171.9 E: - 168.0622 project/cruise: AESOPS KIWI6, APFZ Polar Front Survey cruise 1 ship: Roger Revelle Methodology: Please see Chapter 14 of the JGOFS protocols (1994), "Measurement of Chlorophyll a and Phaeopigments by Fluorometric Analysis" PI-Notes on Methodology: 25mm GF/F filters Extracted volume = 7mL in 90% Acetone 1.2M HCl = 3 drops Turner Design Analog Field Fluorometer. Blank=0. Calibrated by: S. Polk

KIW17

Website	https://www.bco-dmo.org/deployment/57725
Platform	R/V Roger Revelle
Report	http://usjgofs.whoi.edu/aesops/RRp1.html
Start Date	1997-12-02
End Date	1998-01-03
Description	Polar Front Process I Methods & Sampling PI: Walker O. Smith of: University of Tennessee dataset: Fluorometric chlorophyll-a dates: December 04, 1997 to December 30, 1997 location: N: -52.9388 S: -64.7418 W: -174.7303 E: -168.8302 project/cruise: AESOPS KIWI7, APFZ Polar Front Process cruise 1 ship: Roger Revelle Methodology: Please see Chapter 14 of the JGOFS protocols (1994), "Measurement of Chlorophyll a and Phaeopigments by Fluorometric Analysis" PI-Notes on Methodology: 25mm GF/F filters Extracted volume = 7mL in 90% Acetone 1.2M HCl = 3 drops Turner Design Analog Field Fluorometer. Blank=0. Calibrated by: J.Postal

KIW18

IZI VV I O	
Website	https://www.bco-dmo.org/deployment/57726
Platform	R/V Roger Revelle
Report	http://usjgofs.whoi.edu/aesops/RRs2.html
Start Date	1998-01-08
End Date	1998-02-08
Description	Polar Front Survey II Methods & Sampling PI: Walker O. Smith of: University of Tennessee dataset: Fluorometric chlorophyll-a dates: January 10, 1998 to January 28, 1998 location: N: -53.0002 S: -67.7842 W: -175.5483 E: - 169.9983 project/cruise: AESOPS KIWI8, APFZ Polar Front Survey cruise 2 ship: Roger Revelle Methodology: Please see Chapter 14 of the JGOFS protocols (1994), "Measurement of Chlorophyll a and Phaeopigments by Fluorometric Analysis" PI-Notes on Methodology: 25mm GF/F filters Extracted volume = 7mL in 90% Acetone 1.2M HCl = 3 drops Turner Design Analog Field Fluorometer. Blank=0. Calibrated by: Sarah H.

KIW19

Website	https://www.bco-dmo.org/deployment/57727
Platform	R/V Roger Revelle
Report	http://usjgofs.whoi.edu/aesops/RRp2.html
Start Date	1998-02-13
End Date	1998-03-19
Description	Polar Front Process II Methods & Sampling PI: Walker O. Smith of: University of Tennessee dataset: Fluorometric chlorophyll-a dates: February 15, 1998 to March 15, 1998 location: N: -52.966 S: -71.3157 W: -174.7338 E: - 165.9145 project/cruise: AESOPS/KIWI09, APFZ Polar Front Process cruise 2 ship: Roger Revelle Methodology: Please see Chapter 14 of the JGOFS protocols (1994), "Measurement of Chlorophyll a and Phaeopigments by Fluorometric Analysis" PI-Notes on Methodology: 25mm GF/F filters Extracted volume = 7mL in 90% Acetone 1.2M HCl = 3 drops Turner Design Analog Field Fluorometer. Blank=0. Calibrated by: A-M White

Project Information

U.S. JGOFS Antarctic Environment and Southern Ocean Process Study (AESOPS)

Website: http://usigofs.whoi.edu/research/aesops.html

Coverage: Southern Ocean, Ross Sea

The U.S. Southern Ocean JGOFS program, called Antarctic Environment and Southern Ocean Process Study (AESOPS), began in August 1996 and continued through March 1998. The U.S. JGOFS AESOPS program focused on two regions in the Southern Ocean: an east/west section of the Ross-Sea continental shelf along 76.5°S, and a second north/south section of the Southern Ocean spanning the Antarctic Circumpolar Current (ACC) at ~170°W (identified as the Polar Front). The science program, coordinated by Antarctic Support Associates (ASA), comprised eleven cruises using the R.V.I.B Nathaniel B. Palmer and R/V Roger Revelle as observational platforms and for deployment and recovery of instrumented moorings and sediment-trap arrays. The Ross-Sea region was occupied on six occasions and the Polar Front five times. Mapping data were obtained from SeaSoar, ADCP, and bathymetric systems. Satellite coverage was provided by the NASA SeaWiFS and the NOAA/NASA Pathfinder programs.

[table of contents | back to top]

Program Information

U.S. Joint Global Ocean Flux Study (U.S. JGOFS)

Website: http://usjgofs.whoi.edu/

Coverage: Global

The United States Joint Global Ocean Flux Study was a national component of international JGOFS and an integral part of global climate change research.

The U.S. launched the Joint Global Ocean Flux Study (JGOFS) in the late 1980s to study the ocean carbon cycle. An ambitious goal was set to understand the controls on the concentrations and fluxes of carbon and associated nutrients in the ocean. A new field of ocean biogeochemistry emerged with an emphasis on quality measurements of carbon system parameters and interdisciplinary field studies of the biological, chemical and physical process which control the ocean carbon cycle. As we studied ocean biogeochemistry, we learned that our simple views of carbon uptake and transport were severely limited, and a new "wave" of ocean science was born. U.S. JGOFS has been supported primarily by the U.S. National Science Foundation in collaboration with the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration, the Department of Energy and the Office of Naval Research. U.S. JGOFS, ended in 2005 with the conclusion of the Synthesis and Modeling Project (SMP).

[table of contents | back to top]