

Sediment trap deployment summary from R/V Tangaroa cruise 61TG_3052 in the Southern Ocean in 1999 (SOIREE project)

Website: <https://www.bco-dmo.org/dataset/2862>

Version: 20August2008

Version Date: 2008-08-20

Project

» [Southern Ocean Iron Release Experiment](#) (SOIREE)

Program

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
Nodder, Scott	New Zealand National Institute of Water and Atmospheric Research (NIWA)	Principal Investigator
Charette, Matthew A.	Woods Hole Oceanographic Institution (WHOI)	Co-Principal Investigator
Trull, Tom W.	University of Tasmania (UTAS)	Co-Principal Investigator
Waite, Anya	Centre for Water Research, University of Western Australia (CWR)	Co-Principal Investigator
Mackie, Doug	University of Otago	Contact
Gegg, Stephen R.	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

Dataset Description

SOIREE Sediment Trap - Deployments

SOIREE traps - logistics

(Nodder, Charette, Waite, Trull)

Note this T1, T0 for Sediment Traps

Day T1 = 0000 h NZST 10/02/99 + 24 hours to 0000 h NZST 11/02/99

with the nominal start of SOIREE (T0) at 0000 h NZST 10/02/99;

NZST = UTC + 12 hours

MULTI-trap dimensions:

Length (L, m): 0.58 *Trap length with baffles inserted*

L without baffles: 0.525 m

AR without baffles: 7.50

Trap volume without baffles: 2.021 litres

Inside diameter (Di, m): 0.07
Outside diameter (Do, m): 0.08
Aspect ratio (AR): 8.29
Aspect ratio with salt (AR): 7.29 *Basal brine added to a height of 1-trap diameter (7 cm)*
Collection area (A, m²): 0.00385
Trap volume (V, m³): 0.00223
Trap volume (V, litres): 2.232
Baffle length (Lb, m): 0.08
Baffle diameter (Dib, m): 0.01
Baffle aspect ratio (ARb): 5.77

Note: traps were deployed with a 1-D high density brine solution (excess 50 ppt NaCl) which would alter trap AR

Flux calculations:

Flux (mg/m²/d) = Wt sample/(trap collection area x days deployed)

Wt sample = Wt_{filtered} - Wt_{blank}

METHOD

Free-floating arrays of baffled, cylindrical MULTI-sediment traps.
Twelve traps deployed on each cross-frame, 9 back-filled with a 1-trap diameter thick (7 cm), 50ppt excess NaCl basal brine (50 g NaCl/l) and 3 with polyacrylimide gels mounted on base

For all deployments and all depths:
traps 1-3 = gel samples (Anya Waite)
traps 4-6 = POC/PN/13Corg or mass/PSi (Scott Nodder, Tom Trull)
traps 7-9 = phytopigments (Scott Nodder)

For all deployments and 100 m only: traps 10-12 = Th samples (Matt Charette)

For all deployments and 300 m only: traps 10-12 = POC/PN samples (Scott Nodder)

Refer to Nodder & Waite in DSRII

Methods & Sampling

See [SOIREE Preliminary Voyage Report](#)
See comments in dataset description as well

Data Processing Description

See [SOIREE Preliminary Voyage Report](#)
See comments in dataset description as well

BCO-DMO Processing Notes

Generated from original file SOIREE_Export_final.xls, Tab: TrapDeploy provided on the Deep-Sea Research II 48 (2001) accompanying CD-Rom

BCO-DMO Edits

- parameter names modified to conform to BCO-DMO convention
- date formatted as YYYYMMDD
- time formatted as HHMM
- Lat/Lon padded to three decimal places and '-' sign added to lat
- Blank rows in original sheet removed
- 'nd' added to blank cells

- Made column header record one line
- Filled in mostly blank cells for deeper array with data from shallower array (should be the same)
- Added 'Notes' column

[[table of contents](#) | [back to top](#)]

Data Files

File
traps_Deploy.csv (Comma Separated Values (.csv), 1.91 KB) MD5:c1be01f250f1a842959280ae18413656 Primary data file for dataset ID 2862

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Patch	Patch Location (In/Out)	text
Buoy_ID	Buoy Id	integer
SOIREE_Day	SOIREE Day of experiment Note this T1, T0 for Sediment Traps Day T1 = 0000 h NZST 10/02/99 + 24 hours to 0000 h NZST 11/02/99 with the nominal start of SOIREE (T0) at 0000 h NZST 10/02/99; NZST = UTC + 12 hours	text
Array	Sediment trap array number	integer
depth	Nominal depths only, actual depths were 10 m deeper	meters
Array_ID	Array Id - Single Letter	text
Temp_Sensors	Temperature sensor ids	integer
Date_Start_NZST	Start date of trap deployment (NZST)	YYYYMMDD
Julian_Day_Start	Start Julian day of year of trap deployment	DD
Time_Start_NZST	Start time of trap deployment (NZST)	HHMM
lat_Start	Start latitude of trap deployment negative denotes South	decimal degrees
lon_Start	Start longitude of trap deployment negative denotes West	decimal degrees
lat_End	End latitude of trap deployment negative denotes South	decimal degrees
lon_End	End longitude of trap deployment negative denotes West	decimal degrees
Date_End_NZST	End date of trap deployment (NZST)	YYYYMMDD
Time_End_NZST	End time of trap deployment (NZST)	HHMM
Julian_Day_End	End Julian day of trap deployment	DD
Distance_x	X Drift distance in meters	meters
Distance_y	Y Drift distance in meters	meters
Total_distance	Total drift distance in kilometers	kilometers
Total_time_days	Total trap deployment time in days	decimal days
Drift_rate_km_per_d	Drift rate in km/day	km/day
Drift_rate_cm_per_s	Drift rate in cm/sec	cm/sec
Notes	Notes column	text

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Sediment Trap
Generic Instrument Name	Sediment Trap
Dataset-specific Description	MULTI-trap dimensions: Length (L, m): 0.58 Trap length with baffles inserted; L without baffles: 0.525 m AR without baffles: 7.50 Trap volume without baffles: 2.021 litres Inside diameter (Di, m): 0.07 Outside diameter (Do, m): 0.08 Aspect ratio (AR): 8.29 Aspect ratio with salt (AR): 7.29 Basal brine added to a height of 1-trap diameter (7 cm) Collection area (A, m2): 0.00385 Trap volume (V, m3): 0.00223 Trap volume (V, litres): 2.232 Baffle length (Lb, m): 0.08 Baffle diameter (Dib, m): 0.01 Baffle aspect ratio (ARb): 5.77
Generic Instrument Description	Sediment traps are specially designed containers deployed in the water column for periods of time to collect particles from the water column falling toward the sea floor. In general a sediment trap has a jar at the bottom to collect the sample and a broad funnel-shaped opening at the top with baffles to keep out very large objects and help prevent the funnel from clogging. This designation is used when the specific type of sediment trap was not specified by the contributing investigator.

[[table of contents](#) | [back to top](#)]

Deployments

61TG_3052

Website	https://www.bco-dmo.org/deployment/57827
Platform	R/V Tangaroa
Report	http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf
Start Date	1999-01-31
End Date	1999-03-01
Description	Cruise to the Southern Ocean as part of the Fe Sythesis project whose aim was to maintain a coherent patch of iron-enriched seawater for the duration of SOIREE and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period.

[[table of contents](#) | [back to top](#)]

Project Information

Southern Ocean Iron Release Experiment (SOIREE)

Coverage: Southern Ocean

Project in the Southern Ocean aimed at maintaining a coherent patch of iron-enriched seawater for the duration of project and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period of the project.

The Southern Ocean Iron RElease Experiment (SOIREE), was the first in situ iron fertilization experiment performed in the polar waters of the Southern Ocean. SOIREE was an interdisciplinary study involving participants from six countries, and took place in February 1999 south of the Polar Front in the Australasian-Pacific sector of the Southern Ocean.

Approximately 3800 kg of acidified FeSO₄·7H₂O and 165 g of the tracer sulphur hexafluoride (SF₆) were added to a 65-m deep surface mixed layer over an area of ~50 km². Initially, mean dissolved iron concentrations were ~2.7 nM, but decreased to ambient levels within days, requiring subsequent additions of 1550-1750 kg of acidified FeSO₄·7H₂O on days 3, 5 and 7 of the experiment.

During the 13-day site occupation, there were iron-mediated increases in phytoplankton growth rates, with marked increases in chlorophyll a (up to 2 µg l⁻¹) and production rates (up to 1.3 gCm⁻²d⁻¹). These resulted in subsequent changes in the pelagic ecosystem structure, and in the cycling of carbon, silica and sulphur, such as a 10% drawdown of surface CO₂.

The SOIREE bloom persisted for >40 days following our departure from the site, as observed via [SeaWiFS remotely sensed observations of Ocean Colour](#).

BCO-DMO Note:

All original data and metadata provided on a CD-Rom accompanying the Deep-Sea Research II 48 (2001) volume. The CD-Rom contains the main SOIREE datasets and ancillary information including the pre-experiment 'desktop' database study for site-selection, and satellite images of the SOIREE bloom.

© 2001 Elsevier Science Ltd. All rights reserved.

Related files

[SOIREE Preliminary Voyage Report](#)

[SOIREE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

[SOIREE Cruise Track](#)

[[table of contents](#) | [back to top](#)]

Program Information

Iron Synthesis (FeSynth)

Coverage: Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFS (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO₂ budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

http://www.scor-int.org/Working_Groups/wg131.htm

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for

addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

Related file

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[[table of contents](#) | [back to top](#)]