

# Daily light levels, attenuation coefficients and critical depths from the R/V Tangaroa 61TG\_3052 cruise in the Southern Ocean during 1999 (SOIREE project)

**Website:** <https://www.bco-dmo.org/dataset/2881>

**Version:** 04Feb2010

**Version Date:** 2010-02-04

## Project

» [Southern Ocean Iron Release Experiment](#) (SOIREE)

## Program

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
<a href="#">Gall, Mark</a>	New Zealand National Institute of Water and Atmospheric Research (NIWA)	Principal Investigator
<a href="#">Boyd, Philip W.</a>	New Zealand National Institute of Water and Atmospheric Research (NIWA)	Co-Principal Investigator
<a href="#">Mackie, Doug</a>	University of Otago	Contact
<a href="#">Gegg, Stephen R.</a>	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

## Dataset Description

SOIREE Summary of daily light levels, attenuation coefficients and critical depths

## Methods & Sampling

See [SOIREE Preliminary Voyage Report](#)

For more information refer to Gall et al (2001)

"Phytoplankton processes (Part 2): Rates of primary production and factors controlling algal growth during the

Southern Ocean Iron RElease Experiment (SOIREE)".

## Data Processing Description

See [SOIREE Preliminary Voyage Report](#)

For more information refer to Gall et al (2001)  
"Phytoplankton processes (Part 2): Rates of primary  
production and factors controlling algal growth during the  
Southern Ocean Iron Release Experiment (SOIREE)".

## BCO-DMO Processing Notes

Generated from original file Light\_Summary.xls Tab: LightDataSummary  
provided on the  
Deep-Sea Research II 48 (2001) accompanying CD-Rom

## BCO-DMO Edits

- parameter names modified to conform to BCO-DMO convention
- date reformatted to YYYYMMDD
- noontime lat/lon position inserted from ships track data to make these data mapable

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>light_summary.csv</b> (Comma Separated Values (.csv), 628 bytes) MD5:8968a2a19d40a7435568b611a72abfb3 Primary data file for dataset ID 2881

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
PAR	Photosynthetically Active Radiation (PAR) -400-700 nm.	M/m <sup>2</sup> /day
date	Date (GMT)	YYYYMMDD
Kpar	Light Attenuation Estimates Calculated from the beam attenuation relationship with CTDKo.	(tbd)
Ze	Defined as the 1% Light Level (Kirk, 1994).	(tbd)
Zc	Critical Light Depth The critical light depth was calculated from the equations outlined in Nelson & Smith (1991)	(tbd)
lat	latitude, negative denotes South	decimal degrees
lon	longitude, negative denotes West	decimal degrees

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	CTD Seabird 911
<b>Generic Instrument Name</b>	CTD Sea-Bird 911
<b>Dataset-specific Description</b>	NIWA's Seabird 911plus CTD and related instrumentation
<b>Generic Instrument Description</b>	The Sea-Bird SBE 911 is a type of CTD instrument package. The SBE 911 includes the SBE 9 Underwater Unit and the SBE 11 Deck Unit (for real-time readout using conductive wire) for deployment from a vessel. The combination of the SBE 9 and SBE 11 is called a SBE 911. The SBE 9 uses Sea-Bird's standard modular temperature and conductivity sensors (SBE 3 and SBE 4). The SBE 9 CTD can be configured with auxiliary sensors to measure other parameters including dissolved oxygen, pH, turbidity, fluorescence, light (PAR), light transmission, etc.). More information from Sea-Bird Electronics.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### 61TG\_3052

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57827">https://www.bco-dmo.org/deployment/57827</a>
<b>Platform</b>	R/V Tangaroa
<b>Report</b>	<a href="http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf">http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf</a>
<b>Start Date</b>	1999-01-31
<b>End Date</b>	1999-03-01
<b>Description</b>	Cruise to the Southern Ocean as part of the Fe Sythesis project whose aim was to maintain a coherent patch of iron-enriched seawater for the duration of SOIREE and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Southern Ocean Iron Release Experiment (SOIREE)

**Coverage:** Southern Ocean

Project in the Southern Ocean aimed at maintaining a coherent patch of iron-enriched seawater for the duration of project and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period of the project.

The Southern Ocean Iron RElease Experiment (SOIREE), was the first in situ iron fertilization experiment performed in the polar waters of the Southern Ocean. SOIREE was an interdisciplinary study involving participants from six countries, and took place in February 1999 south of the Polar Front in the Australasian-Pacific sector of the Southern Ocean.

Approximately 3800 kg of acidified FeSO<sub>4</sub>·7H<sub>2</sub>O and 165 g of the tracer sulphur hexafluoride (SF<sub>6</sub>) were added to a 65-m deep surface mixed layer over an area of ~50 km<sup>2</sup>. Initially, mean dissolved iron concentrations were ~2.7 nM, but decreased to ambient levels within days, requiring subsequent additions of 1550-1750 kg of acidified FeSO<sub>4</sub>·7H<sub>2</sub>O on days 3, 5 and 7 of the experiment.

During the 13-day site occupation, there were iron-mediated increases in phytoplankton growth rates, with marked increases in chlorophyll a (up to 2 µg l<sup>-1</sup>) and production rates (up to 1.3 gCm<sup>-2</sup>d<sup>-1</sup>). These resulted in subsequent changes in the pelagic ecosystem structure, and in the cycling of carbon, silica and sulphur, such as a 10% drawdown of surface CO<sub>2</sub>.

The SOIREE bloom persisted for >40 days following our departure from the site, as observed via [SeaWiFS remotely sensed observations of Ocean Colour](#).

#### **BCO-DMO Note:**

All original data and metadata provided on a CD-Rom accompanying the Deep-Sea Research II 48 (2001) volume. The CD-Rom contains the main SOIREE datasets and ancillary information including the pre-experiment 'desktop' database study for site-selection, and satellite images of the SOIREE bloom.

© 2001 Elsevier Science Ltd. All rights reserved.

## **Related files**

[SOIREE Preliminary Voyage Report](#)

[SOIREE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

[SOIREE Cruise Track](#)

[ [table of contents](#) | [back to top](#) ]

---

## **Program Information**

### **Iron Synthesis (FeSynth)**

**Coverage:** Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFS (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO<sub>2</sub> budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

[http://www.scor-int.org/Working\\_Groups/wg131.htm](http://www.scor-int.org/Working_Groups/wg131.htm)

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for

addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata  
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations  
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations  
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets  
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

## **Related file**

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[ [table of contents](#) | [back to top](#) ]