

# Model state variables inside of fertilized patch (Var.in) from the R/V Tangaroa 61TG\_3052 cruise in the Southern Ocean during 1999 (SOIREE project)

**Website:** <https://www.bco-dmo.org/dataset/2886>

**Version:** 25August2008

**Version Date:** 2008-08-25

## Project

» [Southern Ocean Iron Release Experiment](#) (SOIREE)

## Program

» [Iron Synthesis](#) (FeSynth)

| Contributors                         | Affiliation  | Role                      |
|--------------------------------------|--|---------------------------|
| <a href="#">Lancelot, Christiane</a> | Université Libre de Bruxelles - Ecologie des Systèmes Aquatiques (ESA-ULB) | Principal Investigator    |
| <a href="#">Boyd, Philip W.</a>      | New Zealand National Institute of Water and Atmospheric Research (NIWA)    | Co-Principal Investigator |
| <a href="#">Mackie, Doug</a>         | University of Otago  | Contact                   |
| <a href="#">Gegg, Stephen R.</a>     | Woods Hole Oceanographic Institution (WHOI)                                | BCO-DMO Data Manager      |

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

## Dataset Description

SOIREE Modeling - Var in  
State variables time series (inside the patch)

### SOIREE Modeling

Updated October 2000

Authors: E. Hannon, C. Lancelot, P.W. Boyd

Contact: Prof. C. Lancelot (ESA-ULB, CP221, Campus Plaine, Bd du Triomphe, B-1050 Brussels, Belgium, [lancelot@ulb.ac.be](mailto:lancelot@ulb.ac.be))

Note: these model runs were performed with the SWAMCO model.

See details relative to model concepts, parameterization and adaptation to SOIREE in:

Lancelot, C., E. Hannon, S. Becquevort, C. Veth, and H. J. W. de Baar. 2000.

Modelling phytoplankton blooms and related carbon export production in the Southern Ocean: application to the Atlantic sector in Austral spring 1992. Deep-Sea Res., Part I, 47, 1621-1662.

Hannon, E., P. W. Boyd, M. Silviso and C. Lancelot. (this issue). Modeling the bloom evolution and carbon flows during SOIREE: Implications for future in situ iron-enrichments in the Southern Ocean. Deep-Sea Res. II.

Model results are compiled in the file "SOIREEmodel.xls"

This file includes 7 sheets:

Var.out: state variables time series (outside the patch)

Var.in: state variables time series (inside the patch)

Act.out: activities (biogenic elements fluxes) time series (outside the patch)

Act.in: activities (biogenic elements fluxes) time series (inside the patch)

BudgetC: integrated carbon budget

pCO2: surface pCO2 time series

Forcing: hydro-meteorological forcings

*Comments:*

*Sheet 1 (Var.out):*

60 days (10/02/1999 to 10/04/1999) time series of the state variables values, averaged daily and over the upper mixed layer (UML)

See state variables list in Table 1 below.

Diatoms and autotrophic nanoflagellates chlorophyll a (columns AH, AI) are estimated on the basis of the functional and structural metabolites concentration (state variables DAF and NFF), using a conversion factor of 0.5 mg chl a / mmol C

*Sheet 2 (Var.in):*

see Sheet 1

*Sheet 3 (Act.out):*

activities time series (outside the patch)

60 days (10/02/1999 to 10/04/1999) time series of the activities (biogenic elements fluxes) values, integrated over the upper mixed layer (UML) and daily averaged.

See activities list in Table 2 below.

*Sheet 4 (Act.in):*

see Sheet 3

*Sheet 5 (BudgetC):*

Carbon budget of the Upper Mixed Layer, calculated over 13 days (site occupation of SOIREE) and 60 days since the initial iron release (inside and outside the fertilized patch)

*Sheet 6 (pCO2):*

Time series of daily averaged surface water pCO2 (in situ temperature, 100% wet) (inside and outside the fertilized patch)

*Sheet 7 (Forcing):*

Time series of daily averaged wind velocity (m s<sup>-1</sup>), sea surface temperature (°C), and mixed layer depth (m).

**Table 1. SWAMCO state variables**

**activity code unit**

|  |         |  |
|--|---------|--|
| diatoms photosynthesis                 | photDA  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| diatoms respiration                    | respDA  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| diatoms growth                         | sDAF    | mmol m <sup>-2</sup> day <sup>-1</sup> |
| diatoms lysis                          | ysDA    | mmol m <sup>-2</sup> day <sup>-1</sup> |
| nanoflagellates photosynthesis         | photNF  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| nanoflagellates respiration            | respNF  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| nanoflagellates growth                 | sNFF    | mmol m <sup>-2</sup> day <sup>-1</sup> |
| nanoflagellates lysis                  | lysNF   | mmol m <sup>-2</sup> day <sup>-1</sup> |
| NO3 uptake                             | NO3upt  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| NH4 uptake by phytoplankton            | NH4upt  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| mesozooplankton grazing on diatoms     | MSgDA   | mmol m <sup>-2</sup> day <sup>-1</sup> |
| mesozooplankton grazing on microzoo    | MSgMC   | mmol m <sup>-2</sup> day <sup>-1</sup> |
| MCZ grazing on autotrophic nanofalg.   | MCgNF   | mmol m <sup>-2</sup> day <sup>-1</sup> |
| MCZ grazing on heterotrophic nanoflag. | MCgHNF  | mmol m <sup>-2</sup> day <sup>-1</sup> |
| HNF grazing on bacteria                | HNFgBAC | mmol m <sup>-2</sup> day <sup>-1</sup> |
| bacteria growth                        | muBAC   | mmol m <sup>-2</sup> day <sup>-1</sup> |
| ammonification                         | NregBAC | mmol m <sup>-2</sup> day <sup>-1</sup> |

NH4 regeneration by protozooplankton NregZOO mmol m-2 day-1  
 PO4 regeneration by protozooplankton Preg mmol m-2 day-1  
 Fe regeneration by protozooplankton Fereg  $\mu$ mol m-2 day-1  
 POC export production (110m) sed mmol m-2 day-1  
 diatoms (C) export production (110m) sedDA mmol m-2 day-1  
 diatoms (Si) export production (110m) sedBSi mmol m-2 day-1  
 sea to air CO2 flux CO2flux mmol m-2 day-1  
 DIC upwelling upwDIC mmol m-2 day-1  
 Fe upwelling upwFe  $\mu$ mol m-2 day-1  
 bacteria autolysis lysBAC mmol m-2 day-1  
 HNF autolysis lysHNF mmol m-2 day-1  
 MCZ autolysis lysMCZ mmol m-2 day-1  
 POC export production out of the UML sedUML mmol m-2 day-1

**Table 2. SWAMCO activities (fluxes of biogenic elements)**

| activity  | code | unit             |
|---|------|------------------|
| Functional and structural metabolites (diatoms)   | DAF  | mmol C m-3       |
| Monomeric substrates (diatoms)                    | DAS  | mmol C m-3       |
| Reserve material (diatoms)                        | DAR  | mmol C m-3       |
| Functional and structural metabolites (nanoflag.) | NFF  | mmol C m-3       |
| Monomeric substrates (nanoflag.)                  | NFS  | mmol C m-3       |
| Reserve material (nanoflag.)                      | NFR  | mmol C m-3       |
| Microzooplankton                                  | MCZ  | mmol C m-3       |
| Heterotrophic nanoflagellates                     | HNF  | mmol C m-3       |
| Bacteria  | BAC  | mmol C m-3       |
| monomeric substrates for bacteria (C)             | BSC  | mmol C m-3       |
| dissolved organic C (high biodeg.)                | DC1  | mmol C m-3       |
| dissolved organic C (low biodeg.)                 | DC2  | mmol C m-3       |
| particulate organic C (high biodeg.)              | PC1  | mmol C m-3       |
| particulate organic C (low biodeg.)               | PC2  | mmol C m-3       |
| monomeric substrates for bacteria (N)             | BSN  | mmol N m-3       |
| dissolved organic N (high biodeg.)                | DN1  | mmol N m-3       |
| dissolved organic N (low biodeg.)                 | DN2  | mmol N m-3       |
| particulate organic N (high biodeg.)              | PN1  | mmol N m-3       |
| particulate organic N (low biodeg.)               | PN2  | mmol N m-3       |
| dissolved organic P (high biodeg.)                | DP1  | mmol P m-3       |
| dissolved organic P (low biodeg.)                 | DP2  | mmol P m-3       |
| particulate organic P (high biodeg.)              | PP1  | mmol P m-3       |
| particulate organic P (low biodeg.)               | PP2  | mmol P m-3       |
| nitrate   | NO3  | mmol N m-3       |
| ammonium  | NH4  | mmol N m-3       |
| phosphate   | PO4  | mmol P m-3       |
| silicic acid                                      | DSi  | mmol Si m-3      |
| biogenic Si                                       | BSi  | mmol Si m-3      |
| total dissolved iron                              | DFe  | $\mu$ mol Fe m-3 |
| total dissolved inorganic carbon                  | DIC  | $\mu$ mol kg-1   |
| total alkalinity                                  | ALK  | $\mu$ mol kg-1   |

## Methods & Sampling

See [SOIRE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

## Data Processing Description

See [SOIRE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

## BCO-DMO Processing Notes

Generated from original file SOIREEmodel.xls, Tab: var\_in provided on the Deep-Sea Research II 48 (2001) accompanying CD-Rom

## BCO-DMO Edits

- parameter names modified to conform to BCO-DMO convention
- date formatted as YYYYMMDD
- number values padded to two decimal places
- deleted blank column

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

| File  |
|---|
| <b>model_varin.csv</b> (Comma Separated Values (.csv), 11.09 KB)<br>MD5:759dcf923548213eacfb1783bf4b068f<br>Primary data file for dataset ID 2886 |

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

| Parameter | Description                                       | Units      |
|-----------|---|------------|
| date      | Date (GMT)  | YYYYMMDD   |
| DAF       | Functional and structural metabolites (diatoms)   | mmol C m-3 |
| DAS       | Monomeric substrates (diatoms)                    | mmol C m-3 |
| DAR       | Reserve material (diatoms)                        | mmol C m-3 |
| NFF       | Functional and structural metabolites (nanoflag.) | mmol C m-3 |
| NFS       | Monomeric substrates (nanoflag.)                  | mmol C m-3 |
| NFR       | Reserve material (nanoflag.)                      | mmol C m-3 |
| MCZ       | Microzooplankton                                  | mmol C m-3 |

|     |                                       |            |
|-----|---------------------------------------|------------|
| HNF | Heterotrophic nanoflagellates         | mmol C m-3 |
| BAC | Bacteria                              | mmol C m-3 |
| BSC | monomeric substrates for bacteria (C) | mmol C m-3 |
| DC1 | dissolved organic C (high biodeg.)    | mmol C m-3 |
| DC2 | dissolved organic C (low biodeg.)     | mmol C m-3 |
| PC1 | particulate organic C (high biodeg.)  | mmol C m-3 |
| PC2 | particulate organic C (low biodeg.)   | mmol C m-3 |
| BSN | monomeric substrates for bacteria (N) | mmol N m-3 |
| DN1 | dissolved organic N (high biodeg.)    | mmol N m-3 |
| DN2 | dissolved organic N (low biodeg.)     | mmol N m-3 |
| PN1 | particulate organic N (high biodeg.)  | mmol N m-3 |
| PN2 | particulate organic N (low biodeg.)   | mmol N m-3 |
| DP1 | dissolved organic P (high biodeg.)    | mmol P m-3 |
| DP2 | dissolved organic P (low biodeg.)     | mmol P m-3 |
| PP1 | particulate organic P (high biodeg.)  | mmol P m-3 |
| PP2 | particulate organic P (low biodeg.)   | mmol P m-3 |
| NO3 | nitrate                               | mmol N m-3 |
| NH4 | ammonium                              | mmol N m-3 |
| PO4 | phosphate                             | mmol P m-3 |
|     |                                       |            |

|           |                                   |                 |
|-----------|-----------------------------------|-----------------|
| DSi       | silicic acid                      | mmol Si m-3     |
| BSi       | biogenic Si                       | mmol Si m-3     |
| DFe       | total dissolved iron              | micromol Fe m-3 |
| DIC       | total dissolved inorganic carbon  | micromol kg-1   |
| ALK       | total alkalinity                  | micromol kg-1   |
| Chla_diat | Chla from diatoms (?????)         | (TBD)           |
| Chla_nano | Chla from nanoflagellates (?????) | (TBD)           |
| Chla_tot  | Total Chla (?????)                | (TBD)           |

[ [table of contents](#) | [back to top](#) ]

## Deployments

### 61TG\_3052

|                    |   |
|--------------------|---|
| <b>Website</b>     | <a href="https://www.bco-dmo.org/deployment/57827">https://www.bco-dmo.org/deployment/57827</a>   |
| <b>Platform</b>    | R/V Tangaroa  |
| <b>Report</b>      | <a href="http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf">http://bcodata.whoi.edu/Fe_Synthesis/SOIREE/SOIREE_cruisereport.pdf</a>   |
| <b>Start Date</b>  | 1999-01-31  |
| <b>End Date</b>    | 1999-03-01  |
| <b>Description</b> | Cruise to the Southern Ocean as part of the Fe Sythesis project whose aim was to maintain a coherent patch of iron-enriched seawater for the duration of SOIREE and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period. |

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Southern Ocean Iron Release Experiment (SOIREE)

**Coverage:** Southern Ocean

Project in the Southern Ocean aimed at maintaining a coherent patch of iron-enriched seawater for the duration of project and to interpret any iron-mediated effects on the patch by conducting measurements and performing experiments during this period of the project.

The Southern Ocean Iron RElease Experiment (SOIREE), was the first in situ iron fertilization experiment

performed in the polar waters of the Southern Ocean. SOIREE was an interdisciplinary study involving participants from six countries, and took place in February 1999 south of the Polar Front in the Australasian-Pacific sector of the Southern Ocean.

Approximately 3800 kg of acidified  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  and 165 g of the tracer sulphur hexafluoride ( $\text{SF}_6$ ) were added to a 65-m deep surface mixed layer over an area of  $\sim 50 \text{ km}^2$ . Initially, mean dissolved iron concentrations were  $\sim 2.7 \text{ nM}$ , but decreased to ambient levels within days, requiring subsequent additions of 1550-1750 kg of acidified  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  on days 3, 5 and 7 of the experiment.

During the 13-day site occupation, there were iron-mediated increases in phytoplankton growth rates, with marked increases in chlorophyll a (up to  $2 \mu\text{g l}^{-1}$ ) and production rates (up to  $1.3 \text{ gCm}^{-2}\text{d}^{-1}$ ). These resulted in subsequent changes in the pelagic ecosystem structure, and in the cycling of carbon, silica and sulphur, such as a 10% drawdown of surface  $\text{CO}_2$ .

The SOIREE bloom persisted for  $>40$  days following our departure from the site, as observed via [SeaWiFS remotely sensed observations of Ocean Colour](#).

#### **BCO-DMO Note:**

All original data and metadata provided on a CD-Rom accompanying the Deep-Sea Research II 48 (2001) volume. The CD-Rom contains the main SOIREE datasets and ancillary information including the pre-experiment 'desktop' database study for site-selection, and satellite images of the SOIREE bloom.

© 2001 Elsevier Science Ltd. All rights reserved.

### **Related files**

[SOIREE Preliminary Voyage Report](#)

[SOIREE Introduction and Summary, Deep-Sea Research II 48 \(2001\) 2425-2438](#)

[SOIREE Cruise Track](#)

[ [table of contents](#) | [back to top](#) ]

---

## **Program Information**

### **Iron Synthesis (FeSynth)**

**Coverage:** Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFs (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry,  $\text{CO}_2$  budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

[http://www.scor-int.org/Working\\_Groups/wg131.htm](http://www.scor-int.org/Working_Groups/wg131.htm)

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata  
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations  
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations  
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets  
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

## Related file

### [Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[ [table of contents](#) | [back to top](#) ]