

Phosphorus -TDP, PO4, DOP, PP from R/V Kaiyo-Maru cruise KY0103-02 in the Northwestern Sub-Arctic Pacific in 2001 (SEEDS I project)

Website: <https://www.bco-dmo.org/dataset/2908>

Version: 03September2008

Version Date: 2008-09-03

Project

» [Subarctic-Pacific Iron Experiment for Ecosystem Dynamics Study I](#) (SEEDS I)

Program

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
Yoshimura, Takeshi	Central Research Institute of Electric Power Industry (CRIEPI)	Principal Investigator
Mackie, Doug	University of Otago	Contact
Gegg, Stephen R.	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

Dataset Description

SEEDS 2001 Phosphorus - TDP, PO4, DOP, PP

Total dissolved phosphorous, PO4, dissolved organic phosphorous (by difference) and particulate phosphorus for all stations IN at 5, 10, 20, 30, 50, 70, 100, 150, 200m.

Dissolved phosphorous, PO4, dissolved organic phosphorous (by difference) but NOT particulate phosphorus for stations OUT at 5, 10, 20, 30, 50, 70, 100, 150, 200m.

Data Processing Description

BCO-DMO Processing Notes

CSV file generated by Doug Mackie from original spreadsheet Phosphorus.xls

Notes from CSV file generated by Doug Mackie:

TDP (total dissolved phosphorus) see Hansen and Koroleff (1999) Determination of nutrients. In Methods of seawater analysis.

PO4 (phosphate) see Hansen and Koroleff (1999) Determination of nutrients. In Methods of seawater analysis.

DOP is TDP - PO4

PP (particulate phosphorus) see Solorzano and Sharp (1980) L and O 25 754-758

BCO-DMO Edits

- Parameter names modified to conform to BCO-DMO convention

[[table of contents](#) | [back to top](#)]

Data Files

File
phos.csv (Comma Separated Values (.csv), 8.94 KB) MD5:e9727d99bc22d0ae32ed7b46daef6122
Primary data file for dataset ID 2908

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
date	Date UTC	YYYYMMDD
lat	latitude, negative denotes South	decimal degrees
lon	longitude, negative denotes West	decimal degrees
station	Station Id	text
time_start	Start time of station	HHMM
time_end	End time of station	HHMM
depth	Sample depth in meters	meters
TDP	total dissolved phosphorus	uM
PO4	phosphate	uM
DOP	DOP = TDP - PO4	uM
PP	particulate phosphorus	uM

[[table of contents](#) | [back to top](#)]

Deployments

KY0103-02

Website	https://www.bco-dmo.org/deployment/57835
Platform	R/V Kaiyo-Maru
Start Date	2001-07-13
End Date	2001-08-06
Description	Patch enrichment = Leg 2: 13 Jul 2001 (Kushiro)--06 Aug 2001 (Tokyo)Note: No cruise track was contributed for this deployment. Data are plotted outside what is displayed as the "best available" cruise track from the data contributed

[[table of contents](#) | [back to top](#)]

Project Information

Subarctic-Pacific Iron Experiment for Ecosystem Dynamics Study I (SEEDS I)

Website: <http://www.seeds-exp.jp/en/index.html>

Coverage: Western subarctic gyre in the North Pacific at 48.5°N, 165°E

An in situ test of the iron limitation hypothesis in the subarctic North Pacific Ocean was performed. First experiment of two (see SEEDS 2004)

A single enrichment of dissolved iron caused a large increase in phytoplankton standing stock and decreases in macronutrients and dissolved carbon dioxide. The dominant phytoplankton species shifted after the iron addition from pennate diatoms to a centric diatom, *Chaetoceros debilis*, that showed a very high growth rate, 2.6 doublings per day. Conclusion was that the bioavailability of iron regulates the magnitude of the phytoplankton biomass and the key phytoplankton species that determine the biogeochemical sensitivity to iron supply of high-nitrate, low-chlorophyll waters.

Data was collected at a total of 13 stations and from 3 moored sediment traps.

- Stations were occupied IN patch for days 0, 2, 4, 7, 9, 11 and 13.
- Stations were occupied OUT patch for days 2, 4, 7, 9, 11, 13.

It is not explicitly stated but it appears that at all stations two CTD sampling rosette casts were made: clean and rms. The clean rosette appears to have typically sampled the mixed layer (<50 m) e.g. 5, 10, 20, 30, 50 m. The rms rosette appears to have typically sampled the euphotic zone (<200m) e.g. 10, 20, 30, 40, 50, 80, 100, 150, 200 m.

Sediment traps were deployed at:

- CENTRE: 20 m
- IN: 40, 60, 100, 200 m
- OUT: 20, 40, 60 and 100 m

Traps were recovered several times. Deployment times (days):

- CENTRE: 3.95, 2.83, 2.02, 1.98, 1.93, 2.05
- IN: 3.99, 2.84, 2.03, 2.00, 1.95, 2.01
- OUT: 5.17, 3.97, 3.42

BCO-DMO/Doug Mackie Note:

Throughout these data, stations are identified as D2-I, D2-O, etc. D2-I indicates "Day 2, in patch station". while D2-O indicates "Day 2, out patch station". This applies to all station identifiers.

Related file

[SEEDS 2001 Project Documentation](#)

Program Information

Iron Synthesis (FeSynth)

Coverage: Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFS (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO₂ budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

http://www.scor-int.org/Working_Groups/wg131.htm

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

Related file

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the

U.S. National Science Foundation (NSF).



[[table of contents](#) | [back to top](#)]