

# Nutrient concentrations and chlorophyll from bottle samples from R/V Melville cruise COOK19MV from the Southern Ocean, south of New Zealand in 2002 (SOFeX project)

**Website:** <https://www.bco-dmo.org/dataset/2944>

**Version:** 14 August 2008

**Version Date:** 2008-08-14

## Project

» [Southern Ocean Iron Experiment](#) (SOFeX)

## Programs

» [Ocean Carbon and Biogeochemistry](#) (OCB)

» [Iron Synthesis](#) (FeSynth)

Contributors	Affiliation	Role
<a href="#">Hiscock, William T.</a>	University of Miami Rosenstiel School of Marine and Atmospheric Science (UM-RSMAS)	Co-Principal Investigator
<a href="#">Millero, Frank</a>	University of Miami Rosenstiel School of Marine and Atmospheric Science (UM-RSMAS)	Co-Principal Investigator
<a href="#">Chandler, Cynthia L.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)

## Dataset Description

Nutrient concentrations and chlorophyll from bottle samples collected during CTD-Niskin and Trace-metal clean rosette casts

## Methods & Sampling

**dates:** 21 January 2002 to 21 February 2002 (20020121-20020221)

**location:** N: -52.3853 S: -66.6115 W: -175.2212 E: -166.9392

**project/cruise:** SOFeX/MV

PI note: All of these samples were run at sea and data are considered preliminary.

## Data Processing Description

Change history:

070501: downloaded original data (MelvilleStationDataNuts.xls) from SOFeX project data web site.

data prepared by Dave DuBois (OCB DMO, WHOI)

080814: added to OCB database by Cyndy Chandler, OCB DMO, (cchandler@whoi.edu)

PI note: All of these samples were run at sea and data are considered preliminary.

OCB DMO Note: 'CRM' and 'TM' type casts were reported in one worksheet called 'Total'. Data are presented as sorted in the contributed Excel file, with records sorted first by cast type (CRM or TM) and then by time. Date, time and position are from the original data file and not the cruise event log. The 'bot' parameter (bottle number) could be used as a proxy for depth, or used to retrieve depth data from the SIO\_bottle data object.

The latitude in the original Excel file for station 7 CTD cast 19 was reported as a different value for each bottle sample. Those values ranged from -0.0470 to 0.4720. The DMO replaced those values with a single latitude from the cruise event log.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>nutrients_bottle.csv</b> (Comma Separated Values (.csv), 131.58 KB) MD5:c4c2b79b2775fc7b0bec1ed7fff9acab
Primary data file for dataset ID 2944

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
date	date sampling began (UTC)	yyyymmdd
time	time sampling began (UTC)	hhmmss
lon	longitude; negative denotes West	decimal degrees
lat	latitude; negative denotes South	decimal degrees
cast_type	CRM=CTD ??, TM=Trace Metal Free	dimensionless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Niskin Bottle
<b>Generic Instrument Name</b>	Niskin bottle
<b>Generic Instrument Description</b>	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

<b>Dataset-specific Instrument Name</b>	Trace Metal Bottle
<b>Generic Instrument Name</b>	Trace Metal Bottle
<b>Generic Instrument Description</b>	Trace metal (TM) clean rosette bottle used for collecting trace metal clean seawater samples.

[ [table of contents](#) | [back to top](#) ]

## Deployments

### COOK19MV

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57826">https://www.bco-dmo.org/deployment/57826</a>
<b>Platform</b>	R/V Melville
<b>Report</b>	<a href="http://ocb.whoi.edu/SOFeX/CRUISES/proj_description.pdf">http://ocb.whoi.edu/SOFeX/CRUISES/proj_description.pdf</a>
<b>Start Date</b>	2002-01-19
<b>End Date</b>	2002-02-26
<b>Description</b>	<p>Brief cruise plan description: Three ships were involved in the SOFeX experiment. Each ship operated in the study area at a different time to afford the longest observation time. The designations SOFeX-N and SOFeX-S are sometimes used to distinguish between two iron enriched patches - one in low silicate waters north of the polar front (SOFEX-N), and the other in high silicate waters south of the polar front (SOFEX-S). All three ships, Melville (MV), Revelle (RR) and Polar Star (PS), worked in SOFeX-S, but only the Revelle and Melville worked in the SOFeX N patch and shuttled between the two patches. The R/V MELVILLE sailed several weeks after the R/V REVELLE to arrive in the study area just as the 'patches' were forming in response to iron fertilization. The MELVILLE's team planned to make detailed measurements of phytoplankton physiology and rate processes, and to sample daily for phytoplankton growth rates and biomass, soluble and particulate iron and zooplankton biomass. A cruise logbook includes daily entries filed by the Chief Scientist aboard each vessel.</p> <p><b>Methods &amp; Sampling</b>  dates: 21 January 2002 to 21 February 2002 (20020121-20020221) location: N: -52.3853 S: -66.6115 W: -175.2212 E: -166.9392 project/cruise: SOFeX/MV PI note: All of these samples were run at sea and data are considered preliminary.</p> <p><b>Processing Description</b>  Change history: 070501: downloaded original data (MelvilleStationDataNuts.xls) from SOFeX project data web site. data prepared by Dave DuBois (OCB DMO, WHOI) 080814: added to OCB database by Cyndy Chandler, OCB DMO, (<a href="mailto:cchandler@whoi.edu">cchandler@whoi.edu</a>) PI note: All of these samples were run at sea and data are considered preliminary. OCB DMO Note: 'CRM' and 'TM' type casts were reported in one worksheet called 'Total'. Data are presented as sorted in the contributed Excel file, with records sorted first by cast type (CRM or TM) and then by time. Date, time and position are from the original data file and not the cruise event log. The 'bot' parameter (bottle number) could be used as a proxy for depth, or used to retrieve depth data from the SIO_bottle data object. The latitude in the original Excel file for station 7 CTD cast 19 was reported as a different value for each bottle sample. Those values ranged from -0.0470 to 0.4720. The DMO replaced those values with a single latitude from the cruise event log.</p>

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Southern Ocean Iron Experiment (SOFeX)

**Website:** <http://www.mbari.org/expeditions/SOFeX2002/>

**Coverage:** Southern Ocean, south of New Zealand

Before he passed away in 1993, John Martin suggested that an increase in the flow of iron-rich dust to the ocean causes phytoplankton (single celled algae) to grow. The increased photosynthesis removes carbon dioxide from surface waters as the algae create biomass. This carbon dioxide is replaced by carbon dioxide gas that flows into the sea from the atmosphere. Reduced carbon dioxide in the atmosphere cools the planet (CO<sub>2</sub> is a greenhouse gas that warms the earth). The results of this work, funded by the National Science Foundation, the Department of Energy, and the US Coast Guard, will be a much better understanding of how biological processes may regulate climate. (see Related Info: Fe cycle)

A direct test of the 'Martin Hypothesis' that trace concentrations of Fe are responsible for phytoplankton's ability to grow by direct experimental addition of Fe to the surface waters. Consequently the distribution of bioavailable Fe in the surface waters determines large geographical areas primary production and the following flux of fixed organic matter to the deep sea. The aim of the SOFeX project is to investigate the effects of iron fertilization on the productivity of the Southern Ocean. The results of this work will contribute significantly to our understanding of important biogeochemical processes which bear directly on the global carbon cycle, atmospheric carbon dioxide concentration, and climate control.

The SOFeX-N and SOFeX-S designations are sometimes used to distinguish between two iron enriched patches - one in low silicate waters north of the polar front (SOFEX-N), and the other in high silicate waters south of the polar front (SOFEX-S). All three ships, Melville (MV), Revelle (RR) and Polar Star (PS), worked in SOFEX-S, but only the Revelle and Melville worked in the SOFeX N patch and shuttled between the two patches.

[ [table of contents](#) | [back to top](#) ]

---

## **Program Information**

### **Ocean Carbon and Biogeochemistry (OCB)**

**Website:** <http://us-ocb.org/>

**Coverage:** Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO<sub>2</sub> and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

## Iron Synthesis (FeSynth)

**Coverage:** Global

The two main objectives of the Iron Synthesis program (SCOR Working Group proposal, 2005), are:

1. Data compilation: assembling a common open-access database of the *in situ* iron experiments, beginning with the first period (1993-2002; Ironex-1, Ironex-2, SOIREE, EisenEx, SEEDS-1; SOFeX, SERIES) where primary articles have already been published, to be followed by the 2004 experiments where primary articles are now in progress (EIFEX, SEEDS-2; SAGE, FeeP); similarly for the natural fertilizations S.O.JGOFS (1992), CROZEX (2004/2005) and KEOPS (2005).

2. Modeling and data synthesis of specific aspects of two or more such experiments for various topics such as physical mixing, phytoplankton productivity, overall ecosystem functioning, iron chemistry, CO<sub>2</sub> budgeting, nutrient uptake ratios, DMS(P) processes, and combinations of these variables and processes.

SCOR Working Group proposal, 2005. "The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling".

[http://www.scor-int.org/Working\\_Groups/wg131.htm](http://www.scor-int.org/Working_Groups/wg131.htm)

See also: SCOR Proceedings Vol. 42 Concepcion, Chile October 2006, pgs: 13-16 2.3.3 Working Group on The Legacy of *in situ* Iron Enrichments: Data Compilation and Modeling.

The first objective of the Iron Synthesis program involves a data recovery effort aimed at assembling a common, open-access database of data and metadata from a series of *in-situ* ocean iron fertilization experiments conducted between 1993 and 2005. Initially, funding for this effort is being provided by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).

Through the combined efforts of the principal investigators of the individual projects and the staff of Biological and Chemical Oceanography Data Management Office (BCO-DMO), data currently available primarily through individuals, disparate reports and data agencies, and in multiple formats, are being collected and prepared for addition to the BCO-DMO database from which they will be freely available to the community.

As data are contributed to the BCO-DMO office, they are organized into four overlapping categories:

1. Level 1, basic metadata  
(e.g., description of project/study, general location, PI(s), participants);
2. Level 2, detailed metadata and basic shipboard data and routine ship's operations  
(e.g., CTDs, underway measurements, sampling event logs);
3. Level 3, detailed metadata and data from specialized observations  
(e.g., discrete observations, experimental results, rate measurements) and
4. Level 4, remaining datasets  
(e.g., highest level of detailed data available from each study).

Collaboration with BCO-DMO staff began in March of 2008 and initial efforts have been directed toward basic project descriptions, levels 1 and 2 metadata and basic data, with detailed and more detailed data files being incorporated as they become available and are processed.

## Related file

[Program Documentation](#)

The Iron Synthesis Program is funded jointly by the Scientific Committee on Oceanic Research (SCOR) and the U.S. National Science Foundation (NSF).



[ [table of contents](#) | [back to top](#) ]