

Photographs of net collections before any picking from R/V Polarstern (ANT-XXIV_1) from October to November 2007 (CMarZ_2004-2010 project)

Website: <https://www.bco-dmo.org/dataset/2996>

Version:

Version Date: 2010-11-01

Project

» [Census of Marine Zooplankton-2004-2010](#) (CMarZ_2004-2010)

Program

» [Census of Marine Life](#) (CoML)

Contributors	Affiliation	Role
Schiel, Sigi	Alfred Wegener Institute for Polar and Marine Research (AWI)	Chief Scientist
Wiebe, Peter H.	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI)	Scientist
Allison, Dicky	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Dataset Description

Tow Photos from Polarstern Cruise ANT-XXIV-1 Oct - Nov, 2007 Photographs were taken in the "Cold Container", alias the cold wet lab, of the Polarstern of the contents of each net in a tray before any animals were picked out. Nancy Copley used her own equipment. See parameter list for settings.

[[table of contents](#) | [back to top](#)]

Data Files

File
tow_photos_PS24.csv (Comma Separated Values (.csv), 12.94 KB) MD5:9a13c67f6a96dfb0606131960a9b0823
Primary data file for dataset ID 2996

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
net_size	Size of opening of each net on that instrument.	square meters
station	station number	
tow	tow number	
lat	Latitude. Negative = South.	decimal degrees
lon	Longitude. Negative = West.	decimal degrees
net	Number of the applicable net.	
depth_range	Range each net fished. Used with MOCNESS tows.	meters
depth_min	Depth where net opened on the way down (net 0) and depth where the net closed on the way up. Used with MOCNESS data.	meters
depth_max	Depth where the net closed when net going down (net 0) and depth where net opened coming up. Used with MOCNESS data.	meters
comments	notes relative to the functioning of the nets during the tow.	
link_html	Where to find the full-sized pictures. Includes the thumbnail for a quick look.	

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	MOCNESS10
Generic Instrument Name	MOCNESS10
Dataset-specific Description	Mocness 10 with .333micron mesh in nets 1-8 and 3mm mesh in net 0.
Generic Instrument Description	The Multiple Opening/Closing Net and Environmental Sensing System (MOCNESS) is based on the Tucker Trawl principle (Tucker, 1951). The MOCNESS-10 (with 10 m ² nets) carries 6 nets of 3.0-mm circular mesh which are opened and closed sequentially by commands through conducting cable from the surface (Wiebe et al., 1976). In this system, "the underwater unit sends a data frame, comprising temperature, depth, conductivity, net-frame angle, flow count, time, number of open net, and net opening/closing, to the deck unit in a compressed hexadecimal format every 2 seconds and from the deck unit to a microcomputer every 4 seconds" (Wiebe et al., 1985).

[[table of contents](#) | [back to top](#)]

Deployments

ANT-XXIV_1

Website	https://www.bco-dmo.org/deployment/57857
Platform	R/V Polarstern
Report	http://epic.awi.de/28985/1/Sch2009ad.pdf
Start Date	2007-10-26
End Date	2007-11-27

[[table of contents](#) | [back to top](#)]

Project Information

Census of Marine Zooplankton-2004-2010 (CMarZ_2004-2010)

Website: <http://www.cmarz.org/>

Coverage: Global ocean

The Census of Marine Zooplankton (CMarZ) is a field project of the Census of Marine Life (see www.CoML.org). CMarZ is working toward a taxonomically comprehensive assessment of biodiversity of animal plankton throughout the world ocean. The project goal is to produce accurate and complete information on zooplankton species diversity, biomass, biogeographical distribution, genetic diversity, and community structure by 2010. Our taxonomic focus is the animals that drift with ocean currents throughout their lives (i.e., the holozooplankton, Fig. 1). This assemblage currently includes ~6,800 described species in fifteen phyla; our expectation is that at least that many new species will be discovered as a result of our efforts. The census encompasses unique marine environments and those likely to be inhabited by endemic and undescribed zooplankton species.

Program Information

Census of Marine Life (CoML)

Website: <http://www.coml.org/>

Coverage: global

The Census of Marine Life is a global network of researchers in more than 80 nations engaged in a 10-year scientific initiative to assess and explain the diversity, distribution, and abundance of life in the oceans. The world's first comprehensive Census of Marine Life - past, present, and future - will be released in 2010.

The stated purpose of the Census of Marine Life is to assess and explain the diversity, distribution, and abundance of marine life. Each plays an important role in what is known, unknown, and may never be known about what lives in the global ocean.

First, diversity. The Census aims to make for the first time a comprehensive global list of all forms of life in the sea. No such unified list yet exists. Census scientists estimate that about 230,000 species of marine animals have been described and reside in jars in collections in museums of natural history and other repositories. Since the Census began in 2000, researchers have added more than 5600 species to the lists. They aim to add many thousands more by 2010. The database of the Census already includes records for more than 16 million records, old and new. By 2010, the goal is to have all the old and the new species in an on-line encyclopedia with a webpage for every species. In addition, we will estimate how many species remain unknown, that is, remain to be discovered. The number could be astonishingly large, perhaps a million or more, if all small animals and protists are included. For comparison, biologists have described about 1.5 million terrestrial plants and animals.

Second, distribution. The Census aims to produce maps where the animals have been observed or where they could live, that is, the territory or range of the species. Knowing the range matters a lot for people concerned about, for example, possible consequences of global climate change.

Third, abundance. No Census is complete without measures of abundance. We want to know not only that there is such a thing as a Madagascar crab but how many there are. For marine life, populations are being estimated either in numbers or in total kilos, called biomass.

To complete the context, it is important to understand the top motivations for the Census of Marine Life. Most importantly, much of the ocean is unexplored. Most of the records in its database are for observations near the surface, and down to 1000 meters. No observations have been made in most of the deep ocean, while most of the ocean is deep.

Another important issue is that diversity varies in space. Marine hot spots, like the rain forests of the land, exist off for large fish off the coasts of Brazil and Australia. The goal is to know much more about marine hot spots, to help conserve these large fish. Their abundance and thus their diversity is changing, especially for commercially important species. Between 1952 and 1976, for example, fishermen and their customers emptied many areas of the ocean of tuna.

The Census has evolved a strategy of 14 field projects to touch the major habitats and groups of species in the global ocean. Eleven field projects address habitats, such as seamounts or the Arctic Ocean. Three field projects look globally at animals that either traverse the seas or appear globally distributed: the top predators such as tuna and the plankton and the microbes. The projects employ a mix of technologies. These include acoustics or sound, optics or cameras, tags placed on individual animals that store or report data, and genetics, as well as some actual capture of animals. The technologies complement one another. Sound can survey large areas in the ocean, while light cannot. Light can capture detail and characters that sound cannot. And genetics can make identifications from fragments of specimens or larvae where pictures tell little.

This mix of curiosity, need to know, technology, and scientists willing to investigate the unexplored and undiscovered will result in a Census of Marine Life in 2010 that provides a much clearer picture of what lives below the surface around the globe. Several reasons make such a report timely, indeed urgent. Crises in the sea are reported regularly. One recent study predicted the end of commercial fishery globally by 2050, if

current trends persist. Better information is needed to fashion the management that will sustain fisheries, conserve diversity, reverse losses of habitat, reduce impacts of pollution, and respond to global climate change. Hence, there are biological, economic, philosophical and political reasons to push for greater exploration and understanding of the ocean and its inhabitants. Indeed, the United Nations Convention on Biological Diversity requires signatories to collect information on living resources, but, as yet, no nation has a complete baseline of such information. The Census of Marine Life's global network of researchers will help to fill this knowledge gap, providing critical information to help guide decisions on how to manage global marine resources for the future.

[Text copied from the CoML web site, November 5, 2008]

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
Alfred P. Sloan Foundation (Sloan)	unknown CMarZ_2004-2010 Sloan

[[table of contents](#) | [back to top](#)]