

# Temperature logger data from MAPCO2 buoy deployed on NOAA Ship Ronald H. Brown cruise RB-08-02 in the Southwest Atlantic sector of the Southern Ocean near South Georgia Island in 2008 (SO\_GasEx project)

**Website:** <https://www.bco-dmo.org/dataset/3288>

**Version:** 07 Jan 2010

**Version Date:** 2010-01-07

## Project

» [Southern Ocean Gas Exchange Experiment](#) (SO\_GasEx)

## Programs

» [Ocean Carbon and Biogeochemistry](#) (OCB)

» [United States Surface Ocean Lower Atmosphere Study](#) (U.S. SOLAS)

Contributors	Affiliation	Role
<a href="#">McGillis, Wade</a>	Lamont-Doherty Earth Observatory (LDEO)	Principal Investigator
<a href="#">DeGrandpre, Michael</a>	University of Montana	Co-Principal Investigator
<a href="#">Zappa, Christopher J.</a>	Lamont-Doherty Earth Observatory (LDEO)	Co-Principal Investigator
<a href="#">Hales, Burke</a>	Oregon State University (OSU-CEOAS)	Contact
<a href="#">Gegg, Stephen R.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Dataset Description

SO-GasEx MAPCO2 Buoy - Temperature Logger data

Temperature logger datasets from 18 temperature loggers deployed on the MAPCO2 buoy

## Methods & Sampling

**For MAPCO2 Buoy Information See:** [SO-GasEx cruise report, Section 5.5.1 pgs 30-33](#)

Onset PRO V2 temperature loggers were placed at depths between 5 and 105 m depth on the MapCO2 buoy. These loggers have an accuracy of  $0.1 \pm 0.02$  °C.

## NOAA Buoy Temperature Sensors Summary - Probe Locations

### Deployment 1    Deployment 2    Deployment 3

Depth Sensor Comment    Depth Sensor Comment    Depth Sensor Comment

5 O SAMI-52    5 O SAMI-52    5 O SAMI-52

19 P SAMI-46    19 P SAMI-46    19 P SAMI-46

25 L Top Drogue 25 L Top Drogue 25 L Top Drogue  
 34 K Bottom Drogue 34 K Bottom Drogue 34 K Bottom Drogue  
 34.5 Q SAMI-11 34.5 Q SAMI-11 34.5 Q SAMI-11  
 40.5 J Top Drogue 40.5 J Top Drogue 40.5 J Top Drogue  
 49.5 I Bottom Drogue 49.5 I Bottom Drogue 49.5 I Bottom Drogue  
 50 R SAMI-48 50 R SAMI-48 50 R SAMI-48  
 56 H Top Drogue 56 F Top Drogue 56 F Top Drogue  
 65 G Bottom Drogue 65 E Bottom Drogue 65 E Bottom Drogue  
 65.75 F Top Drogue 65.75 N Top Cable 65.75 N Top Cable  
 74.75 E Bottom Drogue 74.75 M Bottom Cable 74.75 M Bottom Cable  
 75.25 S SAMI-16 75.25 S SAMI-16 75.25 S SAMI-16  
 81.25 D Top Drogue 81.25 B Top Drogue 81.25 B Top Drogue  
 90.25 C Bottom Drogue 90.25 A Bottom Drogue 90.25 A Bottom Drogue  
 91 B Top Drogue 96 T SAMI-15 96 T SAMI-15  
 100 A Bottom Drogue  
 105 T SAMI-15

## Data Processing Description

For MAPCO2 Buoy Information See: [SO-GasEx cruise report, Section 5.5.1 pgs 30-33](#)

### BCO-DMO Processing Notes

- Generated from original multi sheet file Temperature Probes.xls

### BCO-DMO Edits

- multiple sheet .xls file converted to single .xls file containing all three deployments
- DeploymentId added to each data record as Deployment\_1, etc.
- single date/time field converted to separate fields and formatted to BCO-DMO convention
- lat/lon added from SAMI CO2, O2, T data files
- any blank fields changed to 'nd' (no data)
- data parameter names reformatted to BCO-DMO convention and depth added to each

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>MAPCO2_Temp.csv</b> (Comma Separated Values (.csv), 133.18 KB) MD5:06f950ef6b1f96940f244c4aa0cfd1bf
Primary data file for dataset ID 3288

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
DeploymentId	MAPCO2 Buoy Deployment Id	text
date	Date (UTC)	YYYYMMDD
time	Time (UTC)	HHMM
lon	buoy longitude position in decimal degrees (West is negative)	decimal degrees
lat	buoy latitude position in decimal degrees (South is negative)	decimal degrees
O_Temp_5m	Temperature from sensor O/SAMI-52 mounted at 5meters depth	degrees celcius

P_Temp_19m	Temperature from sensor P/SAMI-46 mounted at 19meters depth	degrees celcius
L_Temp_25m	Temperature from sensor L/Top Drogue mounted at 25meters depth	degrees celcius
K_Temp_34m	Temperature from sensor K/Bottom Drogue mounted at 34meters depth	degrees celcius
Q_Temp_34point5m	Temperature from sensor Q/SAMI-11 mounted at 34.5meters depth	degrees celcius
J_Temp_40point5m	Temperature from sensor J/Top Drogue mounted at 40.5meters depth	degrees celcius
I_Temp_49point5m	Temperature from sensor I/Bottom Drogue mounted at 49.5meters depth	degrees celcius
R_Temp_50m	Temperature from sensor R/SAMI-48 mounted at 50meters depth	degrees celcius
F_Temp_56m	Temperature from sensor F/Top Drogue mounted at 56meters depth	degrees celcius
E_Temp_65m	Temperature from sensor E/Bottom Drogue mounted at 65meters depth	degrees celcius
N_Temp_65point75m	Temperature from sensor N/Top Cable mounted at 65.75meters depth	degrees celcius
M_Temp_74point75m	Temperature from sensor M/Bottom Cable mounted at 75.75meters depth	degrees celcius
S_Temp_75point25m	Temperature from sensor S/SAMI-16 mounted at 75.25meters depth	degrees celcius
B_Temp_81point25m	Temperature from sensor B/Top Drogue mounted at 81.25meters depth	degrees celcius
A_Temp_90point25m	Temperature from sensor A/Bottom Drogue mounted at 90.25meters depth	degrees celcius
T_Temp_96m	Temperature from sensor T/SAMI-15 mounted at 96meters depth	degrees celcius
H_Temp_56m	Temperature from sensor H/Top Drogue mounted at 56meters depth	degrees celcius
G_Temp_65m	Temperature from sensor G/Bottom Drogue mounted at 65meters depth	degrees celcius
F_Temp_65point75m	Temperature from sensor F/Top Drogue mounted at 65.75meters depth	degrees celcius
E_Temp_74point75m	Temperature from sensor E/Bottom Drogue mounted at 74.75meters depth	degrees celcius
D_Temp_81point25m	Temperature from sensor D/Top Drogue mounted at 81.25meters depth	degrees celcius
C_Temp_90point25m	Temperature from sensor C/Bottom Drogue mounted at 90.25meters depth	degrees celcius
B_Temp_91m	Temperature from sensor B/Top Drogue mounted at 91meters depth	degrees celcius
A_Temp_100m	Temperature from sensor A/Bottom Drogue mounted at 100meters depth	degrees celcius
T_Temp_105m	Temperature from sensor T/SAMI-15 mounted at 105meters depth	degrees celcius

## Instruments

<b>Dataset-specific Instrument Name</b>	MAPCO2 Drifting Buoy
<b>Generic Instrument Name</b>	MAPCO2 Drifting Buoy
<b>Dataset-specific Description</b>	For SO-GasEx MAPCO2 Buoy Configuration See: SO-GasEx MAPCO2 Metadata Report
<b>Generic Instrument Description</b>	The Moored Autonomous pCO <sub>2</sub> (MAPCO <sub>2</sub> ) surface drifting buoy designed by NOAA/PMEL is a low profile, high payload buoy. It was used in the SO GasEx project as a drogued drifter instrumented with a variety of autonomous instruments capable of making a coordinated set of physical, geochemical, and biological measurements at high temporal resolutions. These measurements provide a key component in the study of processes controlling air-sea CO <sub>2</sub> exchange.
<b>Dataset-specific Instrument Name</b>	Onset PRO V2 temperature logger
<b>Generic Instrument Name</b>	Onset HOBO Pro v2 temperature logger
<b>Dataset-specific Description</b>	NOAA Buoy Temperature Sensors Summary - Probe Locations Deployment 1 Deployment 2 Deployment 3 Depth Sensor Comment Depth Sensor Comment Depth Sensor Comment 5 O SAMI-52 5 O SAMI-52 5 O SAMI-52 19 P SAMI-46 19 P SAMI-46 19 P SAMI-46 25 L Top Drogue 25 L Top Drogue 25 L Top Drogue 34 K Bottom Drogue 34 K Bottom Drogue 34 K Bottom Drogue 34.5 Q SAMI-11 34.5 Q SAMI-11 34.5 Q SAMI-11 40.5 J Top Drogue 40.5 J Top Drogue 40.5 J Top Drogue 49.5 I Bottom Drogue 49.5 I Bottom Drogue 49.5 I Bottom Drogue 50 R SAMI-48 50 R SAMI-48 50 R SAMI-48 56 H Top Drogue 56 F Top Drogue 56 F Top Drogue 65 G Bottom Drogue 65 E Bottom Drogue 65 E Bottom Drogue 65.75 F Top Drogue 65.75 N Top Cable 65.75 N Top Cable 74.75 E Bottom Drogue 74.75 M Bottom Cable 74.75 M Bottom Cable 75.25 S SAMI-16 75.25 S SAMI-16 75.25 S SAMI-16 81.25 D Top Drogue 81.25 B Top Drogue 81.25 B Top Drogue 90.25 C Bottom Drogue 90.25 A Bottom Drogue 90.25 A Bottom Drogue 91 B Top Drogue 96 T SAMI-15 96 T SAMI-15 100 A Bottom Drogue 105 T SAMI-15
<b>Generic Instrument Description</b>	The HOBO Water Temp Pro v2 temperature logger, manufactured by Onset Computer Corporation, has 12-bit resolution and a precision sensor for ±0.2°C accuracy over a wide temperature range. It is designed for extended deployment in fresh or salt water. Operation range: -40° to 70°C (-40° to 158°F) in air; maximum sustained temperature of 50°C (122°F) in water Accuracy: 0.2°C over 0° to 50°C (0.36°F over 32° to 122°F) Resolution: 0.02°C at 25°C (0.04°F at 77°F) Response time: (90%) 5 minutes in water; 12 minutes in air moving 2 m/sec (typical) Stability (drift): 0.1°C (0.18°F) per year Real-time clock: ± 1 minute per month 0° to 50°C (32° to 122°F) Additional information ( <a href="http://www.onsetcomp.com/">http://www.onsetcomp.com/</a> ) Onset Computer Corporation 470 MacArthur Blvd Bourne, MA 02532

## Deployments

RB-08-02

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/57846">https://www.bco-dmo.org/deployment/57846</a>
<b>Platform</b>	NOAA Ship Ronald H. Brown
<b>Report</b>	<a href="http://bcodata.whoi.edu/SO-GasEx/SO_GasEx_Cruise_Report.pdf">http://bcodata.whoi.edu/SO-GasEx/SO_GasEx_Cruise_Report.pdf</a>
<b>Start Date</b>	2008-02-29
<b>End Date</b>	2008-04-12
<b>Description</b>	<p>The Southern Ocean GasEx experiment was conducted aboard the NOAA ship Ronald H. Brown with 31 scientists representing 22 institutions, companies and government labs. The cruise departed Punta Arenas, Chile on 29 February, 2008 and transited approximately 5 days to the nominal study region at 50°S, 40°W in the Atlantic sector of the Southern Ocean. The scientific work concentrated on quantifying gas transfer velocities using deliberately injected tracers, measuring CO<sub>2</sub> and DMS fluxes directly in the marine air boundary layer, and elucidating the physical, chemical, and biological processes controlling air-sea fluxes with measurements in the upper-ocean and marine air. The oceanic studies used a Lagrangian approach to study the evolution of chemical and biological properties over the course of the experiment using shipboard and autonomous drifting instruments. The first tracer patch was created and studied for approximately 6 days before the ship was diverted from the study site, 350 miles to the south, to wait near South Georgia Island for calmer seas. After more than 4 days away, we returned to the study area and managed to find some remnants of the tracer patch. After collecting one final set of water column samples and recovering the two drifting buoys deployed with the patch, we relocated to the northwest, closer to the area where the first patch was started. A second tracer patch was created and studied for approximately 15 days before we had to break off the experiment and transit to Montevideo, Uruguay for the completion of the cruise.</p>

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Southern Ocean Gas Exchange Experiment (SO\_GasEx)

**Website:** <http://so-gasex.org/>

**Coverage:** Southwest Atlantic sector of the Southern Ocean (nominally at 50°S, 40°W, near South Georgia Island)

The Southern Ocean Gas Exchange Experiment (SO-GasEx; also known as GasEx III) took place in the Southwest Atlantic sector of the Southern Ocean (nominally at 50°S, 40°W, near South Georgia Island) in austral fall of 2008 (February 29-April 12, 2008) on the [NOAA ship Ronald H. Brown](#). SO-GasEX is funded by NOAA, NSF and NASA.

The research objectives for Southern Ocean GasEx are to answer the following questions:

- What are the gas transfer velocities at high winds?
- What is the effect of fetch on the gas transfer?
- How do other non-direct wind effects influence gas transfer?
- How do changing pCO<sub>2</sub> and DMS levels affect the air-sea CO<sub>2</sub> and DMS flux, respectively in the same locale?
- Are there better predictors of gas exchange in the Southern Ocean other than wind?
- What is the near surface horizontal and vertical variability in turbulence, pCO<sub>2</sub>, and other relevant biochemical and physical parameters?
- How do biological processes influence pCO<sub>2</sub> and gas exchange?
- Do the different disparate estimates of fluxes agree, and if not why?
- With the results from Southern Ocean GasEx, can we reconcile the current discrepancy between model based CO<sub>2</sub> flux estimates and observation based estimates?

## Related files

[SO-GasEx cruise report](#)  
[SO-GasEx Science Plan](#)  
[SO-GasEx Implementation Plan](#)

The SO-GasEx cruise report and Science and Implementation plans, may also be available at [the SO-GasEx science Web page](#).

[ [table of contents](#) | [back to top](#) ]

---

## Program Information

### Ocean Carbon and Biogeochemistry (OCB)

**Website:** <http://us-ocb.org/>

**Coverage:** Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO<sub>2</sub> and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

### United States Surface Ocean Lower Atmosphere Study (U.S. SOLAS)

**Website:** <http://www.us-solas.org/>

**Coverage:** Global

The Surface Ocean Lower Atmosphere Study (SOLAS) program is designed to enable researchers from different disciplines to interact and investigate the multitude of processes and interactions between the coupled ocean and atmosphere.

Oceanographers and atmospheric scientists are working together to improve understanding of the fate, transport, and feedbacks of climate relevant compounds, and also weather and hazards that are affected by processes at the surface ocean.

Oceanographers and atmospheric scientists are working together to improve understanding of the fate, transport, and feedbacks of climate relevant compounds.

Physical, chemical, and biological research near the ocean-atmosphere interface must be performed in synergy to extend our current knowledge to adequately understand and forecast changes on short and long time frames and over local and global spatial scales.

The findings obtained from SOLAS are used to improve knowledge at process scale that will lead to better quantification of fluxes of climate relevant compounds such as CO<sub>2</sub>, sulfur and nitrogen compounds, hydrocarbons and halocarbons, as well as dust, energy and momentum. This activity facilitates a fundamental understanding to assist the societal needs for climate change, environmental health, weather prediction, and national security.

The US SOLAS program is a component of the International SOLAS program where collaborations are forged with investigators around the world to examine SOLAS issues ubiquitous to the world's oceans and atmosphere.

[Â» International SOLAS Web site](#)

### Science Implementation Strategy Reports

[US-SOLAS](#) (4 MB PDF file)

[Other SOLAS reports](#) are available for download from the US SOLAS Web site

[ [table of contents](#) | [back to top](#) ]

---

### Funding

Funding Source	Award
National Oceanic and Atmospheric Administration (NOAA)	<a href="#">unknown SO_GasEx NOAA</a>
National Aeronautics & Space Administration (NASA)	<a href="#">unknown SO_GasEx NASA</a>
National Science Foundation (NSF)	<a href="#">unknown SO_GasEx NSF</a>

[ [table of contents](#) | [back to top](#) ]