CTD Profile Data from R/V Knorr KN197-08 in the Amazon River plume, NE coast of South America from May to June 2010 (ANACONDAS project)

Website: https://www.bco-dmo.org/dataset/3374 Version: 14 October 2010 Version Date: 2010-10-14

Project

» <u>Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses</u> (ANACONDAS)

Programs

- » Integrated Marine Biogeochemistry and Ecosystem Research -US (IMBER-US)
- » <u>Ocean Carbon and Biogeochemistry</u> (OCB)
- » Emerging Topics in Biogeochemical Cycles (ETBC)
- » Marine Microbiology Initiative (MMI)

Contributors	Affiliation	Role
<u>Coles, Victoria</u>	University of Maryland Center for Environmental Science (UMCES/HPL)	Principal Investigator, Contact
<u>Gegg, Stephen</u> <u>R.</u>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Parameters
- Instruments
- Deployments
- <u>Project Information</u>
- <u>Program Information</u>
- <u>Funding</u>

Dataset Description

CTD profile data

Methods & Sampling

SBE9+ s/n 785 Pressure calibration 08 Aug 2008 Channel 1: SBE4c s/n 04 3089 Conductivity calibration 16 Mar 2010 SBE3T s/n 03P 4502 Temperature calibration 24 Mar 2010 SBE43 s/n 43 1642 Dissolved Oxygen calibration 12 June 2009

Channel 2: SBE4c s/n 04 3042 Conductivity calibration 24 Mar 2010 SBE3T s/n 03P 4507 Temperature calibration 12 Mar 2010

Transmissometer WetLabs Model CSTar s/n CST - 1117DR Calibrated 30 April 2008

Fluorometer WetLabs Model FLNTU s/n FLNTURD-304 Calibrated 10 March 2008

Surface PAR Biospherical Instruments QSR-240 s/n 6294 Calibrated 13 March 2008

Aquatic PAR Biospherical Instruments s/n QSP-200L4 s/n 4550 Calibrated 13 March 2008

Channel 2 is used for most profiles as there were problems with the pump for channel 1 on some stations. Oxygen is taken from channel 1, however there are some bad values which are not yet flagged.

Sample header file:

* Sea-Bird SBE 9 Data File: * FileName = C:Datactd19708001.hdr * Software Version Seasave V 7.18d * Temperature SN = 4502* Conductivity SN = 3089* Number of Bytes Per Scan = 40 * Number of Voltage Words = 5 * Number of Scans Averaged by the Deck Unit = 1* System UpLoad Time = May 23 2010 08:09:54 * NMEA Latitude = 11 33.61 N * NMEA Longitude = 056 47.89 W * NMEA UTC (Time) = May 23 2010 08:09:51 * Store Lat/Lon Data = Append to Every Scan ** Ship: R/V Knorr ** Station: 001 ** Operator: Victoria ** station 001 cast 01 event 01.03 # nguan = 23 # nvalues = 1994# units = specified # name 0 = prDM: Pressure, Digiguartz [db] # name 1 = depSM: Depth [salt water, m] # name 2 = t190C: Temperature, 2 [ITS-90, deg C] # name 3 = c1S/m: Conductivity, 2 [S/m] # name 4 = sbeox0V: Oxygen Voltage, SBE 43 # name 5 = xmiss: Beam Transmission, Chelsea/Seatech/Wetlab CStar [%] # name 6 = bat: Beam Attenuation, Chelsea/Seatech/Wetlab CStar [1/m]# name 7 = flECO-AFL: Fluorescence, Wetlab ECO-AFL/FL [mg/m^3] # name 8 = upoly0: Upoly 0, FLNTU turbidity # name 9 = spar: SPAR/Surface Irradiance # name 10 = par: PAR/Irradiance, Biospherical/Licor # name 11 = cpar: CPAR/Corrected Irradiance [%] # name 12 = t090C: Temperature [ITS-90, deg C] # name 13 = c0S/m: Conductivity [S/m] # name 14 = depSM: Depth [salt water, m], lat = 11.5602 # name 15 = density11: Density, 2 [density, Kg/m^3] # name 16 = sigma-t11: Density, 2 [sigma-t, Kg/m^3] # name 17 = potemp190C: Potential Temperature, 2 [ITS-90, deg C] # name 18 = sal11: Salinity, Practical, 2 [PSU] # name 19 = svWM: Sound Velocity [Wilson, m/s] # name 20 = sbeox0Mm/Kg: Oxygen, SBE 43 [umol/Kg], WS = 2 # name 21 = sbeox0PS: Oxygen, SBE 43 [% saturation], WS = 2 # name 22 = flag; flag # span 0 = 5.029, 2019.162 # span 1 = 4.978, 1997.855 # span 2 = 3.4786, 28.9724 # span 3 = 3.293110, 5.786188 # span 4 = 1.2362, 2.4303 # span 5 = 93.5592, 97.7670 # span 6 = 0.0903, 0.2663 # span 7 = -0.0113, 0.4303 # span 8 = 1.0607965, 1.3869948 # span 9 = 1.5656e+01, 1.7824e+01 # span 10 = 1.0000e-12, 1.0000e-12

```
\# span 11 = 5.6114e-12, 6.3893e-12
# span 12 = 3.4787, 28.9723
# span 13 = 3.293104, 5.786650
# span 14 = 5.000, 1998.000
# span 15 = 1020.0771, 1037.0397
# span 16 = 20.0517, 27.8118
# span 17 = 3.3170, 28.9661
# span 18 = 32.1813, 37.0067
# span 19 = 1486.29, 1544.67
# span 20 = 112.419, 249.669
# span 21 = 38.99671, 96.04706
\# span 22 = 0.0000e+00, 0.0000e+00
# interval = meters: 1
# start time = May 23 2010 08:09:51
\# bad flag = -9.990e-29
# datcnv date = Jun 23 2010 19:27:22, 7.20c
# datcnv in = Z:Data aw19708001.hex Z:Data aw19708001.CON
# datcnv skipover = 0
# datcnv ox hysteresis correction = yes
# filter date = Jun 23 2010 19:27:59, 7.20c
# filter in = Z:Datavicprocess19708001.cnv
# filter low pass tc A = 0.500
# filter low pass to B = 0.150
# filter low pass A vars = depSM t190C c1S/m sbeox0V xmiss bat fIECO-AFL upoly0 spar cpar t090C c0S/m
# filter low pass B vars = prDM
# alignctd date = Jun 23 2010 19:28:04, 7.20c
# alignctd in = Z:Datavicprocess19708001.cnv
# alignctd adv = c1S/m 0.073, sbeox0V 2.000
# celltm date = Jun 23 2010 19:28:09, 7.20c
# celltm in = Z:Datavicprocess19708001.cnv
\# celltm alpha = 0.0000, 0.0300
# celltm tau = 0.0000, 7.0000
# celltm_temp_sensor_use_for_cond = , secondary
# loopedit_date = Jun 23 2010 19:28:14, 7.20c
# loopedit in = Z:Datavicprocess19708001.cnv
# loopedit minVelocity = 0.250
# loopedit surfaceSoak: minDepth = 5.0, maxDepth = 20, useDeckPress = 1
# loopedit excl bad scans = yes
# wildedit date = Jun 23 2010 19:28:19, 7.20c
# wildedit in = Z:Datavicprocess19708001.cnv
# wildedit pass1 nstd = 2.0
# wildedit pass2 nstd = 20.0
# wildedit pass2 mindelta = 0.000e+000
# wildedit npoint = 100
# wildedit vars = prDM depSM t190C c1S/m xmiss bat fIECO-AFL upoly0 spar par cpar t090C c0S/m
# wildedit excl bad scans = yes
# wfilter date = Jun 23 2010 19:28:23, 7.20c
# wfilter in = Z:Datavicprocess19708001.cnv
# wfilter excl bad scans = yes
# wfilter action xmiss = gaussian, 200, 100, 0
# wfilter action bat = gaussian, 200, 100, 0
# wfilter action flECO-AFL = gaussian, 200, 100, 0
# Derive date = Jun 23 2010 19:28:31, 7.20c
# Derive in = Z:Datavicprocess19708001.cnv Z:Data aw19708001.con
# derive time window docdt = seconds: 2
# derive ox tau correction = yes
# binavg_date = Jun 23 2010 19:28:45, 7.20c
# binavg in = Z:Datavicprocess19708001.cnv
# binavg bintype = meters
# binavg binsize = 1
# binavg excl bad scans = no
# binavg skipover = 0
# binavg_surface_bin = yes, min = 0.000, max = 0.000, value = 0.000
```

Data Processing Description

The data have not yet been processed to include hand salinity and oxygen samples.

BCO-DMO Processing Notes

- Awk written to reformat original .cnv files contributed by Victoria Coles
- AWK: ANACONDAS_CTDcnv_2_bcodmo.awk
- Header data for CTD data generated from .avg.btl.csv files
- space delimited reformatted to tab delimited
- all records with "#" or "*" ignored
- blank lines ignored
- BCO-DMO header o/p from routine
- Parameter names same as Bottle Data wherever possible

[table of contents | back to top]

Data Files

File

CTD_Profiles.csv(Comma Separated Values (.csv), 16.97 MB) MD5:fe19ca4c9f1af12e3bf16cc81afec3fd

Primary data file for dataset ID 3374

[table of contents | back to top]

Parameters

Parameter	Description	Units
CTD_DataSet	CTD profile data dataset id (19708001)	text
CTD	unique CTD number	integer
Station	Station number	integer
Cast	CTD cast at the same station	integer
Event	Event number = station.event.number at station	xx.xx
date_gmt	Station date (GMT)	YYYYMMDD
time_gmt	Station time (GMT)	HHMMSS
lon	Station longitude (West is negative)	decimal degrees
lat	Station latitude (South is negative)	decimal degrees
Press	Pressure	decibars
Depth	Depth of instrument	meters
Temp2	Temperature from sensor 2	degrees Celsius
Cond2	Conductivity from sensor 2	Siemens/meter
02_v	Oxygen in volts	volts
BeamTrans	Beam Transmission	percentage
BeamAtten	Beam Attenuation	m-1
Fluorescence	Fluorescence	mgm-3
Turbidity	Turbidity	Nephelometric Turbidity Units (NTU)
Spar	photosynthetically active radiation (S PAR)	uEm-2sec-1
PAR	photosynthetically active radiation (PAR)	uEm-2sec-1
Cpar	photosynthetically active radiation (C PAR)	uEm-2sec-1
Temp1	Temperature from sensor 1	degrees Celsius
Cond1	Conductivity from sensor 1	Siemens/meter
Depth_Lat	Latitude corrected depth of instrument lat - 11.2883	meters
Density	Density	kgm-2
Sigma_theta	Sigma theta	kgm-2
Pot_Temp	Potential Temperature	degrees Celsius
Salinity	Salinity	practical salinity units (PSU)
Sound_Velocity	Sound Velocity	m/sec
02	Oxygen	umolkg-1
O2_PS	Oxygen Percent Saturation (percentage)	percentage
Flag	Data Flag (always 0 in these data)	integer

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	CTD Sea-Bird SBE 911plus
Generic Instrument Name	CTD Sea-Bird SBE 911plus
Generic Instrument Description	The Sea-Bird SBE 911 plus is a type of CTD instrument package for continuous measurement of conductivity, temperature and pressure. The SBE 911 plus includes the SBE 9plus Underwater Unit and the SBE 11plus Deck Unit (for real-time readout using conductive wire) for deployment from a vessel. The combination of the SBE 9 plus and SBE 11 plus is called a SBE 911 plus. The SBE 9 plus uses Sea-Bird's standard modular temperature and conductivity sensors (SBE 3 plus and SBE 4). The SBE 9 plus CTD can be configured with up to eight auxiliary sensors to measure other parameters including dissolved oxygen, pH, turbidity, fluorescence, light (PAR), light transmission, etc.). more information from Sea-Bird Electronics

Dataset- specific Instrument Name	Fluorometer
Generic Instrument Name	Fluorometer
Generic Instrument	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.

Dataset- specific Instrument Name	Photosynthetically Available Radiation Sensor
Generic Instrument Name	Photosynthetically Available Radiation Sensor
Generic Instrument	A PAR sensor measures photosynthetically available (or active) radiation. The sensor measures photon flux density (photons per second per square meter) within the visible wavelength range (typically 400 to 700 nanometers). PAR gives an indication of the total energy available to plants for photosynthesis. This instrument name is used when specific type, make and model are not known.

Dataset- specific Instrument Name	Wet Labs CSTAR Transmissometer
Generic Instrument Name	WET Labs {Sea-Bird WETLabs} C-Star transmissometer
Generic Instrument Description	The C-Star transmissometer has a novel monolithic housing with a highly intgrated opto- electronic design to provide a low cost, compact solution for underwater measurements of beam transmittance. The C-Star is capable of free space measurements or flow-through sampling when used with a pump and optical flow tubes. The sensor can be used in profiling, moored, or underway applications. Available with a 6000 m depth rating. More information on Sea-Bird website: <u>https://www.seabird.com/c-star-transmissometer/product?id=60762467717</u>

Deployments

KN197-08	
Website	https://www.bco-dmo.org/deployment/58043
Platform	R/V Knorr
Report	http://bcodata.whoi.edu/ANACONDAS/ANACONDAS1-FullCruiseReport.pdf
Start Date	2010-05-22
End Date	2010-06-24
Description	ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom SymbiosesROCA: River Ocean Continuum of the Amazon WHOI cruise planning synopsis Cruise information and original data are available from the NSF R2R data catalog. Cruise Track over Salinity Climatology (Image from Dr. Ajit Subramaniam)

[table of contents | back to top]

Project Information

Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses (ANACONDAS)

Website: <u>http://amazoncontinuum.org/</u>

Coverage: Amazon River plume; NE coast of South America; Western Tropical North Atlantic - 15N-Equator and 60W to 45W - Region surrounding the Amazon River Plume

ANACONDAS is an IMBER endorsed project. <u>View list of all IMBER endorsed projects</u>

View the ANACONDAS project GCMD DIF record

The ANACONDAS project was funded as part of the US National Science Foundation (NSF) Emerging Topics in Biogeochemical Cycles (ETBC) program (Directorate for Geosciences, NSF 07 -049, September 19, 2007) explicitly intended to support emerging areas of interdisciplinary research. The ETBC program aimed to foster transformational advances in the quantitative or mechanistic understanding of biogeochemical cycles that integrated physical-chemical-biological processes over the range of temporal and/or spatial scales in Earth's environments. The program especially sought proposals that addressed emerging topics in biogeochemical cycles, the water cycle or their coupling, across the interfaces of atmosphere, land, and oceans.

The ANACONDAS investigators hypothesize that large tropical river plumes with low N: P ratios provide an ideal niche for diatom-diazotroph assemblages (DDAs). They suggest that the ability of these organisms to fix N2 within the surface ocean is responsible for significant C export in the Amazon River plume. Their previous observations in the Amazon River plume helped reveal that blooms comprised of the endosymbiotic N2-fixing cyanobacterium Richelia and its diatom hosts (e.g. Hemiaulus) were a significant source of new production and carbon export. The previous work focused largely on the sensitivity of DDAs to external forcing from dust and riverine inputs, so the ecology of these organisms and the fate of their new production were largely unstudied. It is now known that DDAs are responsible for a significant amount of CO2 drawdown in the Amazon River plume, and floating sediment traps at 200 m measured 4x higher mass fluxes beneath the plume than outside the plume. This led the researchers to hypothesize that this greater export is due either to aggregation and sinking of DDAs themselves or to grazing of DDAs by zooplankton.

In this study the researchers will undertake a suite of field, satellite and modeling studies aimed at understanding the ecology and tracing the fate of C and N fixed by DDAs and other phytoplankton living in the plume. By examining C and silicate (Si) export from offshore surface waters, through the upper oceanic food web, the mesopelagic, and down to the deep sea floor, they will quantify the impact of the Amazon River on biological processes that control C sequestration and the implications of these regional processes on C, N and Si budgets. The study will go beyond previous research because they will quantify 1) the distribution, nutrient demands, and activity of DDAs in the context of phytoplankton species succession, 2) the sensitivity of the CO2 drawdown to the mix of phytoplankton, 3) the grazing and aggregation processes contributing to the sinking flux, 4) the composition of this flux, and 5) the proportion of this material that reaches the seafloor. This effort truly represents a measure of C sequestration and pump efficiency. Ecological modeling will be used to place observational results from field studies and satellites into the context of the larger Atlantic basin with tropical climate variability on interannual and longer time scales.

Three cruises were carried out during the ANACONDAS project: AN10/KN197-08 - R/V KNORR - May/June 2010 - <u>Cruise Track over Salinity Climatology</u> (*Image: Yager, et al, 2007*) AN11/MV1110 - R/V MELVILLE - September/October 2011 - <u>Cruise Track over Salinity Climatology</u> (*Image: Yager, et al, 2007*) AN12/AT21-04 - R/V ATLANTIS - July/2012 - <u>Cruise Track over Salinity Climatology</u> (*Image: Yager, et al, 2007*)

The ANACONDAS project builds on observations made by MANTRA/PIRANA in 2001 and 2003 (RV Knorr and Seward Johnson I cruises to the same region) to address specifically 1) how carbon cycling and sequestration in the western tropical North Atlantic (WTNA) is influenced by the Amazon River through its impact on pelagic ecosystem dynamics and 2) the sensitivity of this ecosystem to anthropogenic climate change. PIRANA revealed the importance of both riverine and atmospheric inputs for driving the high productivity of the WTNA through N2-fixation, and demonstrated the significance of the region to basin-wide biogeochemistry and C cycling. ANACONDAS will now focus on what drives phytoplankton community succession through the plume, light and nutrient requirements, factors limiting productivity, and the fate of production. These components are critical to understand the role of the plume in the regional C cycle, and to predict its response to climate variability and change.

The NSF-funded ANACONDAS project will also serve as a platform for additional measurements supported by the Gordon and Betty Moore Foundation's Marine Microbiology Initiative. ROCA (River-Ocean Continuum of the Amazon) brings additional focus on marine microbial community structure and activities, along with high-resolution measurements of organic matter along the river-ocean continuum.

ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses **ROCA:** River Ocean Continuum of the Amazon

The project is funded by NSF-OCE-0934095 and NSF-OCE-0934036: Collaborative Research: ETBC: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses and by the Gordon and Betty Moore Foundation through GBMF-MMI-2293: River Ocean Continuum of the Amazon.

Planned Cruise Sampling

Water Column Characterization (hydrographic sampling with CTD/Rosette):

Nutrient (NO2, NO2+NO3, PO4, SiO4) concentrations Chlorophyll a and pigments concentrations Inorganic carbon (discrete DIC, ALK, underway pCO2) Organic carbon, nitrogen, phosphorus Phytoplankton and Diazotroph Abundance (using rosette and also small nets to collect) Carbon and Nitrogen Fixation by plankton Kinetic and Physiological Measurements of phytoplankton Stable Isotopic Measurements of particulate material Microbial heterotrophy Microbial community structure and gene expression Organic carbon and biomarker characterization

MOCNESS tows for zooplankton

Zooplankton collection for abundance and biomass Zooplankton grazing and POC flux measurements

Multicorer for deep sea sediment analyses

Solid phase analysis Pore water chemistry Isotopic composition (Pb, Th, C)

Other instrumentation over the side:

The in-water light field will be characterized with a free-falling 14 channel spectroradiometer Two "Carbon Explorers" - autonomous Sounding Oceanographic Lagrangian Observer profilers Sediment Trap Studies (using 48h deployments of floating Particle Interceptor Traps; PITs) Surface water pumps - directly bring large volumes of surface water to the deck of the ship for processing.

Shipboard Instrumentation:

ADCP 75 kHz Bathymetry System 12 kHz Bathymetry System 3.5 kHz Deionized Water System Fume Hood HiSeasNet Multibeam Uncontaminated Seawater System CTD/Water Sampling: 911+ Rosette 24-position, 10-liter bottle Rosette with dual T/C sensors Biospherical underwater PAR (1000m depth limit) SBE43 oxygen sensor Wet Labs C*Star transmissometer (660nm wavelength) Wet Labs ECO-AFL fluorometer Dissolved Oxygen Titration System (Portable modified Winkler titration system)

[table of contents | back to top]

Program Information

Integrated Marine Biogeochemistry and Ecosystem Research -US (IMBER-US)

Website: http://www.imber.info/

Coverage: global

The BCO-DMO database includes data from IMBER endorsed projects lead by US funded investigators. There is no dedicated US IMBER project or data management office. Those functions are provided by US-OCB and BCO-DMO respectively.

The information in this program description pertains to the Internationally coordinated IMBER research program. The projects contributing data to the BCO-DMO database are those funded by US NSF only. The full IMBER data catalog is hosted at the Global Change Master Directory (GCMD).

IMBER Data Portal: The IMBER project has chosen to create a metadata portal hosted by the NASA's Global Change Master Directory (GCMD). The GCMD IMBER data catalog provides an overview of all IMBER endorsed and related projects and links to datasets, and can be found at URL http://gcmd.nasa.gov/portals/imber/.

IMBER research will seek to identify the mechanisms by which marine life influences marine biogeochemical cycles, and how these, in turn, influence marine ecosystems. Central to the IMBER goal is the development of a predictive understanding of how marine biogeochemical cycles and ecosystems respond to complex forcings, such as large-scale climatic variations, changing physical dynamics, carbon cycle chemistry and nutrient fluxes, and the impacts of marine harvesting. Changes in marine biogeochemical cycles and ecosystems due to global change will also have consequences for the broader Earth System. An even greater challenge will be drawing together the natural and social science communities to study some of the key impacts and feedbacks between the marine and human systems.

To address the IMBER goal, four scientific themes, each including several issues, have been identified for the IMBER project: Theme 1 - Interactions between Biogeochemical Cycles and Marine Food Webs; Theme 2 - Sensitivity to Global Change: How will key marine biogeochemical cycles, ecosystems and their interactions, respond to global change?; Theme 3 - Feedback to the Earth System: What are the roles of the ocean biogeochemistry and ecosystems in regulating climate?; and Theme 4 - Responses of Society: What are the relationships between marine biogeochemical cycles, ecosystems, and the human system?

Ocean Carbon and Biogeochemistry (OCB)

Website: http://us-ocb.org/

Coverage: Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO2 and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

Emerging Topics in Biogeochemical Cycles (ETBC)

Website: http://www.nsf.gov/pubs/2007/nsf07049/nsf07049.jsp

Coverage: global

The original call for proposals for Emerging Topics in Biogeochemical Cycles (ETBC) was issued in September 2007 by the US NSF Directorate for Geosciences (NSF 07-049).

The Geosciences Directorate (GEO) is substantially augmenting our past funding sources to explicitly support emerging areas of interdisciplinary research. We seek to foster transformational advances in our quantitative or mechanistic understanding of biogeochemical cycles that integrate physical-chemical-biological processes over the range of temporal and/or spatial scales in Earthâ€[™]s environments. We encourage submission of proposals that address emerging topics in biogeochemical cycles, the water cycle or their coupling, across the interfaces of atmosphere, land, and oceans. Proposals must cross the disciplinary boundaries of two or more divisions in Geosciences (e.g. ATM, EAR, OCE) or of at least one division in Geosciences and a division in another NSF directorate.

Although funding programmatic disciplines continues to provide a robust and adaptable framework to build our understanding of the geosciences and the earth as a system, there are classes of emerging and challenging problems that require integration of concepts and observations across diverse fields. Our goal is to enhance such integration. Successful proposals need to develop intellectual excitement in the participating disciplinary communities. Also encouraged are proposals that have broader educational, diversity, societal, or infrastructure impacts that capitalize on this interdisciplinary opportunity.

Website: https://www.moore.org/initiative-strategy-detail?initiativeId=marine-microbiology-initiative

A Gordon and Betty Moore Foundation Program.

Forging a new paradigm in marine microbial ecology:

Microbes in the ocean produce half of the oxygen on the planet and remove vast amounts of carbon dioxide, a greenhouse gas, from the atmosphere. Yet, we have known surprisingly little about these microscopic organisms. As we discover answers to some long-standing puzzles about the roles that marine microorganisms play in supporting the ocean's food webs and driving global elemental cycles, we realized that we still need to learn much more about what these organisms do and how they do it—including how they evolved and contribute to our ocean's health and productivity.

The Marine Microbiology Initiative seeks to gain a comprehensive understanding of marine microbial communities, including their diversity, functions and behaviors; their ecological roles; and their origins and evolution. Our focus has been to enable researchers to uncover the principles that govern the interactions among microbes and that govern microbially mediated nutrient flow in the sea. To address these opportunities, we support leaders in the field through investigator awards, multidisciplinary team research projects, and efforts to create resources of broad use to the research community. We also support development of new instrumentation, tools, technologies and genetic approaches.

Through the efforts of many scientists from around the world, the initiative has been catalyzing new science through advances in methods and technology, and to reduce interdisciplinary barriers slowing progress. With our support, researchers are quantifying nutrient pools in the ocean, deciphering the genetic and biochemical bases of microbial metabolism, and understanding how microbes interact with one another. The initiative has five grant portfolios:

Individual investigator awards for current and emerging leaders in the field.

Multidisciplinary projects that support collaboration across disciplines.

New instrumentation, tools and technology that enable scientists to ask new questions in ways previously not possible.

. Community resource efforts that fund the creation and sharing of data and the development of tools, methods and infrastructure of widespread utility.

Projects that advance genetic tools to enable development of experimental model systems in marine microbial ecology.

We also bring together scientists to discuss timely subjects and to facilitate scientific exchange.

Our path to marine microbial ecology was a confluence of new technology that could accelerate science and an opportunity to support a field that was not well funded relative to potential impact. Around the time we began this work in 2004, the life sciences were entering a new era of DNA sequencing and genomics, expanding possibilities for scientific research – including the nascent field of marine microbial ecology. Through conversations with pioneers inside and outside the field, an opportunity was identified: to apply these new sequencing tools to advance knowledge of marine microbial communities and reveal how they support and influence ocean systems.

After many years of success, we will wind down this effort and close the initiative in 2021. We will have invested more than \$250 million over 17 years to deepen understanding of the diversity, ecological activities and evolution of marine microbial communities. Thanks to the work of hundreds of scientists and others involved with the initiative, the goals have been achieved and the field has been profoundly enriched; it is now positioned to address new scientific questions using innovative technologies and methods.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-0934095</u>
Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)	<u>GBMF2293</u>

[table of contents | back to top]