

Plankton tow log from Alvin dives from R/V Atlantis AT15-59 in the Pacific Ocean off Costa Rica from January 2010 (Seep Carbonate Ecology CROCKS II project)

Website: <https://www.bco-dmo.org/dataset/3617>

Version:

Version Date: 2012-01-30

Project

» [Short-term colonization processes at Costa Rica methane seeps](#) (Seep Carbonate Ecology CROCKS II)

Contributors	Affiliation	Role
Levin, Lisa A.	University of California-San Diego (UCSD-SIO)	Chief Scientist
Orphan, Victoria J.	California Institute of Technology (Caltech)	Co-Chief Scientist
Rathburn, Anthony E	Indiana State University (ISU)	Co-Chief Scientist
Rouse, Gregory	University of California-San Diego (UCSD-SIO)	Co-Chief Scientist
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Atlantis/Alvin cruise off Costa Rica

The primary goal of the cruise was to recover biological experiments deployed at active and inactive seep areas during Feb./March 2009. We successfully recovered 23/24 experimental units deployed on Mound 12, Costa Rica. One was simply missing on the sea floor. By using a gear elevator each day to maximize sample collection (and reserve ALVIN basket space for experiments) we were able to recover our experiments rapidly. This left us time for exploration of unusual biological communities at Jaco Scar.

Methods & Sampling

Equipment used: standard Alvin tubecores; Ocean Instruments multicorer; Seabird CTD with O2 sensor.

Where indicated, sediment samples from tube cores were sectioned vertically and preserved in buffered formaldehyde (standard procedures) or frozen at -80°C.

Colonization substrates had macrofauna (>0.3 mm) and meiofauna (>0.42 microns) removed and preserved. Representative individuals were frozen for subsequent stable isotope analyses.

[[table of contents](#) | [back to top](#)]

Data Files

File

planktontow_log.csv(Comma Separated Values (.csv), 640 bytes)
MD5:52f71ec15fbef7e694e29863d4bb31d5

Primary data file for dataset ID 3617

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
tow	tow number	integer
year	year of sampling	integer
month	month of sampling	1 to 12
day	day of sampling	1 to 31
yrday_gmt	GMT day and decimal time; as 326.5 for the 326th day of the year or November 22 at 1200 hours (noon)	decimal date
time	GMT time	hhmm
lat_start	latitude at start of tow; North is positive	decimal degrees
lon_start	longitude at start of tow; East is positive	decimal degrees
lat_end	latitude at end of tow; North is positive	decimal degrees
lon_end	longitude at end of tow; East is positive	decimal degrees
duration	duration of tow	minutes
mesh	mesh size	microns
speed_kts	speed of ship during tow	knots
comments_speed	comments pertaining to ship's speed	text
wire_out	meters of wire out	meters
comments	comments	text

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Plankton Net
Generic Instrument Name	Plankton Net
Generic Instrument Description	A Plankton Net is a generic term for a sampling net that is used to collect plankton. It is used only when detailed instrument documentation is not available.

[[table of contents](#) | [back to top](#)]

Deployments

AT15-59

Website	https://www.bco-dmo.org/deployment/58765
Platform	R/V Atlantis
Start Date	2010-01-06
End Date	2010-01-13
Description	Costa Rica seafloor methane seeps 8 deg 55 N 84 depth 990m. Included Alvin dives 4586-4591. The primary goal of the cruise was to recover biological experiments deployed at active and inactive seep areas during Feb./March 2009. We successfully recovered 23/24 experimental units deployed on Mound 12, Costa Rica. One was simply missing on the sea floor. By using a gear elevator each day to maximize sample collection (and reserve ALVIN basket space for experiments) we were able to recover our experiments rapidly. This left us time for exploration of unusual biological communities at Jaco Scar. Cruise information and original data are available from the NSF R2R data catalog. Cruise dates changed Feb 2015 to match WHOI ship schedule and R2R.

[[table of contents](#) | [back to top](#)]

Project Information

Short-term colonization processes at Costa Rica methane seeps (Seep Carbonate Ecology CROCKS II)

Coverage: Costa Rica seafloor methane seeps 8 deg 55 N 84 deg 18 W depth 990m

This RAPID project will conduct 5 submersible or ROV dives to collect a series of colonization experiments deployed in March 2009 on Mound 12 off Costa Rica (997 m). These experiments were deployed opportunistically, and to optimize the information that could be obtained, the PIs needed to recover them within a 12 month time frame. Early colonization of rock, wood, shell and tube substrates will be studied. The microbes, foraminiferans and metazoans present after 6-12 mo will be compared to those colonizing similar experiments to be deployed at Hydrate Ridge, where seeps occur within an oxygen minimum zone. The overall project goal is to integrate physical, geological, chemical and biological data to develop a holistic view of the influence of seep-generated carbonate hard-ground ecosystems on margins.

The objectives of the research are to (a) Compare colonizers at seeps off Costa Rica and Hydrate Ridge to assess the importance of different oxygen regimes in the development of anaerobic methane oxidation, sulfide oxidizers and other microbial metabolisms on hard substrates, and to evaluate their roles in driving protozoan and metazoan succession at methane seeps. (b) Deploy a suite of biotic and abiotic substrates to distinguish the specific roles of carbonate substrate from those of other hard substrates (wood, clam and mussel shells, worm tubes) available. (c) Explore the similarity of vent and seep colonization processes by comparing colonization at the Costa Rica seeps, where vent species are common, to the Hydrate Ridge seeps, where they are not. (d) Determine whether there are diagnosable biogeographic isotope or other biomarker signatures from newly recruited Costa Rica microbial, foraminiferal and animal populations at active vs. inactive seeps, and whether these differ from those at Hydrate Ridge.

This research will involve international participation from Costa Rican scientists at the Univ. of Costa Rica.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0825791
NSF Division of Ocean Sciences (NSF OCE)	OCE-0826254
NSF Division of Ocean Sciences (NSF OCE)	OCE-0939559
NSF Division of Ocean Sciences (NSF OCE)	OCE-0825436
NSF Division of Ocean Sciences (NSF OCE)	OCE-0939232
NSF Division of Ocean Sciences (NSF OCE)	OCE-0939557

[[table of contents](#) | [back to top](#)]