Temperature Data along the southern California coast around 32N 117W and northwest coast of Mexico from 2003-2009 (Bivalve Connectivity project)

Website: https://www.bco-dmo.org/dataset/3637 Version: 02 April 2012 Version Date: 2012-04-02

Project

» Temporal and spatial scales of variability in bivalve connectivity (Bivalve Connectivity)

Contributors	Affiliation	Role
<u>Levin, Lisa A.</u>	University of California-San Diego (UCSD-SIO)	Principal Investigator
<u>Muccino, Julia C.</u>	Arizona State University (ASU)	Co-Principal Investigator
Rasmussen, Linda L.	University of California-San Diego (UCSD-SIO)	Co-Principal Investigator
<u>Gonzalez, Jennifer</u>	University of California-San Diego (UCSD-SIO)	Contact
<u>Gegg, Stephen R.</u>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
 - <u>Methods & Sampling</u>
 - <u>Data Processing Description</u>
- Data Files
- Parameters
- Instruments
- <u>Deployments</u>
- <u>Project Information</u>
- <u>Funding</u>

Dataset Description

Multi year temperature data in degs C May/2003 - Nov/2009 From coastal deployment on moorings approx. 1-2 meters below surface

Methods & Sampling

Hobo temperature loggers deployed on anchored moorings at 1-3 m water depth in inshore locations having approximately 10 m total water depth.

Deployments generally occurred for 2-4 weeks in May and November between 2003 and 2009.

Data Processing Description

No Processing of the temperature data was done.

BCO-DMO Processing

Generated from original file "BC_TemperatureData.xls" contributed by Lisa Levin

- Simple awk (BC_Temperature_Transpose.awk) written to reorganize the data by Site
- Site Id, Site Abbreviation and Site Lat/Lon added
- Date converted to YYYYMMDD
- Time converted to HHMMSS

Data Files

File
Temperature_Data.csv(Comma Separated Values (.csv), 7.92 MB) MD5:a7a3c71cef3715267bfa6cbc8e6ec8b7
Primary data file for dataset ID 3637

[table of contents | back to top]

Parameters

Parameter	Description	Units
Latitude	latitude in decimal degrees (positive is North)	decimal degrees
Longitude	longitude in decimal degrees (negative is West)	decimal degrees
Abbr	Site Abbreviation	text
Site	Site Name	dimensionless
Date	Date (GMT -8)	YYYYMMDD
Temp	Temperature	degrees Celsius
Time	Time (GMT -8)	HHMMSS

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	Onset Pro v2 temperature logger
Generic Instrument Name	Onset HOBO Pro v2 temperature logger
Generic Instrument	The HOBO Water Temp Pro v2 temperature logger, manufactured by Onset Computer Corporation, has 12-bit resolution and a precision sensor for ±0.2°C accuracy over a wide temperature range. It is designed for extended deployment in fresh or salt water. Operation range: -40° to 70°C (-40° to 158°F) in air; maximum sustained temperature of 50°C (122°F) in water Accuracy: 0.2°C over 0° to 50°C (0.36°F over 32° to 122°F) Resolution: 0.02°C at 25°C (0.04°F at 77°F) Response time: (90%) 5 minutes in water; 12 minutes in air moving 2 m/sec (typical) Stability (drift): 0.1°C (0.18°F) per year Real-time clock: ± 1 minute per month 0° to 50°C (32° to 122°F) Additional information (http://www.onsetcomp.com/) Onset Computer Corporation 470 MacArthur Blvd Bourne, MA 02532

[table of contents | back to top]

Deployments

BC_May2003-Nov2009

Website	https://www.bco-dmo.org/deployment/58788
Platform	Bivalve Connectivity
Start Date	2003-05-01
End Date	2009-11-30
Description	Life history data collection and Inshore larval connectivity studies May/2003 - Nov/2009 A series of sites re-visited on separate occasions from May/2003 - Nov/2009 General Location of the Sites: Southern California Coast around 32N, 117W Northwest Coast of Mexico - Bajamar, MX and Punta Morro, MX

[table of contents | back to top]

Project Information

Temporal and spatial scales of variability in bivalve connectivity (Bivalve Connectivity)

Website: http://levin.ucsd.edu/

Coverage: Southern California Coast around 32N, 117W, NW Mexico

Temporal and spatial scales of variability in bivalve connectivity

Larval connectivity, the extent to which sub-populations exchange larvae, has emerged as a fundamental concept within the diverse arenas of population ecology, biotic resource management, biodiversity conservation, invasive species control, and habitat restoration. However, determining dispersal trajectories of larvae and their scales of variability remain a major challenge. Project investigators Levin, Muccino and Rasmussen will use an integration of prospective modeling and retrospective (elemental fingerprinting) approaches to assess variability in larval connectivity and its demographic consequences for mytilid mussel populations in southern California. This project builds on initial studies of mytilid connectivity to address in greater depth and with a more strongly coupled physical/biological approach, questions of variability and its causes.

These collaborating scientists will address hypotheses concerning the spatial and temporal scales of connectivity for mussels, Mytilus californianus and Mytilus galloprovincialis, examining their consistency among sites and species. This will be accomplished through (a) larval out-planting at 18 locations in San Diego County several times a year to generate reference signatures for trace elemental fingerprinting, (b) collection of recruits and elemental analyses of their larval shells to determine sites and regions of origins, and (c) high frequency data collection at 2 bay locations for M. galloprovincialis and 2 open coast locations for M. californianus to carry out weekly analysis of recruitment variability, its link to chemical signals and recruit origins, and for collection of demographic data (size-specific survivorship, growth and fecundity).

Through numerical dispersal simulations of the coastal ocean and bays, and subsequent comparisons to fingerprinting based assessment of recruit origins, they will examine the roles of circulation, local vs. remote forcing, bay-ocean interaction, episodic events, and larval attributes (vertical behavior, release times, planktonic duration) in defining the variability of connectivity. Demographic data will be combined with connectivity data to model the population and meta-population level fitness consequences of observed mytilid connectivity patterns.

The results of connectivity studies have direct applicability to conservation and management of commercial and natural bivalve populations, and given the key structural role of mussels, to the conservation of rocky shore habitats.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-0648656</u>
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-0648193</u>

[table of contents | back to top]