

# Dissolved inorganic carbon from R/V Thomas G. Thompson and R/V Kilo Moana cruises TN277, KM1301 in the Eastern North Pacific Ocean from 2012-2013 (POWOW project)

**Website:** <https://www.bco-dmo.org/dataset/3752>

**Data Type:** Cruise Results

**Version:** 2

**Version Date:** 2014-05-13

## Project

» [Seasonal and decadal changes in temperature drive Prochlorococcus ecotype distribution patterns](#)

(POWOW)

Contributors	Affiliation	Role
<a href="#">Johnson, Zackary I.</a>	Duke University	Principal Investigator
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Dissolved inorganic carbon from R/V Thomas G. Thompson and R/V Kilo Moana cruises TN277, KM1301 in the Eastern North Pacific Ocean from 2012-2013 (POWOW project)

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:41.5004 E:-120.6976 S:21.3428 W:-160.6166

**Temporal Extent:** 2012-03-01 - 2013-02-06

---

## Dataset Description

Dissolved inorganic carbon from samples collected during the POWOW1 (TN277) and POWOW2 (KM1301) cruises.

## Methods & Sampling

DIC was measured on mercuric chloride poisoned samples by acidification and subsequent quantification of released CO<sub>2</sub> using a CO<sub>2</sub> detector (Li-Cor 7000). DIC samples were collected following recommended procedures (Dickson et al., 2007) and measurements were calibrated against Certified Reference Materials provided by Dr. A. G. Dickson at Scripps Institution of Oceanography (SIO), University of California, San Diego (UCSD).

## Data Processing Description

### BCO-DMO edits made:

- Parameter names have been changed to conform to BCO-DMO conventions.
- month\_utc, day\_utc, year, and time\_utc were added, based on the original ISO\_DateTime\_UTC field.
- Rosette bottle numbers were added from the CTD cast sheets.
- Replaced 'NaN' with 'nd' to indicate 'no data'.

NOTE: During cast CTD01 of POWOW1, all bottles were fired at 25m depth. The bottle numbers displayed in the data for CTD01 of POWOW1 are those bottles from which DIC was measured.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>DIC.csv</b> (Comma Separated Values (.csv), 78.02 KB) MD5:614180e0b34da4f0120f6d5dfffb17e8b Primary data file for dataset ID 3752

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Dickson, A.G., Sabine, C.L. and Christian, J.R. (Eds.) 2007. Guide to best practices for ocean CO2 measurements. PICES Special Publication 3, 191 pp. ISBN: 1-897176-07-4. URL: [https://www.nodc.noaa.gov/ocads/oceans/Handbook\\_2007.html](https://www.nodc.noaa.gov/ocads/oceans/Handbook_2007.html) <https://hdl.handle.net/11329/249>  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
cruise_name	Cruise identifier (POWOW1 = TN277 = R/V Thomas G. Thompson cruise 277; POWOW2 = KM1301 = R/V Kilo Moana cruise 1301).	text
cast	Consecutive CTD cast number. CTD numbers are unique and sequential across stations.	unitless
lat	Latitude at start of CTD cast. Positive = North.	decimal degrees
lon	Longitude at start of CTD cast. Positive = East.	decimal degrees
depth_w	Depth of the water (bottom depth) at sampling station.	meters
month_utc	2-digit month of year, UTC.	mm (01 to 12)
day_utc	2-digit day of month, UTC.	dd (01 to 31)
year	4-digit year. in YYYY format	unitless
time_utc	Time (UTC) at start of sample collection, 24-hour clock.	HHMM.mm
ISO_DateTime_UTC	Date/Time (UTC) ISO8601 formatted. T indicates start of time string; Z indicates UTC.	YYYY-mm-ddTHH:MM:SS.ssZ
depth	Sample depth.	meters
bot	Rosette position of the bottle.	unitless
DIC	Dissolved inorganic carbon, uM Carbon.	uM

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	LI-COR LI-7000 Gas Analyzer
<b>Generic Instrument Name</b>	LI-COR LI-7000 Gas Analyzer
<b>Dataset-specific Description</b>	Released CO2 was quantified using a Li-Cor 7000 CO2 detector.
<b>Generic Instrument Description</b>	The LI-7000 CO2/H2O Gas Analyzer is a high performance, dual cell, differential gas analyzer. It was designed to expand on the capabilities of the LI-6262 CO2/ H2O Gas Analyzer. A dichroic beam splitter at the end of the optical path provides radiation to two separate detectors, one filtered to detect radiation absorption of CO2 and the other to detect absorption by H2O. The two separate detectors measure infrared absorption by CO2 and H2O in the same gas stream. The LI-7000 CO2/ H2O Gas Analyzer is a differential analyzer, in which a known concentration (which can be zero) gas is put in the reference cell, and an unknown gas is put in the sample cell.

[ [table of contents](#) | [back to top](#) ]

## Deployments

### TN277

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/58867">https://www.bco-dmo.org/deployment/58867</a>
<b>Platform</b>	R/V Thomas G. Thompson
<b>Report</b>	<a href="http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW1-cruise_report.pdf">http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW1-cruise_report.pdf</a>
<b>Start Date</b>	2012-02-29
<b>End Date</b>	2012-03-11
<b>Description</b>	The POWOW #1 cruise was a trip of opportunity to sample along temperature gradients and test out new protocols. The primary goal of this cruise was to measure the abundance, diversity and activity of Prochlorococcus and associated bacterial and viral communities across temperature (and other environmental) gradients to understand how climate change may impact ocean ecology and biogeochemistry. There are many additional scientific and broader impact goals including characterizing oxidative stress and investigating nitrogen uptake/utilization molecular diversity. Cruise information and original data are available from the NSF R2R data catalog.

### KM1301

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/505095">https://www.bco-dmo.org/deployment/505095</a>
<b>Platform</b>	R/V Kilo Moana
<b>Report</b>	<a href="http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW2-cruise_report.pdf">http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW2-cruise_report.pdf</a>
<b>Start Date</b>	2013-01-10
<b>End Date</b>	2013-02-08
<b>Description</b>	From the cruise report: The POWOW #2 cruise was the second in a series of cruises to study the influence of temperature and other environmental variables on <i>Prochlorococcus</i> , its viruses and other members of the microbial community. The primary goal of this cruise was to measure the abundance, diversity and activity of <i>Prochlorococcus</i> and associated bacterial and viral communities across temperature (and other environmental) gradients to understand how climate change may impact ocean ecology and biogeochemistry. Cruise information and original data are available from the NSF R2R data catalog.

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Seasonal and decadal changes in temperature drive *Prochlorococcus* ecotype distribution patterns (POWOW)

**Website:** <http://oceanography.ml.duke.edu/johnson/research/powow/>

**Coverage:** Eastern North Pacific Ocean

Project also known as '*Prochlorococcus* Of Warming Ocean Waters' (POWOW).

The two numerically-dominant ecotypes of the marine cyanobacterium *Prochlorococcus* partition the surface ocean niche latitudinally, with ecotype eMIT9312 dominant in the 30 degree N to 30 degree S region and eMED4 dominant at higher latitudes. These ecotypes may account for 25-50% of primary production in open ocean ecosystems, but this percentage is dependent on which ecotype dominates. The relative abundance of the two ecotypes follows a log-linear relationship with temperature, with the transition from eMIT9312 to eMED4 occurring at approx. 18 degrees C. From these descriptive data, it has been hypothesized that temperature is the primary driver of relative abundance. Their contribution to net primary production, however, appears to be independent of temperature, suggesting temperature regulates ecotype dominance through photosynthesis-independent mechanisms.

To test these hypotheses, the PIs are undertaking a series of field and lab studies to investigate the effect of temperature change on the distribution of these ecotypes. Two cruises in the North Pacific will trace the transitions from eMIT9312- to eMED4-dominated regions, with one cruise during the winter and the other during summer. They have hypothesized that the ratio of ecotype abundance will move latitudinally with the seasonal shift in temperature gradient: migration of the 18 degrees C isotherm northward in the summer will be matched by a similar migration of the 1:1 ecotype transition point. Multiple crossings of the 18 degrees C isotherm are proposed, and the summer cruise will also follow the isotherm to the Western US coast to gain insight on physical and geochemical influences. Environmental variables such as nutrient concentrations, light/mixing depths, and virus /grazing based mortality, which may impinge on the relationship between temperature and ecotype ratio, will be assessed through a series of multivariate analyses of the collected suite of physical, chemical and biological data. Seasonal comparisons will be complemented with on-deck incubations and lab competition assays (using existing and new isolates) that will establish, for the first time, how fitness coefficients of these ecotypes relate to temperature. As latitudinal shifts in temperature gradient and migration of ecotypes during seasonal warming likely share common features with high latitude warming as a consequence of climate change, the investigator's analyses will contribute important biological parameters (e.g., abundances, production rates, temperature change coefficients) for modeling biological and biogeochemical responses to climate change. This research will be integrated with that of committed collaborators, generating data sufficient for ecosystem-scale characterizations of the contributions of temperature (relative to other forcing factors) in constraining the range and seasonal migration of these

numerically dominant marine phototrophs.

**Publications produced as result of this research:**

Rowe, J.M., DeBruyn, J.M., Poorvin, L., LeClerc, G.R., Johnson, Z.I., Zinser, E.R., and Wilhelm, S.W. 2012. Viral and bacterial abundance and production in the Western Pacific Ocean and the relation to other oceanic realms. *FEMS Microbiology Ecology*, 72, p. 359. DOI: [10.1111/j.1574-6941.2011.01223.x](https://doi.org/10.1111/j.1574-6941.2011.01223.x)

Morris, J.J., Lenski, R.E. and E.R. Zinser. 2012. The Black Queen Hypothesis: Evolution of Dependencies through Adaptive Gene Loss. *mBio*, 3, p. e00036-12. DOI: [10.1128/mBio.00036-12](https://doi.org/10.1128/mBio.00036-12)

Morris, J.J., Johnson, Z.I., Szul, M.J., Keller, M., and Zinser, E.R. 2011. Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS One*, 6(2), p. 16805. DOI:[10.1371/journal.pone.0016805](https://doi.org/10.1371/journal.pone.0016805)

Ringuet, S., Sassano, L., and Johnson, Z.I. 2011. A suite of microplate reader-based colorimetric methods to quantify ammonium, nitrate, orthophosphate and silicate concentrations for aquatic nutrient monitoring. *Journal of Environmental Monitoring*. DOI:[10.1039/C0EM00290A](https://doi.org/10.1039/C0EM00290A)

Ritchie, A.E. and Johnson, Z.I. 2012. Abundance and genetic diversity of aerobic anoxygenic phototrophic bacteria of coastal regions of the Pacific Ocean. *Applied and Environmental Microbiology*, 78, p. 2858. DOI: [10.1128/AEM.06268-11](https://doi.org/10.1128/AEM.06268-11)

[ [table of contents](#) | [back to top](#) ]

---

**Funding**

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1031064</a>

[ [table of contents](#) | [back to top](#) ]