Ammonium (NH4) data from CTD rosette bottles from R/V Thomas G. Thompson and R/V Kilo Moana cruises TN277, KM1301, KM1312 in the Eastern North Pacific Ocean from 2012-2013 (POWOW project)

Website: https://www.bco-dmo.org/dataset/3754

Data Type: Cruise Results

Version: 2

Version Date: 2014-07-10

Proiect

» <u>Seasonal and decadal changes in temperature drive Prochlorococcus ecotype distribution patterns</u> (POWOW)

Contributors	Affiliation	Role
Johnson, Zackary I.	Duke University	Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Ammonium (NH4) data from CTD rosette bottles from R/V Thomas G. Thompson and R/V Kilo Moana cruises TN277, KM1301, KM1312 in the Eastern North Pacific Ocean from 2012-2013 (POWOW project)

Table of Contents

- Coverage
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Related Publications
- Parameters
- <u>Instruments</u>
- Deployments
- Project Information
- Funding

Coverage

Spatial Extent: N:46.6888 E:-118.316 S:21.3428 W:-160.6166

Temporal Extent: 2012-03-01 - 2013-07-27

Dataset Description

Ammonium (NH4) concentration in nM from 3 POWOW cruises (TN277, KM1301, and KM1312). Samples that had a mean concentration (mean of replicated samples) below the nominal detection limit are reported as zero.

Also see related inorganic nutrient data from this project.

Nitrite and Nitrate (NO2, NO3) https://www.bco-dmo.org/dataset/3755

Phosphate and Silicate (PO4, SiOH4) https://www.bco-dmo.org/dataset/3755

Methods & Sampling

Ammonium was measured on fresh, unfiltered samples following Holmes et al. 1999. Certified reference materials were used to verify protocols (Inorganic Ventures: QCP-NT, QCP-NUT-1, CGSI1-1). The detection

limits of NH4 were:
9 nM for POWOW1 samples
5 nM for POWOW2 and POWOW3 samples

Data Processing Description

BCO-DMO edits made:

- Parameter names have been changed to conform to BCO-DMO conventions.
- month utc, day utc, year, and time utc were added, based on the original ISO DateTime UTC field.
- Rosette bottle numbers were added from the CTD cast sheets for POWOW1 data.
- 'NaN' was replaced with 'nd' to indicate 'no data'.

[table of contents | back to top]

Data Files

File

ammonium.csv(Comma Separated Values (.csv), 159.79 KB)

MD5:f75bfea19ef4b949f2bebf5ef3f741da

Primary data file for dataset ID 3754

[table of contents | back to top]

Related Publications

Holmes, R. M., Aminot, A., Kerouel, R., Hooker, B. A., & Peterson, B. J. (1999). A simple and precise method for measuring ammonium in marine and freshwater ecosystems. Canadian Journal of Fisheries and Aquatic Sciences, 56(10), 1801-1808. doi:10.1139/f99-128 https://doi.org/10.1139/cjfas-56-10-1801 *Methods*

[table of contents | back to top]

Parameters

Parameter	Description	Units
cruise_name	Cruise identifier (POWOW1 = TN277 = R/V Thomas G. Thompson cruise 277; POWOW2 = KM1301 = R/V Kilo Moana cruise 1301; POWOW3 = KM1312 = R/V Kilo Moana cruise 1312).	text
cast	Consecutive CTD cast number. CTD numbers are unique and sequential across stations.	unitless
lat	Latitude at start of CTD cast. Positive = North.	decimal degrees
lon	Longitude at start of CTD cast. Positive = East.	decimal degrees
depth_w	Depth of the water (bottom depth) at sampling station.	meters
month_utc	2-digit month of year, UTC.	mm (01 to 12)
day_utc	2-digit day of month, UTC.	dd (01 to 31)
year	4-digit year. in YYYY format	unitless
time_utc	Time (UTC) at start of sample collection, 24-hour clock.	HHMM.mm
ISO_DateTime_UTC	Date/Time (UTC) ISO8601 formatted. T indicates start of time string; Z indicates UTC.	YYYY-mm- ddTHH:MM:SS.ssZ
depth	Sample depth.	meters
bot	Rosette position of the bottle.	unitless
NH4	Ammonium (NH4) concentration in nM.	nanomolar (nM)

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	Niskin bottle
Generic Instrument Name	Niskin bottle
	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

[table of contents | back to top]

Deployments

TN277

Website	https://www.bco-dmo.org/deployment/58867	
Platform	R/V Thomas G. Thompson	
Report	http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW1-cruise_report.pdf	
Start Date	2012-02-29	
End Date	2012-03-11	
Description	The POWOW#1 cruise was a trip of opportunity to sample along temperature gradients and test out new protocols. The primary goal of this cruise was to measure the abundance, diversity and activity of Prochlorococcus and associated bacterial and viral communities across temperature (and other environmental) gradients to understand how climate change may impact ocean ecology and biogeochemistry. There are many additional scientific and broader impact goals including characterizing oxidative stress and investigating nitrogen uptake/utilization molecular diversity. Cruise information and original data are available from the NSF R2R data catalog.	

KM1301

	4.1701	
Website	https://www.bco-dmo.org/deployment/505095	
Platform	R/V Kilo Moana	
Report	http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW2-cruise_report.pdf	
Start Date	2013-01-10	
End Date	2013-02-08	
Description	From the cruise report: The POWOW#2 cruise was the second in a series of cruises to study the influence of temperature and other environmental variables on Prochlorococcus, its viruses and other members of the microbial community. The primary goal of this cruise was to measure the abundance, diversity and activity of Prochlorococcus and associated bacterial and viral communities across temperature (and other environmental) gradients to understand how climate change may impact ocean ecology and biogeochemistry. Cruise information and original data are available from the NSF R2R data catalog.	

KM1312

Website	https://www.bco-dmo.org/deployment/515629	
Platform	R/V Kilo Moana	
Report	http://dmoserv3.whoi.edu/data_docs/POWOW/POWOW3-cruise_report.pdf	
Start Date	2013-07-01	
End Date	2013-07-28	
Description	From the cruise report: The POWOW#3 cruise was the third in a series of cruises to study the influence of temperature and other environmental variables on Prochlorococcus, its viruses and other members of the microbial community in the Northern Pacific Ocean. The primary goal of this cruise was to measure the abundance, diversity and activity of Prochlorococcus and associated bacterial and viral communities across temperature (and other environmental) gradients to understand how climate change may impact ocean ecology and biogeochemistry. There are many additional scientific and broader impact goals including characterizing oxidative stress and investigating nitrogen uptake/utilization molecular diversity. The official title of the project is "Collaborative Research: Seasonal and decadal changes in temperature drive Prochlorococcus ecotype distribution patterns" and it is part of NSF #1031064 (Duke) and 1030518 (UTK). Cruise information and original data are available from the NSF R2R data catalog.	

Project Information

Seasonal and decadal changes in temperature drive Prochlorococcus ecotype distribution patterns (POWOW)

Website: http://oceanography.ml.duke.edu/johnson/research/powow/

Coverage: Eastern North Pacific Ocean

Project also known as 'Prochlorococcus Of Warming Ocean Waters' (POWOW).

The two numerically-dominant ecotypes of the marine cyanobacterium *Prochlorococcus* partition the surface ocean niche latitudinally, with ecotype eMIT9312 dominant in the 30 degree N to 30 degree S region and eMED4 dominant at higher latitudes. These ecotypes may account for 25-50% of primary production in open ocean ecosystems, but this percentage is dependent on which ecotype dominates. The relative abundance of the two ecotypes follows a log-linear relationship with temperature, with the transition from eMIT9312 to eMED4 occurring at approx. 18 degrees C. From these descriptive data, it has been hypothesized that temperature is the primary driver of relative abundance. Their contribution to net primary production, however, appears to be independent of temperature, suggesting temperature regulates ecotype dominance through photosynthesis-independent mechanisms.

To test these hypotheses, the PIs are undertaking a series of field and lab studies to investigate the effect of temperature change on the distribution of these ecotypes. Two cruises in the North Pacific will trace the transitions from eMIT9312- to eMED4-dominated regions, with one cruise during the winter and the other during summer. They have hypothesized that the ratio of ecotype abundance will move latitudinally with the seasonal shift in temperature gradient: migration of the 18 degrees C isotherm northward in the summer will be matched by a similar migration of the 1:1 ecotype transition point. Multiple crossings of the 18 degrees C isotherm are proposed, and the summer cruise will also follow the isotherm to the Western US coast to gain insight on physical and geochemical influences. Environmental variables such as nutrient concentrations, light/mixing depths, and virus /grazing based mortality, which may impinge on the relationship between temperature and ecotype ratio, will be assessed through a series of multivariate analyses of the collected suite of physical, chemical and biological data. Seasonal comparisons will be complemented with on-deck incubations and lab competition assays (using existing and new isolates) that will establish, for the first time, how fitness coefficients of these ecotypes relate to temperature. As latitudinal shifts in temperature gradient and migration of ecotypes during seasonal warming likely share common features with high latitude warming as a consequence of climate change, the investigator's analyses will contribute important biological parameters (e.g., abundances, production rates, temperature change coefficients) for modeling biological and biogeochemical responses to climate change. This research will be integrated with that of committed collaborators, generating data sufficient for ecosystem-scale characterizations of the contributions of temperature (relative to other forcing factors) in constraining the range and seasonal migration of these numerically dominant marine phototrophs.

Publications produced as result of this research:

Rowe, J.M., DeBruyn, J.M., Poorvin, L., LeCleir, G.R., Johnson, Z.I., Zinser, E.R., and Wilhelm, S.W. 2012. Viral and bacterial abundance and production in the Western Pacific Ocean and the relation to other oceanic realms. FEMS Microbiology Ecology, 72, p. 359. DOI: 10.1111/j.1574-6941.2011.01223.x

Morris, J.J., Lenski, R.E. and E.R. Zinser. 2012. The Black Queen Hypothesis: Evolution of Dependencies through Adaptive Gene Loss. mBio, 3, p. e00036-12. DOI: 10.1128/mBio.00036-12

Morris, J.J., Johnson, Z.I., Szul, M.J., Keller, M., and Zinser, E.R. 2011. Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. PLoS One, 6(2), p. 16805. DOI:10.1371/journal.pone.0016805

Ringuet, S., Sassano, L., and Johnson, Z.I. 2011. A suite of microplate reader-based colorimetric methods to quantify ammonium, nitrate, orthophosphate and silicate concentrations for aquatic nutrient monitoring. Journal of Environmental Monitoring. DOI:10.1039/C0EM00290A

Ritchie, A.E. and Johnson, Z.I. 2012. Abundance and genetic diversity of aerobic anoxygenic phototrophic bacteria of coastal regions of the Pacific Ocean. Applied and Environmental Microbiology, 78, p. 2858. DOI: 10.1128/AEM.06268-11

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1031064

[table of contents | back to top]