

# Concentrations of anions, cations, and trace elements/metals from vents near Milos Island, Greece. from May 2012 (Hydrothermal Autotrophic Carbon Fixation project)

Website: <https://www.bco-dmo.org/dataset/474322>

Version: 2014-01-08

## Project

» [Autotrophic carbon fixation at a shallow-water hydrothermal system: Constraining microbial activity, isotopic and geochemical regimes](#) (Hydrothermal Autotrophic Carbon Fixation)

## Program

» [Center for Dark Energy Biosphere Investigations](#) (C-DEBI)

Contributors	Affiliation	Role
<a href="#">Foustoukos, Dionysis</a>	Carnegie Institution for Science (CIS)	Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Dataset Description

Data from samples collected at the shallow-water hydrothermal vent sites at Paleochori Bay, Milos Island, Greece, and includes the chemical composition (anion, cation, trace elements/metals) of vent fluids sampled.

This is also available at [http://people.gl.ciw.edu/dfoustoukos/Site/Data\\_Repository.html](http://people.gl.ciw.edu/dfoustoukos/Site/Data_Repository.html)

## Methods & Sampling

Fluid samples were collected from the water column close to the shallow-water venting sites and from the interstitial waters of all cores at 2 cm intervals. Fluids and gases were also sampled from the subaerial sites. Samples returned to the Geophysical Lab were analyzed to determine dissolved concentrations of: organic acids and a range of major anions/cations species (e.g. SO<sub>4</sub><sup>2-</sup>, PO<sub>4</sub><sup>2-</sup>, NH<sub>3</sub>, NO<sub>3</sub><sup>-</sup>, Cl<sup>-</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, and Mg<sup>2+</sup>) including trace elements and metals. We used ion chromatography (Metrohm "MIC-3 Advanced IC, Metrosep A supp 7-250/16-250; C4- 250 columns). Fluids were analyzed for metal and trace element concentrations at the MC-ICP-MS facility of Prof. Michael Bizimis at the University of South Carolina.

## Relevant References:

Mustafa Yücel, Stefan M. Sievert, Costantino Vetriani, Dionysis I. Foustoukos, Donato Giovannelli, Nadine Le Bris. Eco-geochemical dynamics of a shallow-water hydrothermal vent system at Milos Island, Aegean Sea (Eastern Mediterranean). *Chemical Geology* 356 (2013) 11-20.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>vent_chem.csv</b> (Comma Separated Values (.csv), 3.99 KB) MD5:b47acd181a5ddcf589ce2d12bfb8ecb3 Primary data file for dataset ID 474322

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
date	sampling date	mm/dd/yyyy
year	year of sampling	yyyy
month	month of sampling	1 to 12
day	day of sampling	1 to 31
yrday	yearday of sampling; e.g. 1.5 means January 1 at 1200 hours (noon time)	1 to 365
site	sample location identification	unitless
lat	latitude; north is postive	decimal degrees
lon	longitude; east is postive	decimal degrees
distance	distance from the vent site	centimeters
depth_cm	sampling depth	centimeters
sample_descrip	brief description of the sample name	text
sample_id	sample code	alphanumeric
temp	temperature at sampling location	degrees Celsius
activity	vent or subaerial	text
Li	Lithium ion concentration as measured by the ICW lab	millimoles
Na	Sodium ion concentration as measured by the ICW lab	millimoles
NH4	Ammonium ion concentration as measured by the ICW lab	millimoles
K	Potassium ion concentration as measured by the ICW lab	millimoles
Mg_IC	Magnesium ion concentration as measured by the ICW lab	millimoles
Mg_USC	Magnesium ion concentration as measured by the USC lab	millimoles
Ca_IC	Calcium ion concentration as measured by the ICW lab	millimoles
Ca_USC	Calcium ion concentration as measured by the USC lab	millimoles
Cl	Chloride ion concentration as measured by the ICW lab	millimoles
Br	Bromide ion concentration as measured by the ICW lab	millimoles
F	Flouride ion concentration as measured by the ICW lab	millimoles
NO3	Nitrate concentration as measured by the ICW lab	millimoles
PO4	Phosphate concentration as measured by the ICW lab	millimoles
SO4	Sulphate concentration as measured by the ICW lab	millimoles
Sr	Strontium ion concentration as measured by the USC lab	micromoles
Cs	Cesium ion concentration as measured by the USC lab	micromoles
Rb	Rubidium ion concentration as measured by the USC lab	micromoles
Ba	Barium ion concentration as measured by the USC lab	micromoles
Mn	Manganese ion concentration as measured by the USC lab	micromoles
Fe	Iron ion concentration as measured by the USC lab	micromoles
Ni	Nickel ion concentration as measured by the USC lab	micromoles
Cu	Copper ion concentration as measured by the USC lab	micromoles
Si	Silicon ion concentration as measured by the USC lab	micromoles

## Instruments

<b>Dataset-specific Instrument Name</b>	temperature probe
<b>Generic Instrument Name</b>	Water Temperature Sensor
<b>Dataset-specific Description</b>	Hand-held temperature probe
<b>Generic Instrument Description</b>	General term for an instrument that measures the temperature of the water with which it is in contact (thermometer).

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### Milos\_vents\_2012

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/474391">https://www.bco-dmo.org/deployment/474391</a>
<b>Platform</b>	shoreside Milos_vents
<b>Start Date</b>	2012-05-21
<b>End Date</b>	2012-05-30
<b>Description</b>	Vent fluids and sediment cores were collected at shallow-water hydrothermal vent sites at Paleochori Bay, Milos Island, Greece, and analyzed for their chemical composition (anion, cation, trace elements/metals).

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### **Autotrophic carbon fixation at a shallow-water hydrothermal system: Constraining microbial activity, isotopic and geochemical regimes (Hydrothermal Autotrophic Carbon Fixation)**

**Coverage:** Shallow-water hydrothermal vents, Paleochori Bay, Milos Island, Greece

In this project we studied the shallow-water hydrothermal vent sites at Milos Island (Greece) to better understand the extent of autotrophic carbon fixation and its chemical and isotopic signature along environmental (redox/thermal) gradients. This was a 12-day long expedition (May 18 to 30, 2012) to sample vent fluids, gases and retrieve sediment cores at Paleochori Bay by using SCUBA diving at 8-10 m depth. In addition to the submarine vent sites, two subaerial locations of venting were identified at 36° 40' 28"N - 24° 31' 14" E and 36° 40' 25" N - 24° 30' 44" E. Both the subaerial and submarine sites are located on the same fracture zone that likely controls the hydrothermal circulation of evolved meteoritic water and seawater within the magmatic zone of Milos Island. To this end, the geochemistry of the fluids and gases emitted from subaerial sites provide important information towards identifying the linkage between the subaerial and submarine magmatic activity and provide insights on the metabolic functions (e.g. H<sub>2</sub> oxidation, Fe(III) reduction, C and S cycling) of the subsurface microbial community.

#### Abstract:

Currently, there is only limited information on the identity and activity of the microorganisms carrying out CO<sub>2</sub>-fixation in situ, despite the fact that these organisms form the basis of their respective ecosystems. Representatives that are able to grow autotrophically are known to exist in almost all major groups of prokaryotes, and these organisms play essential roles in ecosystems by providing a continuous supply of organic carbon for heterotrophs. Microorganisms present in extreme environments utilize CO<sub>2</sub>-fixation pathways other than the Calvin-Benson-Bassham (CBB) cycle. At present, five alternative autotrophic CO<sub>2</sub>

fixation pathways are known. Different carbon fixation pathways result in distinct isotopic signatures of the produced biomass due to the isotopic discrimination between light ( $^{12}\text{C}$ ) and heavy ( $^{13}\text{C}$ ) carbon by the carboxylating enzymes. Thus, inferences about the carbon fixation pathway predominantly utilized by the microbial community can also be made based on the stable carbon isotopic composition of the organic matter, in extant systems as well as in the geological record. However, at present little is known about the systematics and extents of fractionation during carbon fixation by prokaryotic organisms, and to our knowledge no studies exist that have systematically studied the relationship between the operation of different carbon fixation pathways and how this is reflected in the stable carbon isotopic composition in a natural system. This is a 2-year interdisciplinary, international research program that employs a powerful combination of cutting-edge research tools aiming to improve our understanding of autotrophic carbon fixation and its chemical and isotopic signature along environmental gradients in a natural hydrothermal system. The following hypotheses are addressed:

1. The diversity of microorganisms present along a thermal and redox gradient, and rates of  $\text{CO}_2$  fixation, will reflect adaptation to in situ temperatures and geochemical conditions
2. Microorganisms utilizing the CBB cycle for autotrophic  $\text{CO}_2$ -fixation will represent a smaller percentage of the chemolithoautotrophic community at higher temperatures, where microorganisms utilizing alternative  $\text{CO}_2$ -fixation pathways dominate
3. Isotopic values of biomass and specific biomarker molecules will vary along a thermal and redox gradient from zones characterized by a higher hydrothermal fluid flux and thus higher temperatures to the surrounding, cooler areas, corresponding to the physiology of the microorganisms utilizing different pathways for carbon fixation

The PIs will use a multidisciplinary approach to delineate the relative contribution of the different carbon fixation pathways along an environmental gradient by combining metagenomic analyses coupled with: 1) an assessment of the frequency and the expression of specific key genes involved in carbon fixation, and 2) with the measurement of carbon fixation rates. These data will be integrated with the determination of stable C isotopic composition of biomass, DIC, and specific hydrocarbons/lipids. Due to its easy accessibility, well-established environmental gradients, and extensive background information, the shallow-water vents off Milos (Greece) will be used as a natural laboratory to perform these studies.

Intellectual Merit. The data generated in this study will allow constraints on the relationship between autotrophic carbon fixation and the resulting isotopic signatures of biomass and specific biomarker molecules (e.g.  $\text{CH}_4$ ,  $\text{C}_2+$  alkanes, lipids) in a natural system. This has implications for assessing the importance of carbon fixation in extant ecosystems, and it will also provide a tool to improve the interpretation of isotopic values in the geological record.

[ [table of contents](#) | [back to top](#) ]

---

## Program Information

### Center for Dark Energy Biosphere Investigations (C-DEBI)

**Website:** <http://www.darkenergybiosphere.org>

**Coverage:** Global

The mission of the Center for Dark Energy Biosphere Investigations (C-DEBI) is to explore life beneath the seafloor and make transformative discoveries that advance science, benefit society, and inspire people of all ages and origins.

C-DEBI provides a framework for a large, multi-disciplinary group of scientists to pursue fundamental questions about life deep in the sub-surface environment of Earth. The fundamental science questions of C-DEBI involve exploration and discovery, uncovering the processes that constrain the sub-surface biosphere below the oceans, and implications to the Earth system. What type of life exists in this deep biosphere, how much, and how is it distributed and dispersed? What are the physical-chemical conditions that promote or limit life? What are the important oxidation-reduction processes and are they unique or important to humankind?

How does this biosphere influence global energy and material cycles, particularly the carbon cycle? Finally, can we discern how such life evolved in geological settings beneath the ocean floor, and how this might relate to ideas about the origin of life on our planet?

C-DEBI's scientific goals are pursued with a combination of approaches:

- (1) coordinate, integrate, support, and extend the research associated with four major programs—Juan de Fuca Ridge flank (JdF), South Pacific Gyre (SPG), North Pond (NP), and Dorado Outcrop (DO)—and other field sites;
- (2) make substantial investments of resources to support field, laboratory, analytical, and modeling studies of the deep seafloor ecosystems;
- (3) facilitate and encourage synthesis and thematic understanding of submarine microbiological processes, through funding of scientific and technical activities, coordination and hosting of meetings and workshops, and support of (mostly junior) researchers and graduate students; and
- (4) entrain, educate, inspire, and mentor an interdisciplinary community of researchers and educators, with an emphasis on undergraduate and graduate students and early-career scientists.

Note: Katrina Edwards was a former PI of C-DEBI; James Cowen is a former co-PI.

### **Data Management:**

C-DEBI is committed to ensuring all the data generated are publically available and deposited in a data repository for long-term storage as stated in their [Data Management Plan \(PDF\)](#) and in compliance with the [NSF Ocean Sciences Sample and Data Policy](#). The data types and products resulting from C-DEBI-supported research include a wide variety of geophysical, geological, geochemical, and biological information, in addition to education and outreach materials, technical documents, and samples. All data and information generated by C-DEBI-supported research projects are required to be made publically available either following publication of research results or within two (2) years of data generation.

To ensure preservation and dissemination of the diverse data-types generated, C-DEBI researchers are working with BCO-DMO Data Managers make data publicly available online. The partnership with BCO-DMO helps ensure that the C-DEBI data are discoverable and available for reuse. Some C-DEBI data is better served by specialized repositories (NCBI's GenBank for sequence data, for example) and, in those cases, BCO-DMO provides dataset documentation (metadata) that includes links to those external repositories.

[ [table of contents](#) | [back to top](#) ]

---

## **Funding**

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1123871</a>

[ [table of contents](#) | [back to top](#) ]