

Phytoplankton cell counts at 3 pCO₂ levels and 2 temperatures before and after 12 month conditioning off New Zealand South Island, near Taiaroa Head, 2011 (Plankton acclimation project)

Website: <https://www.bco-dmo.org/dataset/515271>

Version: 2014-05-15

Project

» [Experimental studies to understand and evaluate acclimation of marine plankton assemblages to increased CO₂ and temperature](#) (Plankton acclimation)

Contributors	Affiliation	Role
Hutchins, David A.	University of Southern California (USC-HIMS)	Principal Investigator
Caron, David	University of Southern California (USC-HIMS)	Co-Principal Investigator
Fu, Feixue	University of Southern California (USC-HIMS)	Co-Principal Investigator
Schnetzer, Astrid	North Carolina State University - Center for Marine Science and Technology (NCSU CMAST)	Co-Principal Investigator
Tatters, Avery	University of Southern California (USC-HIMS)	Student, Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

The study included a short-term two-week temperature/pCO₂ factorial matrix incubation experiment using a natural, mixed diatom assemblage, the isolation of clonal cultures from each treatment and conditioning of the clones to the pCO₂ and temperature combinations from which they were isolated for 1 year. Finally, the conditioned clones were recombined into artificial communities and allowed to compete, followed by a comparison of final community structure with that observed in the original two-week natural community experiment.

For this dataset, cell abundances of *Cylindrotheca fusiformis*, *Coscinodiscus* sp., *Thalassiosira* sp., *Pseudonitzschia delicatissima*, *Navicula* sp and *Chaetoceros criophilus* assessed after the initial natural community experiment and the 12-month conditioned artificial community incubations at low, medium and high CO₂ concentrations and temperatures of 14°C and 19°C. Values given include the three replicate sample counts, the mean of counts and standard deviations on the three replicate bottles.

[Experimental Design Chart](#)

Relevant References:

These data are published as Supplemental Table 2 in Avery O. Tatters, Michael Y. Roleda, Astrid Schnetzer,

Methods & Sampling

Experimental design

An overview of the experimental design is depicted in figure 1, and followed the general protocols for recently published dinoflagellate community experiments [21]. Sequentially, the study included a short-term two-week temperature/pCO₂ factorial matrix incubation experiment using a natural, mixed diatom assemblage, the isolation of clonal cultures from each treatment and conditioning of the clones to the pCO₂ and temperature combinations from which they were isolated for 1 year. Finally, the conditioned clones were recombined into artificial communities and allowed to compete, followed by a comparison of final community structure with that observed in the original two-week natural community experiment.

Short-term natural community incubation experiment

A mixed diatom assemblage that consisted primarily of *Cylindrotheca fusiformis* Reimann and Lewin 1964, *Coscinodiscus* spp., *Thalassiosira* spp., *Pseudo-nitzschia delicatissima* (Cleve) Heiden 1928, *Navicula* sp. and *Chaetoceros criophilus* (Castracene) sensu Hust 1886 was collected off the city of Dunedin on the South Island of New Zealand in January of 2011. The water was collected approximately 3 km offshore from Tairua Head at the mouth of Otago Harbour halfway to Munida (45, 45.098 S 170, 48.68 E). The ambient sea surface temperature was 14.88C.

Seawater was collected for both the initial incubations and the short-term experimental dilution water. All water was combined into an approximately 500 l container and subsampled after filtering through 80 mm mesh to remove large zooplankton. Volumes (800 ml) were added to triplicate polycarbonate bottles and spiked with an f/50 nutrient derivative (10 mM NaNO₃ 2, 0.8 mM NaH₂PO₄ 32, 10 mM Na₂SiO₃ and f/50 vitamin and trace metal concentrations [27,28]) to promote diatom growth. The bottles were incubated on a 12 L : 12 D cycle under 140 mE of cool white fluorescent illumination in free-standing laboratory incubators at 14 or 19.8C. The temperatures (ambient and 19.8C) were selected based on predicted sea surface warming from the IPCC [29]. Triplicate sterilized 1 l polycarbonate bottles were gently bubbled at each temperature using commercially prepared air/ CO₂ mixtures (Alphagaz, Air Liquide) at three concentrations also based on IPCC scenarios (approx. 210 matm ¹/₄ pre-industrial pCO₂; approx. 370 matm ¹/₄ current pCO₂; and approx. 560 matm ¹/₄ future, year 2050 projected pCO₂) [29]. Cellular abundances in an unbubbled control treatment did not significantly deviate from results of the current pCO₂-bubbled treatment (data not shown). This methodology has been used for other CO₂ experiments [21,30,31], including previous diatom studies [10,16,20].

The six pCO₂/temperature treatments were maintained in active growth using semicontinuous culture methods [21]. Each bottle was diluted to the original time-zero in vivo chlorophyll a fluorescence value every 2 days with nutrient-amended 0.2 mmfiltered seawater. Aliquots were removed initially and after one and two weeks for examination of carbonate system parameters and community structure using microscopic cell counts.

Establishment of clonal cultures

Two to four individual cells from the six dominant diatom species were isolated from each of the short-term incubation bottles at the termination of the experiment. Inverted light microscopy was used to make taxonomic determinations based on morphological characteristics to make sure the isolates for each cell line were from the same species [32]. These monospecific clones were propagated in 24-well plates prior to being transferred to tissue culture flasks for long-term maintenance under pCO₂ and temperature conditions identical to those from which they were isolated. A set of the culture isolates were transported under controlled temperature conditions to the University of Southern California in Los Angeles, CA, USA, where conditioning of the isolates and the 12-month community recombination experiments presented here were carried out. The culture isolates were maintained unreplicated for the first few weeks until they were verified to be established and growing well, at which time they were transferred into triplicate cultures for long-term maintenance; initial growth rates were obtained from these original unreplicated cell lines. These cultures were then maintained for a period of 1 year in exponential growth phase using the same recipe of autoclave-sterilized enriched seawater growth medium, and with other environmental variables such as light, pCO₂ bubbling, temperature etc., maintained as in the two-week natural community experiment. Semicontinuous weekly dilutions were performed based on specific growth rates within each bottle, calculated as in [21]. The approximate number of generations during this time period was: *C. fusiformis* (185-212), *Coscinodiscus* sp. (169-229), *Thalassiosira* sp. (179-200), *P. delicatissima* (178-221), *Navicula* sp. (188-212) and *C. criophilus* (194-236).

Artificial community competition experiments

After the 12-month pCO₂/temperature conditioning period, the cultures were recombined into artificial communities in the same relative proportions and abundance as in the original natural assemblage collected from Otago Harbour. The incubations of these artificial communities were performed under experimental conditions, duration and dilution frequencies identical to those of the original short-term natural community experiment.

Cell counts and growth rates

Samples for cell counts were obtained at the time of collection, before and after dilution and upon termination of the natural and artificial community incubations to determine abundances of each species. Cell-specific growth rates for each clonal culture were determined in individual culture flasks at the beginning of the 1 year conditioning period and in triplicate replicates after approximately 10 months of conditioning. These were calculated from samples taken 3 days apart using the growth rate equation $m = \frac{1}{t_1 - t_0} \ln(N_{t_1}/N_{t_0})$ (where N is the number of cells at time t₁ and t₀ (in days)) and represent a long-term steady-exponential state of growth. Algal cells were collected in 30 ml borosilicate glass scintillation vials, preserved with acidified Lugol's solution and enumerated using an Accu-Scope v. 3032 inverted microscope using the Utermöhl method [33].

Carbonate system characterization

Samples for carbonate system parameter analysis were taken at the time of the natural sample collection and at the termination of the short- and long-term experiments. Spectrophotometric pH for the initial community incubations was measured after [34] as described in [35] using a UV-vis spectrophotometer (Ocean Optics USB4000). For samples from the 12-month community incubations, spectrophotometric pH was determined using a Shimadzu 1800UV spectrophotometer according to a similar method [36]. Temperature was monitored using standard laboratory incubator thermometers and salinity by conductivity with an interchangeable probe using an Orion 5-star plus pH meter. For pH measurements, temperature and salinity values for the initial experiment were 23.68C and 35, respectively. For the conditioned experiment, the temperature was 25.8C and salinity 35. Dissolved inorganic carbon was analysed using a CM5230 CO₂ coulometer (UIC) [37]. Experimental pCO₂ was calculated using CO₂SYS software [22] with dissociation constants from Dickson & Millero [38] using the combined data of [39,40] and KSO₄ from [41] (table 1).

Data Processing Description

Statistics

Multivariate analyses were conducted using the PRIMER v6 statistics package [42] with the PERMANOVA add-on [43]. Bray-Curtis similarities were computed following square-root transformation of final cell abundances (cells m⁻²) for all six diatom species from replicate bottles. PERMANOVA was used to test for significant differences among and within predefined groups in response to differing pCO₂ competition levels and differing temperature. Data from the original natural community experiment and from the artificial community competition trials 12 months later were analysed. Pseudo-F values of 1 are typical of a large overlap among sample groups that are being compared (confirmation of H₀ hypothesis), whereas pseudo-F values greater than 1 indicate little or no overlap between the compared groups [43]. Observed interactions between pCO₂ and temperature were interrogated using PERMANOVA as well as pairwise comparisons (one-way ANOSIM [38]). R-values close to zero were indicative of no difference among groups, whereas R-values close to 1 meant that dissimilarities among groups were larger than any dissimilarity within groups [42].

We used a two-way crossed design for the ANOSIM routine to examine the comparative effects of differing pCO₂ competition levels and differing temperature on algal assemblages. This approach tests the average effect of pCO₂ levels during competition removing differences in temperature and the average effect of temperature levels removing differences in competition pCO₂ [42]. Cell abundance information for these analyses was taken from the final time points of our initial natural community experiment and the artificial community competition trial after 12 months.

Differences between specific growth rates after 10 months of conditioning in addition to cell abundances from the original natural community and the final artificial community experiments under the four temperature and pCO₂ combinations were tested using one-way ANOVA using Microsoft EXCEL 2013.

Literature Cited

21. Tatters AO, Schnetzer A, Fu F-X, Lie AAY, Caron DA, Hutchins DA. 2013 Short- versus long-term responses to changing CO₂ in a coastal dinoflagellate bloom: implications for interspecific competitive

- interactions and community structure. *Evolution* 67, 1879-1891. (doi:10.1111/evo.12029)
22. Müller MN, Schulz KG, Riebesell U. 2010 Effects of long-term high CO₂ exposure on two species of coccolithophores. *Biogeosciences* 7, 1109-1116. (doi:10.5194/bg-7-1109-2010)
23. Collins S, Bell G. 2004 Phenotypic consequences of 1000 generations of selection at elevated CO₂ in a green alga. *Nature* 431, 566-569. (doi:10.1038/nature02945)
24. Collins S. 2010 Competition limits adaptation and productivity in a photosynthetic alga at elevated CO₂. *Proc. R. Soc. B* 278, 247-255. (doi:10.1098/rspb.2010.1173)
25. Lohbeck KT, Riebesell U, Reusch TBH. 2012 Adaptive evolution of a key phytoplankton species to ocean acidification. *Nat. Geosci.* 5, 346-351. (doi:10.1038/NGEO1441)
26. Lohbeck KT, Riebesell U, Collins S, Reusch TBH. 2012 Functional genetic divergence in high CO₂ adapted *Emiliania huxleyi* populations. *Evolution* 67, 1892- 1900. (doi:10.1111/j.1558-5646.2012.01812.x)
27. Guillard RRL. 1975 Culture of phytoplankton for feeding marine invertebrates. In *Culture of marine invertebrate animals* (eds WL Smith, MH Chanley), pp. 26-60. New York, NY: Plenum Press.
28. Guillard RRL, Ryther JH. 1962 Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* Cleve. *Can. J. Microbiol.* 8, 229-239. (doi:10.1139/m62-029)
29. Solomon S et al. 2007 Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK: Cambridge University Press.
30. Fu F-X, Warner ME, Zhang Y, Feng Y, Hutchins DA. 2007 Effects of increased temperature and CO₂ on photosynthesis, growth and elemental ratios of marine *Synechococcus* and *Prochlorococcus* (Cyanobacteria). *J. Phycol.* 43, 485-496. (doi:10.1111/j.1529-8817.2007.00355.x)
31. Fu F-X, Mulholland MR, Garcia N, Beck A, Bernhardt PW, Warner ME, Sanudo-Wilhelmy SA, Hutchins DA. 2008 Interactions between changing pCO₂, N₂ fixation, and Fe limitation in the marine unicellular cyanobacterium *Crocospaera*. *Limnol. Oceanogr.* 53, 2472-2484. (doi:10.4319/lo.2008.53.6.2472)
32. Tomas C. (ed.) 1997 Identifying marine phytoplankton. San Diego, CA: Academic Press.
33. Utermöhl H. 1931 Neue Wege in der quantitativen Erfassung des Planktons. (Mit besonderer Berücksichtigung des Ultraplanktons). *Verh. Int. Verein. Limnol.* 5, 567-596.
34. Zhang H, Byrne RH. 1996 Spectrophotometric pH measurements of surface seawater at in-situ conditions: absorbance and protonation behaviour of thymol blue. *Mar. Chem.* 52, 17-25. (doi:10.1016/0304-4203(95)00076-3)
35. McGraw CM, Cornwall C, Reid MR, Currie K, Hepburn CD, Boyd PW, Hurd CL, Hunter KA. 2010 An automated pH-controlled culture system for laboratory-based ocean acidification experiments. *Limnol. Oceanogr. Methods* 8, 686-694. (doi:10.4319/lom.2010.8.686)
36. Clayton TD, Byrne RH. 1993 Spectrophotometric seawater pH measurements: total hydrogen ion concentration scale calibration of m-cresol purple and at-sea results. *Deep Sea Res. I* 40, 2115-2129. (doi:10.1016/0967-0637(93)90048-8)
37. King AL, Sanudo-Wilhelmy SA, Leblanc K, Hutchins DA, Fu F-X. 2011 CO₂ and vitamin B12 interactions determine bioactive trace metal requirements of a subarctic Pacific diatom. *ISME J.* 5, 1388-1396. (doi:10.1038/ismej.2010.211)
38. Dickson AG, Millero FJ. 1987 A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res.* 34, 1733-1743. (doi:10.1016/0198-0149(87)90021-5)
39. Mehrbach C, Culbertson CH, Hawley JE, Pytkowicz RM. 1973 Measurement of the apparent dissociation constants of carbonic acid in seawater at rstb.royalsocietypublishing.org *Phil Trans R Soc B* 368: 20120437 13 Downloaded from rstb.royalsocietypublishing.org on May 19, 2014 atmospheric pressure. *Limnol. Oceanogr.* 18, 897-907. (doi:10.4319/lo.1973.18.6.0897)
40. Hansson I. 1973 A new set of acidity constants for carbonic acid and boric acid in seawater. *Deep Sea Res.* 20, 461-478.
41. Dickson AG. 1990 Standard potential of the reaction: $\text{AgCl(s)} + \frac{1}{2}\text{H}_2\text{(g)} \rightleftharpoons \frac{1}{4}\text{Ag(s)} + \text{pHCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15 K. *J. Chem. Thermodyn.* 22, 113-127. (doi:10.1016/0021-9614(90)90074-Z)
42. Clarke KR, Warwick RM. 2001 Change in marine communities: an approach to statistical analysis and interpretation, 2nd edn. Ivybridge, UK: Primer-E.

[[table of contents](#) | [back to top](#)]

Data Files

File
phyto_abund.csv (Comma Separated Values (.csv), 10.39 KB) MD5:5d8b705f3325b722b52c430cfa23f5e7
Primary data file for dataset ID 515271

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
species	species name	unitless
temp_init	temperature before conditioning period	degrees Celsius
pCO2_level	relative level of pCO2	unitless
sample	sample name: A, B, C, average, or standard deviation of the 3 samples	unitless
incubation_init	initial period of incubation	months
pCO2_init	initial partial pressure of carbon dioxide	micro-atmospheres
abundance_init	abundance of cells before conditioning period	cells/milliliter
incubation_cond	duration of pCO2/temperaure conditioning period of incubation	months
temp_cond	temperature during conditioning period	degrees Celsius
pCO2_cond	partial pressure of carbon dioxide during conditioning period	micro-atmospheres
abundance_cond	abundance of cells after conditioning period	cells/milliliter

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	CO2 coulometer
Generic Instrument Name	CO2 Coulometer
Dataset-specific Description	Dissolved inorganic carbon was analysed using a CM5230 CO2 coulometer (UIC)
Generic Instrument Description	A CO2 coulometer semi-automatically controls the sample handling and extraction of CO2 from seawater samples. Samples are acidified and the CO2 gas is bubbled into a titration cell where CO2 is converted to hydroxyethylcarbonic acid which is then automatically titrated with a coulometrically-generated base to a colorimetric endpoint.

Dataset-specific Instrument Name	Conductivity Meter
Generic Instrument Name	Conductivity Meter
Dataset-specific Description	Orion 5-star plus pH meter with an interchangeable probe for conductivity measurements to calculate salinity in conjunction with a standard laboratory incubator thermometer.
Generic Instrument Description	Conductivity Meter - An electrical conductivity meter (EC meter) measures the electrical conductivity in a solution. Commonly used in hydroponics, aquaculture and freshwater systems to monitor the amount of nutrients, salts or impurities in the water.

Dataset-specific Instrument Name	spectrophotometer
Generic Instrument Name	Spectrophotometer
Dataset-specific Description	UV-vis spectrophotometer: Ocean Optics USB4000
Generic Instrument Description	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

Dataset-specific Instrument Name	spectrophotometer
Generic Instrument Name	Spectrophotometer
Dataset-specific Description	Shimadzu 1800UV
Generic Instrument Description	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

[[table of contents](#) | [back to top](#)]

Deployments

Hutchins_2011

Website	https://www.bco-dmo.org/deployment/515350
Platform	USC
Start Date	2011-01-01
End Date	2011-01-31
Description	Collections were made off the city of Dunedin on the South Island of New Zealand in January of 2011. The water was collected approximately 3 km offshore from Taiaora Head at the mouth of Otago Harbour halfway to Munida. http://marineregions.org/gazetteer.php?p=details&id=24560

Project Information

Experimental studies to understand and evaluate acclimation of marine plankton assemblages to increased CO₂ and temperature (Plankton acclimation)

Coverage: Bloom sample retrieved from near-shore water near Venice Beach, CA

Progressing ocean acidification and increasing sea surface temperature may significantly impact marine plankton community structure and community-level processes. Yet, our ability to predict specific responses is still limited because of the tremendous taxonomic complexity of microbial assemblages and the limitations of the methodological and experimental tools presently available to test specific hypotheses. Research to study community level effects due to a changing CO₂/temperature regime often involve short-term field incubations that subject organisms to simulated 'greenhouse' conditions. A central question for understanding global climate change is whether the trends and patterns that are observed in communities during short-term manipulations can be extrapolated to the responses of fully acclimated plankton communities over decadal or longer timescales.

The specific objectives of this research program are: 1) to examine how protistan communities restructure in response to increased seawater CO₂ concentrations and temperature in semi-continuous field incubation experiments, and 2) to evaluate if the dominant algal species that are isolated from either ambient or increased CO₂ and temperature treatments in field experiments will re-establish dominance under the same conditions in acclimated laboratory culture competition studies. Changes in community structure of natural protistan assemblages in our experimental treatments will be followed using image-based methods (flow cytometry, FlowCAM and microscopy) in combination with state-of the art molecular tools (DNA fingerprinting). Molecular approaches have begun to reveal an incredible high diversity for marine microbes and stimulate debate in regard to the ubiquitous presence of a microbial 'Rare Biosphere' that is, the presence of a huge number of species that are present at extremely small percentages of the total abundance of microbes, among a much smaller percentage of dominant ones. Little is known about the ecological significance of these rare species, and the investigators hypothesize that change in CO₂ and temperature will select for some of these members that are inconspicuous under ambient conditions.

The unique aspect of this experimental approach is the combined use of field incubations that encompass entire natural microbial assemblages, with a series of laboratory culture competition trials that focus on the same groups of algae after extended acclimation, to evaluate the validity of short-term experiments that examine changing CO₂ and temperature. First, field incubation experiments will be conducted to characterize changes in protistan community structure under ambient and future CO₂/temperature regimes. Second, clonal algal strains will be isolated from dominant taxa in present day and greenhouse treatments, and cultivated for extended periods under their 'preferred' CO₂/temperature conditions. Finally, mixtures of these acclimated strains will be competed against each other, to re-examine their responses to ambient and greenhouse conditions and compare them to the responses observed in the unacclimated field incubation experiments.

Two graduate students will make this project the focus of their Ph.D. research at USC, and undergraduate students will be involved in the field and laboratory work. Results from this research will be incorporated in lesson plans on microbial diversity and global climate change. Dissemination of data and results is planned on a project website. The PIs in this project also participate in an on-going, innovative, NSF-funded program (Centers for Ocean Science Education Excellence; COSEE-West) which focuses on personal involvement of faculty in a custom framework to allow an effective connection with K-12 teachers, thus improving math and science education in disadvantaged parts of Southern California.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0962309

[[table of contents](#) | [back to top](#)]