

Carbon and nitrogen content of *E. huxleyi* at 3 pCO₂ levels, 2011-2012 (E Hux Response to pCO₂ project)

Website: <https://www.bco-dmo.org/dataset/521403>

Data Type: experimental

Version: 2

Version Date: 2016-12-13

Project

» [Planktonic interactions in a changing ocean: Biological responses of *Emiliana huxleyi* to elevated pCO₂ and their effects on microzooplankton](#) (E Hux Response to pCO₂)

Contributors	Affiliation	Role
Olson, M Brady	Western Washington University - Shannon Point Marine Center (SPMC)	Principal Investigator, Contact
Love, Brooke	Western Washington University (WWU)	Co-Principal Investigator
Strom, Suzanne	Western Washington University - Shannon Point Marine Center (SPMC)	Co-Principal Investigator
Wuori, Tristen	Western Washington University - Shannon Point Marine Center (SPMC)	Student
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

These data are not yet available because they are part of a graduate thesis. Please contact the PI for further information.

These data show cellular characterizations of two strains of *Emiliana huxleyi* cultured semi-continuously over a period 13-14 days under three different pCO₂ concentrations (400 ppmv, 750 ppmv, and 1000 ppmv). Cellular characterization measurements were taken throughout the course of the experiments, resulting in a time course data set. CO₂ chemistry was also monitored over the course of the experiment. Cellular characterizations included here: cellular particulate organic carbon and nitrogen, cellular particulate inorganic carbon.

Emiliana huxleyi strains:

Strain NCMA 2668, calcifying phenotype, isolated from Gulf of Maine 2002

Strain NCMA 374, non-calcifying phenotype, isolated from Gulf of Maine 1990

Related Datasets:

[Emiliana huxleyi Chl-a, POC, cell volumes](#)

[Emiliana huxleyi dilution calculations](#)

[Emiliana huxleyi DMSP](#)

[Emiliana huxleyi growth rates](#)

Relevant References:

Methods & Sampling

Culturing conditions:

Cultures of *E. huxleyi* Strain NCMA 2668 and 374 were inoculated at low cell density into media prepared from autoclaved filtered seawater with f/50 nutrient amendment. Cell populations were allowed to acclimate for approximately five generations, until cell density neared levels likely to significantly change the pH/pCO₂. Daily dilutions of cultures with pre-equilibrated media kept cell density low (<1x10⁵ cells/ml), ensured cells remained in exponential growth phase and prevented excessive drawdown of nutrients and CO₂. Cell density was determined by flow cytometry (model described below) and each flask was diluted with media that was continuously sparged with air containing 400, 750 or 1000 ppm CO₂. Air mixtures were created using CO₂ free air (Powerex air compressor, and Twin Towers CO₂ scrubber) and pure CO₂ (Airgas) combined using a system of mass flow controllers (Sierra Instruments) and verified using a non-dispersive infrared CO₂ sensor (Licor 820). Cultures were maintained in 1-liter polycarbonate flasks at 15 degC under a 12/12 light dark cycle. Replicates (n=5) were placed in Plexiglas chambers which were supplied with a flow of the appropriate air mixture for each treatment. Preliminary experiments showed that gas exchange across the air/water surface significantly helped to maintain the target pCO₂ in cultures without the mechanical disturbance of bubbling. Sedimentation was minimized by gentle mixing of the cultures by rotation of the bottles twice a day, during sampling and dilution. Cell densities ranged between about 30,000 cells/ml after dilutions to 80,000 cells/ml on the following day. The culture volume that was removed was used for analyses, and replaced with pre-equilibrated media. Cultures were maintained in this fashion for 14 days. This experiment was carried out twice, in 2011 and 2012.

Cellular carbon and nitrogen:

Samples for cellular particulate carbon and nitrogen were analyzed using a CE Elantech Flash EA 1112 elemental analyzer. In all, analysis blanks were run, and internal standards were inserted between samples, and remained within 1% of standard curve. For the calcifying strain (2668), samples were acid fumed for 24 h to drive off PIC. Values of organic carbon were subtracted from total carbon to yield cellular particulate inorganic carbon.

Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- changed ND to nd, no data

[[table of contents](#) | [back to top](#)]

Data Files

File
CN_v2.csv (Comma Separated Values (.csv), 5.20 KB) MD5:abab0ad9a9ea5ce66baef56432111c7a
Primary data file for dataset ID 521403

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Ehuxleyi_strain	E. huxleyi strain number	unitless
culture_day	Days of semi-continuous culture	days
treatment	CO2 treatment	unitless
sample_rep	Sample date and replicate	unitless
total_C_cell_pg	Total carbon per cell in picograms	picogram/cell
POC_cell_pg	Particulate organic carbon per cell in picograms	picogram/cell
PIC_cell_pg	Particulate inorganic carbon per cell in picograms	picogram/cell
N_cell_pg	Nitrogen per cell in picograms	picogram/cell

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Automatic titrator
Generic Instrument Name	Automatic titrator
Dataset-specific Description	Metrohm 888 Titrande with a Metrohm Ecotrode combined electrode, calibrated with TRIS and AMP buffers on the total H ⁺ ion pH scale.
Generic Instrument Description	Instruments that incrementally add quantified aliquots of a reagent to a sample until the end-point of a chemical reaction is reached.

Dataset-specific Instrument Name	CHN_EA
Generic Instrument Name	CHN Elemental Analyzer
Dataset-specific Description	CE Elantech Flash EA 1112 elemental analyzer
Generic Instrument Description	A CHN Elemental Analyzer is used for the determination of carbon, hydrogen, and nitrogen content in organic and other types of materials, including solids, liquids, volatile, and viscous samples.

Dataset-specific Instrument Name	CO2 Analyzer
Generic Instrument Name	CO2 Analyzer
Dataset-specific Description	Licor 820: a non-dispersive infrared CO2 sensor
Generic Instrument Description	Measures atmospheric carbon dioxide (CO2) concentration.

Dataset-specific Instrument Name	Flow Cytometer
Generic Instrument Name	Flow Cytometer
Dataset-specific Description	BD FACSCalibur flow cytometer
Generic Instrument Description	Flow cytometers (FC or FCM) are automated instruments that quantitate properties of single cells, one cell at a time. They can measure cell size, cell granularity, the amounts of cell components such as total DNA, newly synthesized DNA, gene expression as the amount messenger RNA for a particular gene, amounts of specific surface receptors, amounts of intracellular proteins, or transient signalling events in living cells. (from: http://www.bio.umass.edu/micro/immunology/facs542/facswhat.htm)

Dataset-specific Instrument Name	Fluorometer
Generic Instrument Name	Fluorometer
Dataset-specific Description	Turner Designs 10-AU fluorometer
Generic Instrument Description	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.

Dataset-specific Instrument Name	Gas Chromatograph
Generic Instrument Name	Gas Chromatograph
Dataset-specific Description	Shimadzu GC-14A gas chromatograph
Generic Instrument Description	Instrument separating gases, volatile substances, or substances dissolved in a volatile solvent by transporting an inert gas through a column packed with a sorbent to a detector for assay. (from SeaDataNet, BODC)

Dataset-specific Instrument Name	MFC
Generic Instrument Name	Mass Flow Controller
Dataset-specific Description	Sierra Instruments
Generic Instrument Description	Mass Flow Controller (MFC) - A device used to measure and control the flow of fluids and gases

Dataset-specific Instrument Name	compound microscope
Generic Instrument Name	Microscope - Optical
Dataset-specific Description	Olympus CH30 compound microscope networked to a Photometric CoolSNAP camera
Generic Instrument Description	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

Dataset-specific Instrument Name	spectrophotometer
Generic Instrument Name	Spectrophotometer
Dataset-specific Description	Agilent 5480 UV-VIS spectrophotometer (+/- 0.02)
Generic Instrument Description	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

[[table of contents](#) | [back to top](#)]

Deployments

Lab Olson B

Website	https://www.bco-dmo.org/deployment/521277
Platform	WWU
Start Date	2011-03-31
End Date	2016-09-15
Description	laboratory experiments

[[table of contents](#) | [back to top](#)]

Project Information

Planktonic interactions in a changing ocean: Biological responses of *Emiliana huxleyi* to elevated pCO₂ and their effects on microzooplankton (E Hux Response to pCO₂)

Description from NSF award abstract:

The calcifying Haptophyte *Emiliana huxleyi* appears to be acutely sensitive to the rising concentration of ocean pCO₂. Documented responses by *E. huxleyi* to elevated pCO₂ include modifications to their calcification rate and cell size, malformation of coccoliths, elevated growth rates, increased organic carbon production, lowering of PIC:POC ratios, and elevated production of the active climate gas DMS. Changes in these parameters are mechanisms known to elicit alterations in grazing behavior by microzooplankton, the oceans dominant grazer functional group. The investigators hypothesize that modifications to the physiology and biochemistry of

calcifying and non-calcifying Haptophyte *Emiliana huxleyi* in response to elevated pCO₂ will precipitate alterations in microzooplankton grazing dynamics. To test this hypothesis, they will conduct controlled laboratory experiments where several strains of *E. huxleyi* are grown at several CO₂ concentrations. After careful characterization of the biochemical and physiological responses of the *E. huxleyi* strains to elevated pCO₂, they will provide these strains as food to several ecologically-important microzooplankton and document grazing dynamics. *E. huxleyi* is an ideal organism for the study of phytoplankton and microzooplankton responses to rising anthropogenic CO₂, the effects of which in the marine environment are called ocean acidification; *E. huxleyi* is biogeochemically important, is well studied, numerous strains are in culture that exhibit variation in the parameters described above, and they are readily fed upon by ecologically important microzooplankton.

The implications of changes in microzooplankton grazing for carbon cycling, specifically CaCO₃ export, DMS production, nutrient regeneration in surface waters, and carbon transfer between trophic levels are profound, as this grazing, to a large degree, regulates all these processes. *E. huxleyi* is a model prey organism because it is one of the most biogeochemically influential global phytoplankton. It forms massive seasonal blooms, contributes significantly to marine inorganic and organic carbon cycles, is a large producer of the climatically active gas DMS, and is a source of organic matter for trophic levels both above and below itself. The planned controlled study will increase our knowledge of the mechanisms that drive patterns of change between trophic levels, thus providing a wider array of tools necessary to understand the complex nature of ocean acidification field studies, where competing variables can confound precise interpretation.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0961229

[[table of contents](#) | [back to top](#)]