Seawater carbonate chemistry for brooded coral larval experiments, Taiwan, March 2011 and 2012 (Cumbo, 2013)(MCR LTER project, Climate Coral Larvae project)

Website: https://www.bco-dmo.org/dataset/528806

Version: 2014-08-30

Project

- » Moorea Coral Reef Long-Term Ecological Research site (MCR LTER)
- » The ecophysiological basis of the response of coral larvae and early life history stages to global climate change (Climate_Coral_Larvae)

Program

» Long Term Ecological Research network (LTER)

Contributors	Affiliation	Role
Edmunds, Peter J.	California State University Northridge (CSUN)	Principal Investigator
Cumbo, Vivian R	California State University Northridge (CSUN)	Scientist
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Parameters
- Instruments
- Deployments
- Project Information
- Program Information
- Funding

Dataset Description

To evaluate the effects of temperature and pCO2 on coral larvae, brooded larvae of Pocillopora damicornis from Nanwan Bay, Taiwan (21°56.179' N, 120°44.85' E), were exposed to ambient (419-470 μatm) and high (604-742 μatm) pCO2 at ~25 and ~29 °C in two experiments conducted in March 2010 and March 2012. Larvae were sampled from four consecutive lunar days (LD) synchronized with spawning following the new moon, incubated in treatments for 24 h, and measured for respiration, maximum photochemical efficiency of PSII (F v/F m), and mortality.

The most striking outcome was a strong effect of time (i.e., LD) on larvae performance: respiration was affected by an LD \times temperature interaction in 2010; and 2012, as well as an LD \times pCO2 \times temperature interaction in 2012; F v/F m was affected by LD in 2010 (but not 2012); and mortality was affected by an LD \times pCO2 interaction in 2010, and an LD \times temperature interaction in 2012. There were no main effects of pCO2 in 2010, but in 2012, high pCO2 depressed metabolic rate and reduced mortality. Therefore, differences in larval performance depended on day of release and resulted in varying susceptibility to future predicted environmental conditions. These results underscore the importance of considering larval brood variation across days when designing experiments. Subtle differences in experimental outcomes between years suggest that transgenerational plasticity in combination with unique histories of exposure to physical conditions can modulate the response of brooded coral larvae to climate change and ocean acidification.

These data include the seawater carbonate chemistry monitored from the experimental tanks, March 2011 and 2012.

Related datasets:

brooded coral larvae 2 - larval release March 2003-2008 brooded coral larvae 2 - respiration_photosyth_mortality

These data are published in Vivian R Cumbo, Peter J Edmunds, Christopher B Wall, Tung-Yung Fan. (2013) Brooded coral larvae differ in their response to high temperature and elevated pCO2 depending on the day of release. Marine Biology. See Table 1.

Download complete data for this publication (Excel file)

Data also available from PANGAEA: DOI 10.1007/s00227-013-2280-y

Methods & Sampling

To determine the dissolved inorganic carbon (DIC) chemistry of seawater in the treatments, total alkalinity (TA), pH on the total scale, temperature, and salinity were measured using standard procedures, and used to calculate pCO2 (uatm), HCO3 - (umolkg-1), CO3 2- (umolkg-1), and the aragonite saturation state (XArag) of seawater using CO2SYS (Pierrot et al. 2006). TA (umol kg-1) was measured through potentiometric titration (SOP 3b, Dickson et al. 2007) using an automatic titrator (DL50, Mettler Toledo) filled with certified acid titrant (0.1 M HCl, 0.6 NaCl, Dickson Laboratory, Scripps Institution of Oceanography). The pH probe (DG101-SC, Mettler Toledo) attached to the titrator was 3-point calibrated with pH 4.00, 7.00, and 10.00 buffers (Fisher, NBS). Certified reference material with a known TA (Batch 98, http://andrew.ucsd. edu/index.html) was titrated daily to determine the accuracy and precision of the analyses.

Seawater samples (50 mL) from the treatment tanks were brought to 25C, weighed and titrated in a waterjacketed beaker within 2-3 h of collection. The pH values and the titrant volumes (cm3) obtained from the titrations were sub-sampled for the range between pH 3.0 and 3.5 and inserted into a Microsoft Excel spreadsheet (Fangue et al. 2010), which calculated Gran's function as a product of the mass of titrant added (Dickson et al. 2007). Treatment tank pH was determined spectrophotometrically using m-cresol purple dye (Sigma-Aldrich) following SOP 6b of Dickson et al. (2007) with modification (Fangue et al. 2010). Preliminary sampling of the seawater in the tanks throughout the day confirmed that the pCO2 treatments were stable over a 24-h period.

References:

Dickson AG, Sabine CL, Christian JR (eds) (2007) Guide to best practices for ocean CO2 measurements. PICES Special Publication 3, p 191

Fangue NA, O'Donnell MJ, Sewell MA, Matson PG, MacPherson AC, Hoffman GE (2010) A laboratory-based, experimental system for the study of ocean acidification effects on marine invertebrate larvae. Limnol Oceanogr Methods 8:441-452

Data Processing Description

BCO-DMO processing notes:

- added conventional header with dataset name, PI name, version date, reference information
- added lab, lat, lon, expt columns
- renamed parameters to BCO-DMO standard
- combined data from Table 1, 2012 and 2012 sorted by date, temp, treatment

[table of contents | back to top]

Data Files

File

brood2_carb_chem.csv(Comma Separated Values (.csv), 5.99 KB) MD5:3db9b129ca78efc0405b3f84c1be28

Primary data file for dataset ID 528806

[table of contents | back to top]

Parameters

Parameter	Description	Units
expt	experiment id	unitless
lab	laboratory	unitless
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
month	experiment month	unitless
tank	tank id number	tank
date	local date of measurement	yyyy-mm-dd
pН	pH: The measure of the acidity or basicity of an aqueous solution	pH scale
temp	temperature	degees Celsius
sal	salinity	PSU scale
TA	total alkalinity	umol/kg
pCO2	partial pressure of carbon dioxide by computation from pH and alkalinity	uatm
bicarbonate	concentration of bicarbonate ion ([HCO3]-) in seawater	umol/kg
carbonate	concentration of carbonate ion ([CO3]2-)	umol/kg
omega_Ara	the saturation state of seawater with respect to aragonite	unitless

[table of contents | back to top]

Instruments

Dataset-specific Instrument Name	Automatic titrator
Generic Instrument Name	Automatic titrator
Dataset-specific Description	Model DL50, Mettler-Toledo, Columbus, OH, USA
Generic Instrument Description	Instruments that incrementally add quantified aliquots of a reagent to a sample until the end-point of a chemical reaction is reached.

[table of contents | back to top]

Deployments

lab_Edmunds_NMMBA

Website	https://www.bco-dmo.org/deployment/58892
Platform	Natl Museum Mar. Bio. and Aquar. Taiwan
Start Date	2010-03-18
End Date	2010-03-24
Description	Experiments related to the research project: 'RUI- The ecophysiological basis of the response of coral larvae and early life history stages to global climate change' were conducted at the laboratories of the National Museum of Marine Biology and Aquarium in Southern Taiwan.

[table of contents | back to top]

Project Information

Moorea Coral Reef Long-Term Ecological Research site (MCR LTER)

Website: http://mcr.lternet.edu/

Coverage: Island of Moorea, French Polynesia

From http://www.lternet.edu/sites/mcr/ and http://mcr.lternet.edu/:

The Moorea Coral Reef LTER site encompasses the coral reef complex that surrounds the island of Moorea, French Polynesia (17°30'S, 149°50'W). Moorea is a small, triangular volcanic island 20 km west of Tahiti in the Society Islands of French Polynesia. An offshore barrier reef forms a system of shallow (mean depth ~ 5-7 m), narrow (-0.8-1.5 km wide) lagoons around the 60 km perimeter of Moorea. All major coral reef types (e.g., fringing reef, lagoon patch reefs, back reef, barrier reef and fore reef) are present and accessible by small boat.

The MCR LTER was established in 2004 by the US National Science Foundation (NSF) and is a partnership between the University of California Santa Barbara and California State University, Northridge. MCR researchers include marine scientists from the UC Santa Barbara, CSU Northridge, UC Davis, UC Santa Cruz, UC San Diego, CSU San Marcos, Duke University and the University of Hawaii. Field operations are conducted from the UC Berkeley Richard B. Gump South Pacific Research Station on the island of Moorea, French

MCR LTER Data: The Moorea Coral Reef (MCR) LTER data are managed by and available directly from the MCR project data site URL shown above. The datasets listed below were collected at or near the MCR LTER sampling locations, and funded by NSF OCE as ancillary projects related to the MCR LTER core research themes.

This project is supported by continuing grants with slight name variations: LTER: Long-Term Dynamics of a Coral Reef Ecosystem LTER: MCR II - Long-Term Dynamics of a Coral Reef Ecosystem

LTER: MCR IIB: Long-Term Dynamics of a Coral Reef Ecosystem LTER: MCR III: Long-Term Dynamics of a Coral Reef Ecosystem

The ecophysiological basis of the response of coral larvae and early life history stages to global climate change (Climate Coral Larvae)

Coverage: Moorea, French Polynesia: Southern Taiwan: California State University Northridge

Tropical coral reefs face a suite of environmental assaults ranging from anchor damage to the effects of global climate change (GCC). The consequences are evident throughout the tropics, where many coral reefs have lost a substantial fraction of their coral cover in a few decades. Notwithstanding the importance of reducing the impacts of environmental stresses, the only means by which these ecosystems can recover (or simply persist) is through the recruitment of scleractinians, which is a function of successful larval development, delivery, settlement, metamorphosis, and post-settlement events. Despite wide recognition of the importance of these processes, there are few pertinent empirical data, and virtually none that address the mechanisms mediating the success of early coral life stages in a physical environmental varying at multiple spatio-temporal

The objective of this research is to complete one of the first comprehensive ecophysiological analyses of the early life stages of corals through a description of: (1) their functionality under 'normal' conditions, and (2) their response to the main drivers of GCC. These analyses will be completed for 2 species representative of a brooding life history strategy, and the experiments will be completed in two locations, one (Taiwan) that provides unrivalled experience in coral reproductive biology, and superb microcosm facilities and the other (Moorea), with access to a relatively pristine environment, a well described ecological and oceanographic context (through the MCR-LTER), and the capacity to bring a strong biogeographic contrast to the project. The results of the study will be integrated through modeling to explore the effects of GCC on coral community structure over the

The following publications and data resulted from this project:

2013 Wall CB, Fan TY, Edmunds PJ. Ocean acidification has no effect on thermal bleaching in the coral Seriatopora caliendrum. Coral Reefs 33: 119-130.

Symbiodinium_Seriatopora photosynthesis

Symbiodinium Seriatopora PI curve

Symbiodinium_Seriatopora temp-salinity-light

Symbiodinium_Seriatopora water chemistry

- Download complete data for this publication (Excel file)

2013 Wall CB. Edmunds Pl. In situ effects of low pH and elevated HCO3- on juvenile Parites spp. in Moorea, French Polynesia. Biological Bulletin 225:92-101.

Data at MCR and PANGEA: doi.pangaea.de/10.1594/PANGAEA.833913

- Download complete data for this publication (Excel file)

2013 Vivian R Cumbo, Peter J Edmunds, Christopher B Wall, Tung-Yung Fan. Brooded coral larvae differ in their response to high temperature and elevated pCO2 depending on the day of release. Marine Biology DOI 10.1007/s00227-013-2280-y.

Data also at PANGEA: doi.pangaea.de/10.1594/PANGAEA.831612

brooded coral larvae 2 - carbonate chemistry

brooded coral larvae 2 - larval release March 2003-2008

brooded coral larvae 2 - respiration photosyth mortality

- Download complete data for this publication (Excel file)

2013 Edmunds PJ, Cumbo VR, Fan TY. Metabolic costs of larval settlement and metamorphosis in the coral Seriatopora caliendrum under ambient and elevated pCO2. Journal Experimental Marine Biology and Ecology 443: 33-38 Data also at PANGEA: doi:10.1594/PANGAEA.821644

Coral post-settlement physiology

- Download complete data for this publication (Excel file)

Aaron M Dufault, Aaron Ninokawa, Lorenzo Bramanti, Vivian R Cumbo, Tung-Yung Fan, Peter J Edmunds. The role of light in mediating the effects of ocean acidification on coral calcification. Journal of Experimental Biology 216: 1570-1577. coral-light expt.- PAR

coral-light expt.- carbonate chemistry

coral-light expt.- temp_salinity

coral-light expt.- growth coral-light expt.- protein

coral-light expt.- surviva

Download complete data for this publication (Excel file)

2012 Cumbo, VR, Fan TY, Edmunds PI, Effects of exposure duration on the response of *Pocillopora damicornis* larvae to elevated temperature and high pCO2. I Exp Mar Biol

Ecol 439: 100-107.

Data is also at PANGEA: doi:10.1594/PANGAEA.823582

brooded coral larvae 3 - carbonate chemistry brooded coral larvae 3 - light

brooded coral larvae 3 - mortality

brooded coral larvae 3 - protein

brooded coral larvae 3 - respiration and protein

brooded coral larvae 3 - respiration raw data

brooded coral larvae 3 - symbiont density

brooded coral larvae 3 - tank temperature

- Download part 1 of data for this publication (Excel file)

- Download tank parameters data for this publication (Excel file)

2012 Cumbo, VR, Fan TY, Edmunds PJ. Physiological development of brooded larvae from two pocilloporid corals in Taiwan. Marine Biology 159: 2853-2866.

brooded coral - carbonate chemistry

brooded coral - release brooded coral - respiration

brooded coral - settlement competency

brooded coral - size July

brooded coral - size protein symbionts photosynth

- Download complete data for this publication (Excel file)

Dufault, Aaron M; Vivian R Cumbo; Tung-Yung Fan; Peter J Edmunds. Effects of diurnally oscillating pCO2 on the calcification and survival of coral recruits. Royal Society of London (B) 279: 2951-2958. doi:10.1098/rspb.2011.2545

Data is also at PANGEA: doi:10.1594/PANGAEA.830185

recruit_growth_area

recruit_growth_weight

recruit seawater chemistry

recruit survival

- Download complete data for this publication (Excel file)

2011 Edmunds PJ, Cumbo V, Fan TY. Effects of temperature on the respiration of brooded larvae from tropical reef corals. Journal of Experimental Biology 214: 2783-2790.

CoralLarvae comparison respir

CoralLarvae_release CoralLarvae_respir

CoralLarvae size

- Download complete data for this publication (Excel file)

[table of contents | back to top]

Program Information


Long Term Ecological Research network (LTER)

Website: http://www.lternet.edu/

Coverage: United States

adapted from http://www.lternet.edu/

The National Science Foundation established the LTER program in 1980 to support research on long-term ecological phenomena in the United States. The Long Term Ecological Research (LTER) Network is a collaborative effort involving more than 1800 scientists and students investigating ecological processes over long temporal and broad spatial scales. The LTER Network promotes synthesis and comparative research across sites and ecosystems and among other related national and international research programs. The LTER research sites represent diverse ecosystems with emphasis on different research themes, and cross-site communication, network publications, and research-planning activities are coordinated through the LTER Network Office.



Site Codes

Site Codes		
AND	Andrews Forest LTER	
ARC	Arctic LTER	
BES	Baltimore Ecosystem Stı	
BLE	Beaufort Lagoon	
	Ecosystems LTER	
BNZ	Bonanza Creek LTER	
CCE	California Current	
	Ecosystem LTER	
CDR	Cedar Creek Ecosystem	
	Science Reserve	
CAP	Central Arizona-	
	Phoenix LTER	
CWT	Coweeta LTER	
FCE	Florida Coastal	
	Everglades LTER	
GCE	Georgia Coastal	
	Ecosystems LTER	
HFR	Harvard Forest LTER	
HBR	Hubbard Brook LTER	
JRN	Jornada Basin LTER	
KBS	Kellogg Biological	
	Station LTER	
KNZ	Konza Prairie LTER	
LUQ	Luquillo LTER	
MCM	McMurdo Dry Valleys LT	
MCR	Moorea Coral Reef LTEF	
NWT	Niwot Ridge LTER	
NTL	North Temperate Lakes I	
NES	Northeast U.S. Shelf LTE	
NGA	Northern Gulf of Alaska I	
PAL	Palmer Antarctica LTER	
PIE	Plum Island	
000	Ecosystems LTER	
SBC	Santa Barbara Coastal L	
SEV	Sevilleta LTER	

VCR Virginia Coast Reserve L

2017 LTER research site map obtained from https://lternet.edu/site/lter-network/

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0844785

[table of contents | back to top]