Seawater carbonate chemistry from experiment on brooded coral larval, March 2011, Taiwan (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects)

Website: https://www.bco-dmo.org/dataset/535163

Data Type: experimental Version: 1 Version Date: 2014-10-07

Moorea Coral Reef Long-Term Ecological Research site (MCR LTER)

» The ecophysiological basis of the response of coral larvae and early life history stages to global climate change (Climate Coral Larvae)

Program

» Long Term Ecological Research network (LTER)

Contributors	Affiliation	Role
Edmunds, Peter J.	California State University Northridge (CSUN)	Principal Investigator
Cumbo, Vivian R	California State University Northridge (CSUN)	Co-Principal Investigator
Fan, Tung-Yung	National Museum of Marine Biology and Aquarium (NMMBA)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

The physiological development of brooded larvae from the pocilloporid corals Pocillopora damicornis in southern Taiwan under elevated temperature and pCO2 was examined. These data include seawater carbonate chemistry from experiments on the physiological development of brooded coral larvae conducted in March 2011. Reported parameters include temperature, salinity, pH, total alkalinity, pCO2, bicarbonate and carbonate concentration, and the saturation state (omega) for aragonite. These data were published in Cumbo et al, JEMBE, 2013.

Table of Contents

- Coverage
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Supplemental Files
- **Related Publications**
- Related Datasets
- **Parameters**
- Instruments
- Deployments
- Project Information
- Program Information
- **Funding**

Coverage

Spatial Extent: Lat:21.93817 Lon:120.74602 Temporal Extent: 2011-03-02 - 2011-03-18

Methods & Sampling

Treatments were created in eight 150 L tanks, each filled with 120 L of filtered (1 um) seawater that was changed partially (20%) every day (at ~17:00 h). Tanks were individually heated (300 Wheaters, Taikong Corporation) and chilled (Aquatech Ac11 or Shyeh Duwai Enterprise), with the temperature regulated using programmable, digital controllers (±0.1 °C, AquaControllers, Neptune Systems). Illumination was provided by metal halide (Phillips 150 W 10,000 k) and fluorescent (39 W, Phillips T5 460 nm) bulbs to create a mean light intensity of 268± 17 µmol quanta m-2 s-1 (±SE, n=64). The light intensity was selected to approximate that found at the collection depth of the parent colonies in March.

Treatments were created by blending CO2 with air, and continually assessing the mixture through an Infra Red gas analyzer (S151, Qubit Systems), which dynamically adjusted the flow of CO2 to maintain desired levels. The gas mixture was supplied through an air stone to four of the eight tanks, with others supplied with compressed air (i.e., ambient pCO2). The conditions in the tanks were analyzed for pH, salinity, temperature, and total alkalinity (TA) using standard operating procedures (Dickson et al., 2007), and the program CO2SYS (Lewis and Wallace, 1998) to calculate DIC parameters. The gas mixing technology and the methods for seawater analyses essentially are identical to those we have used before (see Dufault et al., 2012; Edmunds, 2011). In the present analysis, the calculated TA values of certified reference materials supplied by Dr. Andrew Dickson, Scripps Institute of Oceanography (batch no. 98 and 107), were determined within a mean of 1% of the certified value.

The incubation system created target temperatures of ~24.00 °C versus 30.50 °C, and target pCO2 values of ambient versus 86.1 Pa, and an irradiance of ~268 umol quanta m-2 s-1. Conditions in the tanks were measured at least daily using a certified digital thermometer (Model 15-077-8, Fisher Scientific, ±0.05 °C), a cosine-corrected quantum light meter (Li-Cor LI-192 attached to an LI-1400), and a sample ofwater withdrawn from each tank for pH, salinity, and TA analysis.

The 'ambient' and 'high' pCO2 levels: 49.4 Pa versus 86.2 Pa The 'ambient' and 'high' temperatures: 24.00 °C [ambient] versus 30.49 °C

Data also available from PANGAEA: doi:10.1594/PANGAEA.823582

Data Processing Description

BCO-DMO processing notes:

- added conventional header with dataset name, PI name, version date, reference information
- renamed parameters to BCO-DMO standard
- added lab, lat, lon columns
- columns not served: temp out, press out, pH out, fugacity, CO2 out, B Alk out, OH out, P Alk out Si Alk out, Revelle, Omega-Ca, xCO2 out.
- reformatted date from m.d.yyyy to yyyy-mm-dd removed "Tank " from data in tank column

[table of contents | back to top]

Data Files

File

brood3_chem.csv(Comma Separated Values (.csv), 3.41 KB)

Primary data file for dataset ID 535163

[table of contents | back to top]

Supplemental Files

File

Biological data for "brooded coral larvae expt. 3" datasets

filename: Cumbo etal 2012 JEMBE data1 BCODMO.xls

(Octet Stream, 154.50 KB)

MD5:e6c5e6012df9bfc581b9f769f7e52a98

Original biological data for Cumbo et al 2013 including respiration raw data, respiration by mg protein, symbiont densities, protein content, % mortality

Tank physical data

filename: Cumbo_etal_2012_JEMBE_Tank_Parameters_BCODMO.xlsx

(Octet Stream, 57.05 KB)

MD5:0170402805d7c1fe4784a51d2b26fb66

Tank physical data for "brooded coral larvae 3" experiment including seawater chemistry, light and temperature data.

[table of contents | back to top]

Related Publications

Cumbo, V. R., Fan, T. Y., & Edmunds, P. J. (2013). Effects of exposure duration on the response of Pocillopora damicornis larvae to elevated temperature and high pCO2. Journal of Experimental Marine Biology and Ecology, 439, 100–107. doi:10.1016/j.jembe.2012.10.019

Dickson, A.G., Sabine, C.L. and Christian, J.R. (Eds.) 2007. Guide to Best Practices for Ocean CO2 Measurements. PICES Special Publication 3, 191 pp https://isbnsearch.org/isbn/1-897176-07-4 Methods

Dufault, A. M., Cumbo, V. R., Fan, T.-Y., & Edmunds, P. J. (2012). Effects of diurnally oscillating pCO2 on the calcification and survival of coral recruits. Proceedings of the Royal Society B: Biological Sciences, 279(1740), 2951-2958. doi:10.1098/rspb.2011.2545

Edmunds, P. J. (2011). Zooplanktivory ameliorates the effects of ocean acidification on the reef coral Porites spp. Limnology and Oceanography, 56(6), 2402-2410. doi:10.4319/lo.2011.56.6.2402

Methods

Pierrot, D. E. Lewis, and D. W. R. Wallace. 2006. MS Excel Program Developed for CO2 System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. doi: 10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a.

[table of contents | back to top]

Related Datasets

IsRelatedTo

Edmunds, P. J., Cumbo, V. R., Fan, T. (2014) Light data in tanks from experiment on brooded coral larval, Taiwan, March 2011 (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bcodmo.org/id/dataset/535219 [view at BCO-DMO]

, Cumbo, V. R., Fan, T. (2014) Protein content of brooded coral larvae at high and ambient temperature and pCO2, March 2011 (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bco-dmo.org/id/dataset/535425 [view at Bo

Edmunds, P. J., Cumbo, V. R., Fan, T. (2014) Respiration and protein content of brooded coral larvae at high and ambient temperature and pCO2, Taiwan, March 2011 (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bco-dmo.org/id/dataset/535328 [view at BCO-DMO]

., Cumbo, V. R., Fan, T. (2014) Respiration of brooded coral larvae at high and ambient temperature and pCO2, Taiwan, March 2011 (Cumbo et al, **JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bco-dmo.org/id/dataset/535266 [view at BCO-DMO]

Edmunds, P. J., Cumbo, V. R., Fan, T. (2014) Symbiont Symbiodinium density in brooded coral larvae at high and ambient temperature and pCO2, Taiwan, March 2011 (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bco-dmo.org/id/dataset/535358 [view at BCO-DMO]

Edmunds, P. J., Cumbo, V. R., Fan, T. (2014) Temperature data from tanks from experiment on brooded coral larval, Taiwan, March 2011 (Cumbo et al, JEMBE, 2013) (MCR LTER & Climate Coral Larvae projects). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 http://lod.bco-dmo.org/id/dataset/535244 [view at BCO-DMO]

Edmunds, P. J., Cumbo, V. R., Fan, T. (2021) Settling and mortality measurements of brooded coral larvae at high and ambient temperature and pCO2, Taiwan, March 2011 (MCR LTER project, Climate_Coral_Larvae project). Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-10-07 doi:10.26008/1912/bco-dmo.535462.1 [view at BCO-DMO]

[table of contents | back to top]

Parameters

Parameter	Description	Units
lab	laboratory	unitless
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
date	local date	yyyy-mm-dd
tank	tank id number	tank
treatment	temperature: AT=ambient (24 C) or HT=high (30.49 C); pCO2 treatment: AC=ambient (419-470 uatm) or HC=high (604-742 uatm)	unitless
sal	salinity	PSU scale
treatment_temp	target temperature	degrees Celsius
temp_tank	temperature reading in the tank	degrees Celsius
TA	total alkalinity	umol/kg
pH	pH: The measure of the acidity or basicity of an aqueous solution	pH scale
pCO2	partial pressure of carbon dioxide by computation from pH and alkalinity	uatm
bicarbonate	concentration of bicarbonate ion ([HCO3]-) in seawater	umol/kg seawater
carbonate	concentration of carbonate ion ([CO3]2-)	umol/kg seawater
omega_Ara	the saturation state of seawater with respect to aragonite	unitless

[table of contents | back to top]

Instruments

Dataset-specific Instrument Name	Aquarium chiller
Generic Instrument Name	Aquarium chiller
Dataset-specific Description	Aquatech Ac11 or Shyeh Duwai Enterprise
Generic Instrument Description	Immersible or in-line liquid cooling device, usually with temperature control.

Dataset-specific Instrument Name	Gas Analyzer
Generic Instrument Name	Gas Analyzer
Dataset-specific Description	Infra Red gas analyzer (S151, Qubit Systems)
Generic Instrument Description	Gas Analyzers - Instruments for determining the qualitative and quantitative composition of gas mixtures.

Dataset-specific Instrument Name	Immersion heater	
Generic Instrument Name	Immersion heater	
Dataset-specific Description	300 Wheaters, Taikong Corporation	
Generic Instrument Description	Submersible heating element for water tanks and aquaria.	

Dataset- specific Instrument Name	LI-COR LI-192 light sensor
Generic Instrument Name	LI-COR LI-192 PAR Sensor
Dataset- specific Description	cosine-corrected quantum light meter (Li-Cor LI-192 attached to an LI-1400)
Generic Instrument Description	The LI-192 Underwater Quantum Sensor (UWQ) measures underwater or atmospheric Photon Flux Density (PPFD) (Photosynthetically Available Radiation from 360 degrees) using a Silicon Photodiode and glass filters encased in a waterproof housing. The LI-192 is cosine corrected and features corrosion resistant, rugged construction for use in freshwater or saltwater and pressures up to 800 psi (5500 kPa, 560 meters depth). Typical output is in um s-1 m-2. The LI-192 uses computer-tailored filter glass to achieve the desired quantum response. Calibration is traceable to NIST. The LI-192 serial numbers begin with UWQ-XXXX. LI-COR has been producing Underwater Quantum Sensors since 1973. These LI-192 sensors are typically listed as LI-192SA to designate the 2-pin connector on the base of the housing and require an Underwater Cable (LI-COR part number 2222UWB) to connect to the pins on the Sensor and connect to a data recording device. The LI-193 primarily in sensitivity and angular response. 193: Sensitivity: Typically 7 uA per 1000 umol s-1 m-2 in water. Azimuth: < ± 3% error over 360° at 90° from normal axis. Angular Response: < ± 4% error up to ± 90° from normal axis. 192: Sensitivity: Typically 4 uA per 1000 umol s-1 m-2 in water. Azimuth: < ± 1% error over 360° at 45° elevation. Cosine Correction: Optimized for underwater and atmospheric use. (www.licor.com)

Dataset-specific Instrument Name	Water Temp Sensor
Generic Instrument Name	Water Temperature Sensor
Dataset-specific Description certified digital thermometer (Model 15-077-8, Fisher Scientific, ±0.05 °C)	
Generic Instrument Description	General term for an instrument that measures the temperature of the water with which it is in contact (thermometer).

[table of contents | back to top]

Deployments

lab Edmunds NMMBA

Website	https://www.bco-dmo.org/deployment/58892
Platform	Natl Museum Mar. Bio. and Aquar. Taiwan
Start Date	2010-03-18
End Date	2010-03-24
Description	Experiments related to the research project: 'RUI- The ecophysiological basis of the response of coral larvae and early life history stages to global climate change' were conducted at the laboratories of the National Museum of Marine Biology and Aquarium in Southern Taiwan.

[table of contents | back to top]

Project Information

Moorea Coral Reef Long-Term Ecological Research site (MCR LTER)

Website: http://mcr.lternet.edu/

Coverage: Island of Moorea, French Polynesia

From http://www.lternet.edu/sites/mcr/ and http://mcr.lternet.edu/:

The Moorea Coral Reef LTER site encompasses the coral reef complex that surrounds the island of Moorea, French Polynesia (17°30'S, 149°50'W). Moorea is a small, triangular volcanic island 20 km west of Tahiti in the Society Islands of French Polynesia. An offshore barrier reef forms a system of shallow (mean depth ~ 5-7 m), narrow (-0.8-1.5 km wide) lagoons around the 60 km perimeter of Moorea. All major coral reef types (e.g., fringing reef, lagoon patch reefs, back reef, barrier reef and fore reef) are present and accessible by small boat.

The MCR LTER was established in 2004 by the US National Science Foundation (NSF) and is a partnership between the University of California Santa Barbara and California State University, Northridge. MCR researchers include marine scientists from the UC Santa Barbara, CSU Northridge, UC Davis, UC Santa Cruz, UC San Diego, CSU San Marcos, Duke University and the University of Hawaii. Field operations are conducted from the UC Berkeley Richard B. Gump South Pacific Research Station on the island of Moorea, French

MCR LTER Data: The Moorea Coral Reef (MCR) LTER data are managed by and available directly from the MCR project data site URL shown above. The datasets listed below were collected at or near the MCR LTER sampling locations, and funded by NSF OCE as ancillary projects related to the MCR LTER core research themes.

This project is supported by continuing grants with slight name variations:

- . LTER: Long-Term Dynamics of a Coral Reef Ecosystem
- LTER: MCR II Long-Term Dynamics of a Coral Reef Ecosystem
- LTER: MCR IIB: Long-Term Dynamics of a Coral Reef Ecosystem . LTER: MCR III: Long-Term Dynamics of a Coral Reef Ecosystem
- LTER: MCR IV: Long-Term Dynamics of a Coral Reef Ecosystem

The ecophysiological basis of the response of coral larvae and early life history stages to global climate change (Climate_Coral_Larvae)

Coverage: Moorea, French Polynesia: Southern Taiwan: California State University Northridge

Tropical coral reefs face a suite of environmental assaults ranging from anchor damage to the effects of global climate change (GCC). The consequences are evident throughout the tropics, where many coral reefs have lost a substantial fraction of their coral cover in a few decades. Notwithstanding the importance of reducing the impacts of environmental stresses, the only means by which these ecosystems can recover (or simply persist) is through the recruitment of scleractinians, which is a function of successful larval development, delivery, settlement, metamorphosis, and post-settlement events. Despite wide recognition of the importance of these processes, there are few pertinent empirical data, and virtually none that address the mechanisms mediating the success of early coral life stages in a physical environmental varying at multiple spatio-temporal

The objective of this research is to complete one of the first comprehensive ecophysiological analyses of the early life stages of corals through a description of: (1) their functionality under 'normal' conditions, and (2) their response to the main drivers of GCC. These analyses will be completed for 2 species representative of a brooding life history strategy, and the experiments will be completed in two locations, one (Taiwan) that provides unrivalled experience in coral reproductive biology, and superb microcosm facilities, and the other (Moorea), with access to a relatively pristine environment, a well described ecological and oceanographic context (through the MCR-LTER), and the capacity to bring a strong biogeographic contrast to the project. The results of the study will be integrated through modeling to explore the effects of GCC on coral community structure over the next century.

The following publications and data resulted from this project:

Wall CB, Fan TY, Edmunds PJ. Ocean acidification has no effect on thermal bleaching in the coral Seriatopora caliendrum. Coral Reefs 33: 119-130.

<u>Symbiodinium_Seriatopora photosynthesis</u> Symbiodinium_Seriatopora PI curve

Symbiodinium_Seriatopora temp-salinity-light

Symbiodinium_Seriatopora water chemistry

- Download complete data for this publication (Excel file)

2013 Wall CB, Edmunds PJ. In situ effects of low pH and elevated HCO3- on juvenile Porites spp. in Moorea, French Polynesia. Biological Bulletin 225:92-101. Data at MCR and PANGEA: doi.pangaea.de/10.1594/PANGAEA.833913

- Download complete data for this publication (Excel file)

2013 Vivian R Cumbo, Peter J Edmunds, Christopher B Wall, Tung-Yung Fan. Brooded coral larvae differ in their response to high temperature and elevated pCO2 depending on the day of release. Marine Biology DOI 10.1007/s00227-013-2280-y. Data also at PANGEA: doi.pangaea.de/10.1594/PANGAEA.831612

brooded coral larvae 2 - carbonate chemistry

brooded coral larvae 2 - larval release March 2003-2008 brooded coral larvae 2 - respiration, photosyth, mortality

- Download complete data for this publication (Excel file)

2013 Edmunds PJ, Cumbo VR, Fan TY. Metabolic costs of larval settlement and metamorphosis in the coral Seriatopora caliendrum under ambient and elevated pCO2. Journal

Experimental Marine Biology and Ecology 443: 33-38 Data also at PANGEA: doi:10.1594/PANGAEA.821644

Coral post-settlement physiology

- Download complete data for this publication (Excel file)

2013 Aaron M Dufault, Aaron Ninokawa, Lorenzo Bramanti, Vivian R Cumbo, Tung-Yung Fan, Peter J Edmunds. The role of light in mediating the effects of ocean acidification on coral calcification. Journal of Experimental Biology 216: 1570-1577.

coral-light expt.- PÁR

coral-light expt.- carbonate chemistry

coral-light expt.- temp_salinity

coral-light expt.- growth

coral-light expt.- protein

coral-light expt.- survival

- Download complete data for this publication (Excel file)

Cumbo, VR, Fan TY, Edmunds PJ. Effects of exposure duration on the response of Pocillopora damicornis larvae to elevated temperature and high pCO2. J Exp Mar Biol Ecol 439: 100-107.

Data is also at PANGEA: doi:10.1594/PANGAEA.823582

brooded coral larvae 3 - carbonate chemistry

brooded coral larvae 3 - light

brooded coral larvae 3 - mortality

brooded coral larvae 3 - protein

brooded coral larvae 3 - respiration and protein

brooded coral larvae 3 - respiration raw data

brooded coral larvae 3 - symbiont density

brooded coral larvae 3 - tank temperature

- Download part 1 of data for this publication (Excel file) - Download tank parameters data for this publication (Excel file)

2012 Cumbo, VR, Fan TY, Edmunds PI, Physiological development of brooded larvae from two pocilloporid corals in Taiwan. Marine Biology 159: 2853-2866.

brooded coral - carbonate chemistry

brooded coral - release

brooded coral - respiration

brooded coral - settlement competency

brooded coral - size_July

brooded coral - size protein_symbionts_photosynth - Download complete data for this publication (Excel file)

2012 Dufault, Aaron M; Vivian R Cumbo; Tung-Yung Fan; Peter J Edmunds. Effects of diurnally oscillating pCO2 on the calcification and survival of coral recruits. Royal Society of London (B) 279: 2951-2958. doi:10.1098/rspb.2011.2545

Data is also at PANGEA: doi:10.1594/PANGAEA.830185

recruit growth area

recruit growth weight

recruit_seawater_chemistry

recruit survival

- Download complete data for this publication (Excel file)

2011 Edmunds PJ, Cumbo V, Fan TY. Effects of temperature on the respiration of brooded larvae from tropical reef corals. Journal of Experimental Biology 214: 2783-2790.

CoralLarvae_comparison_respir

CoralLarvae release

CoralLarvae_respir CoralLarvae_size

- Download complete data for this publication (Excel file)

[table of contents | back to top]

Program Information


Long Term Ecological Research network (LTER)

Website: http://www.lternet.edu/

Coverage: United States

adapted from http://www.lternet.edu/

The National Science Foundation established the LTER program in 1980 to support research on long-term ecological phenomena in the United States. The Long Term Ecological Research (LTER) Network is a collaborative effort involving more than 1800 scientists and students investigating ecological processes over long temporal and broad spatial scales. The LTER Network promotes synthesis and comparative research across sites and ecosystems and among other related national and international research programs. The LTER research sites represent diverse ecosystems with emphasis on different research themes, and cross-site communication, network publications, and research-planning activities are coordinated through the LTER Network Office.



Site Codes

	Andrews	Forest	17	ΓER
MIND.	Allulews	FULUSI	L	ı⊏n

ARC Arctic LTER

BES Baltimore Ecosystem Stu

BLE Beaufort Lagoon Ecosystems LTER

BNZ Bonanza Creek LTER

CCE California Current Ecosystem LTER

CDR Cedar Creek Ecosystem Science Reserve

CAP Central Arizona-Phoenix LTER

CWT Coweeta LTER

FCE Florida Coastal Everglades LTER

GCE Georgia Coastal Ecosystems LTER

HFR Harvard Forest LTER

HBR Hubbard Brook LTER

JRN Jornada Basin LTER

KBS Kellogg Biological Station LTER

KNZ Konza Prairie LTER

LUQ Luquillo LTER

MCM McMurdo Dry Valleys LT

MCR Moorea Coral Reef LTEF

NWT Niwot Ridge LTER

NTL North Temperate Lakes I

NES Northeast U.S. Shelf LTE

NGA Northern Gulf of Alaska I

PAL Palmer Antarctica LTER

PIE Plum Island Ecosystems LTER

SBC Santa Barbara Coastal L

SEV Sevilleta LTER

VCR Virginia Coast Reserve L

2017 LTER research site map obtained from https://lternet.edu/site/lter-network/

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0844785

[table of contents | back to top]