

Bulk phosphorus uptake by microbes from cruises in the NW Atlantic and western Sargasso Sea 2006-2013 (Biological C:N:P ratios project)

Website: <https://www.bco-dmo.org/dataset/538027>

Data Type: Cruise Results

Version: 1

Version Date: 2014-11-19

Project

» [Biological Controls on the Ocean C:N:P ratios](#) (Biological C:N:P ratios)

Programs

» [Dimensions of Biodiversity](#) (Dimensions of Biodiversity)

» [Ocean Carbon and Biogeochemistry](#) (OCB)

Contributors	Affiliation	Role
Lomas, Michael W.	Bigelow Laboratory for Ocean Sciences	Principal Investigator
Bonachela, Juan A.	University of Strathclyde (U Strathclyde)	Co-Principal Investigator
Levin, Simon	Princeton University	Co-Principal Investigator
Martiny, Adam	University of California-Irvine (UC Irvine)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Bulk phosphorus uptake by microbes from cruises in the NW Atlantic and western Sargasso Sea 2006-2013 (Biological C:N:P ratios project).

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Related Publications](#)
 - [Related Datasets](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:55 E:-40.005 S:19.665 W:-66.097

Temporal Extent: 2006-05-19 - 2013-09-08

Dataset Description

Bulk phosphorus uptake by microbes from cruises in the NW Atlantic and western Sargasso Sea 2006-2013. This data was published in Lomas et al. 2014 PNAS.

Methods & Sampling

Sample collection:

The data presented in this study were collected on 7 cruises throughout the Western North Atlantic Ocean (cruise X0606, X0705, X0804, BVAL 39, BVAL 46, AE1206, and AE1319). Data for kinetics experiments were collected from throughout the western North Atlantic, from roughly 55°N in the Labrador Sea to ~21°N, just north of Puerto Rico. All sample depths were <200m. All samples for Pi uptake rates and kinetics experiments were collected in acid-cleaned Niskin bottles attached to a CTD rosette and kept in subdued lighting until experiments were initiated (< 1 h). Samples for whole community ambient uptake rates were collected from ~4 depths in the upper 60 m, while samples for taxon-specific ambient uptake rates were collected from 5 m, 40 m, and the deep chlorophyll maximum (DCM; ranging from 80 to 120 m) (27). *Trichodesmium* colonies were collected from the near surface (roughly within the top 20 m) by vertically hauling a handheld 100 µm net. Single colonies were transferred a second time into fresh 0.2 µm-filtered water to reduce contamination of closely associated organisms, and subsequently separated by morphotype (either 'puff' with radial trichomes or 'raft' with parallel trichomes); only data for 'rafts' are presented here.

Dissolved inorganic and organic, and particulate nutrients:

Samples for NO₃⁻/NO₂⁻, NO₂⁻ and PO₄³⁻ are gravity filtered through 0.8 µm Nucleopore polycarbonate filters using acid cleaned in-line polycarbonate filter holders, then frozen (-20°C) in HDPE bottles until analysis (Dore et al., 1996). Tests of frozen versus refrigerated samples have indicated no significant difference between storage methods (Dore et al., 1996). Nutrient samples prior to ~2003 were analyzed on a modified Technicon Autoanalyzer and samples post ~2003 were analyzed on an Alpkem Flow Solution IV; both instrumental setups have comparable sensitivity and method detection limits validated by 6-month sample overlap on both instruments. During every sample run, commercially available certified standards, OSIL and Wako Chemical, are analyzed to maintain data quality, as well as 'standard water' from 3000 m which serves as an internal standard.

Soluble reactive phosphate (SRP) concentrations in the euphotic zone of the Sargasso Sea are below analytical detection limits (~20 nmol kg⁻¹) of standard nutrient autoanalyzer configurations. To resolve the low concentrations of SRP in the surface waters at BATS the Magnesium Induced Co-precipitation method, referred to as MAGIC-SRP measurements (Karl and Tien, 1992; Rimmelin and Moutin, 2005), was used starting in late 2004.

Several modifications to the method were made and detailed in Lomas et al. (2010a). Sample accuracy was checked on each run with a certified OSIL nutrient standard. The method detection limit following this protocol is ~1 nmol kg⁻¹ with a precision of + 5% at 5 nmol kg⁻¹.

Particulate organic carbon (POC) and nitrogen (PON) samples are filtered on precombusted (450°C, 4h) Whatman GF/F filters and frozen until analysis Steinberg et al., 2001a. Samples are analyzed on a Control Equipment 240-XA or 440-XA elemental analyzer standardized to acetanilide. Particulate phosphorus samples (PPhos) are analyzed using an ash-hydrolysis method (Lomas et al., 2010a). Oxidation efficiency and standard recovery is tested with each sample run using an ATP standard solution and a certified phosphate standard (OSIL Phosphate Nutrient Standard Solution). Method precision is ~9% at 2.5 nmol kg⁻¹ (the lowest concentrations typically observed well below the euphotic zone), and ~1% at 15 nmol kg⁻¹ (typical euphotic zone concentrations). The method detection limit, defined as three times the standard deviation of the lowest standard (2.5 nmol kg⁻¹) is ~0.5 nmol kg⁻¹.

33P phosphate incubations:

The approach for ambient whole community and population-specific uptake rate measurements were previously published (Casey et al, 2009). Briefly, duplicate aliquots of 10 ml seawater were amended with 0.15 µCi (~80 pmol L⁻¹) additions of H³³PO₄ (3000 Ci mol⁻¹; PerkinElmer, USA), and incubated for 30 - 60 min in subdued lighting (~100 µmol photons m⁻² s⁻¹) at ~23°C. This temperature was within ~3°C of the coolest/warmest in situ temperature from which the samples were collected. The duration of each incubation varied depending on turnover time of the added isotope, such that efforts were made to keep uptake to <25% of the tracer added.

Duplicate killed control incubations were conducted for each station. Killed controls were amended with paraformaldehyde (0.5% final concentration) for 30 min prior to the addition of isotopic tracer and incubation. Whole community incubations were terminated by filtration onto 0.2 µm polycarbonate filters that were subsequently placed in glass scintillation vials. Population-specific ambient uptake incubations were terminated by the addition of paraformaldehyde (0.5% final concentration), and stored at 4°C until sorting (<12 h) as described in the next section.

Whole community and population-specific kinetics experiments were conducted by adding 0.15 µCi (~80 pM) of H³³PO₄ to ~10 replicate 10 ml seawater samples that were further amended by increasing additions of 'cold' KH₂PO₄ up to 100 nM. Samples were incubated as above, but the incubations were terminated by the addition of KH₂PO₄ to a final concentration of 100 µM (28). Whole community samples were filtered onto 0.2 µm polycarbonate filters, and rinsed with an oxalate wash (29). Surface bound phosphate in population-specific samples was accounted for by subtracting 33P counts for sorted populations to which 100 µM phosphate had been added prior to addition of the isotopic tracer. It is assumed that addition of such a high level of phosphate would result in negligible uptake of radioactive phosphate and thus any signal was attributed to surface absorption; this correction was always <2-3%. Population-specific kinetics experiments for samples collected in the deep chlorophyll maximum were first gravity concentrated and resuspended in phosphate-free Sargasso Sea surface water prior to incubation as described. Population-specific samples were stored at 4°C in the dark until

sorting (<3 h) as described in the next section. Kinetics experiments for *Trichodesmium* spp. were conducted in the same manner as above for whole community samples but with picked and rinsed colonies and increasing additions of 'cold' KH₂PO₄ up to 1000 nM.

Flow cytometry analysis and cell sorting:

Samples were sorted on an Influx cell sorter (BD, Seattle, WA) at an average flow rate of ~40 μ L min⁻¹. Samples were sorted for *Prochlorococcus*, *Synechococcus*, and an operationally defined eukaryotic algae size fraction (eukaryotes >2 μ m). A 100 mW blue (488 nm) excitation laser was used. After exclusion of laser noise gated on pulse width and forward scatter, autotrophic cells were discriminated by chlorophyll fluorescence (>650 nm), PE (585/30 nm), and granularity (side scatter). Sheath fluid was made fresh daily from distilled deionized water (Millipore, Billerica, MA) and molecular grade NaCl (Mallinckrodt Baker, Phillipsburg, NJ), pre-filtered through a 0.2 μ m capsule filter (Pall, East Hills, NY), and a STERIVEX sterile 0.22 μ m inline filter (Millipore, Billerica, MA). Mean coincident abort rates were < 1% and mean recovery from secondary sorts (n = 25) was 97.5 \pm 1.1% (data not shown). Spigot™ (BD Seattle, WA) and FCS Express V3™ (DeNovo Software, Seattle, WA) were used for data acquisition and post-acquisition analysis, respectively. Sorted cells from each sample were gently filtered onto 0.2 μ m Nucleopore polycarbonate filters, rinsed with copious amounts of 0.2 μ m filtered seawater, an oxalate wash(29), and placed in a 7 ml scintillation vial for liquid scintillation counting.

Data Processing Description

P uptake Rate Calculations:

Whole community and taxon-specific assimilation rates were calculated using the same equation as follows:

-- [see formula at: [https://datadocs.bco-](https://datadocs.bco-dmo.org/docs/302/lomas/DimBiodiv_CNP/data_docs/Metadata_Data_processing_section_formula.jpg)

[dmo.org/docs/302/lomas/DimBiodiv_CNP/data_docs/Metadata_Data_processing_section_formula.jpg](https://datadocs.bco-dmo.org/docs/302/lomas/DimBiodiv_CNP/data_docs/Metadata_Data_processing_section_formula.jpg)

where V_{pi} is the cell-specific utilization rate (amol 33P cell⁻¹ hr⁻¹); β_{sample} and β_{TA} are the β -emission activities (counts min⁻¹) for the sorted sample and the total activity added, respectively; n is the number of cells sorted; Delta-T is the elapsed time from 33P isotopic tracer addition to counting; T_o is the incubation duration; Lambda is the decay constant of 33P (half life = 25.4 d); P is the ambient concentration of the P source (nmol L⁻¹). The method detection limit following this protocol is ~0.5 nM with a precision of + 5% at 5 nM.

Data analysis:

Parameters for the hyperbolic nutrient uptake curves from all samples were estimated in SigmaPlot (Systat Software, San Jose, CA, Version 10) using an iteratively fit hyperbolic equation.

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- renamed parameters to BCO-DMO standard
- reformatted data from m/d/yyyy to yyyy-mm-dd
- reformatted time from H:MM to HH:MM
- replaced blank cells with nd
- changed positive longitudes to negative values

[[table of contents](#) | [back to top](#)]

Data Files

File
bulk_Puptake.csv (Comma Separated Values (.csv), 42.47 KB) MD5:ae9f516bf164a0e77eb7c87515cca814
Primary data file for dataset ID 538027

[[table of contents](#) | [back to top](#)]

Related Publications

Casey, J., Lomas, M., Michelou, V., Dyhrman, S., Orchard, E., Ammerman, J., & Sylvan, J. (2009). Phytoplankton taxon-specific orthophosphate (Pi) and ATP utilization in the western subtropical North Atlantic. *Aquatic Microbial Ecology*, 58, 31–44. doi:[10.3354/ame01348](https://doi.org/10.3354/ame01348)

General

Dore, J., Houlihan, T., Hebel, D., Tien, G., Tupas, L., and Karl, D. (1996) Freezing as a method of sample preservation for the analysis of dissolved inorganic nutrients in seawater. *Marine Chemistry* 53, 173-185. <http://soest.hawaii.edu/dkarl/misc/dave/Reprints/1996MarChem53-173-185.pdf>

Methods

Karl, D. M., & Tien, G. (1992). MAGIC: A sensitive and precise method for measuring dissolved phosphorus in aquatic environments. *Limnology and Oceanography*, 37(1), 105-116. doi:[10.4319/lo.1992.37.1.0105](https://doi.org/10.4319/lo.1992.37.1.0105)

Methods

Lomas, M. W., Bonachela, J. A., Levin, S. A., & Martiny, A. C. (2014). Impact of ocean phytoplankton diversity on phosphate uptake. *Proceedings of the National Academy of Sciences*, 111(49), 17540-17545. doi:[10.1073/pnas.1420760111](https://doi.org/10.1073/pnas.1420760111)

Methods

Lomas, M. W., Burke, A. L., Lomas, D. A., Bell, D. W., Shen, C., Dyhrman, S. T., & Ammerman, J. W. (2010). Sargasso Sea phosphorus biogeochemistry: an important role for dissolved organic phosphorus (DOP). *Biogeosciences*, 7(2), 695-710. doi:[10.5194/bg-7-695-2010](https://doi.org/10.5194/bg-7-695-2010)

Methods

Rimmelin, P., & Moutin, T. (2005). Re-examination of the MAGIC method to determine low orthophosphate concentration in seawater. *Analytica Chimica Acta*, 548(1-2), 174-182. doi:[10.1016/j.aca.2005.05.071](https://doi.org/10.1016/j.aca.2005.05.071)

Methods

[[table of contents](#) | [back to top](#)]

Related Datasets

IsRelatedTo

Lomas, M. W., Bonachela, J. A., Levin, S., Martiny, A. (2020) **Phosphorus uptake kinetics by microbes to whole communities from R/V Atlantic Explorer AE1206, AE1319 in the Sargasso Sea; Bermuda Atlantic Time-Series Station from 2012-2013 (Biological C:N:P ratios project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-11-10 doi:10.26008/1912/bco-dmo.537996.1 [[view at BCO-DMO](#)]

Lomas, M. W., Bonachela, J. A., Levin, S., Martiny, A. (2020) **Taxon-specific phosphorus uptake by microbes from NW Atlantic and western Sargasso Sea from 2007-2013 (Biological C:N:P ratios project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2014-11-10 doi:10.26008/1912/bco-dmo.538053.1 [[view at BCO-DMO](#)]

Lomas, M. W., Bonachela, J. A., Levin, S., Martiny, A. (2021) **Biogeochemistry of microbial phosphorus uptake from cruises in the Sargasso Sea; Bermuda Atlantic Time-Series Station from 2011-2013 (Biological C:N:P ratios project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 2) Version Date 2021-02-05 doi:10.26008/1912/bco-dmo.538091.2 [[view at BCO-DMO](#)]

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
cruise_id	cruise identification	unitless
station	station number or identification	unitless
cast	cast number	unitless
date	date - local or GMT?	yyyy-mm-dd
time_gmt	GMT time	HH-MM
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
bottle	bottle number	unitless
depth	depth of sample	meters
SRP	Soluble Reactive Phosphorus	nM
P33_PO4_uptake	33P phosphate uptake rate	nmol/L/h
year	sampling year	unitless
mon_start	month of start of cruise	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Niskin bottle
Generic Instrument Name	Niskin bottle
Dataset-specific Description	Acid washed Niskin bottles attached to a CTD rosette
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset-specific Instrument Name	Nutrient Autoanalyzer
Generic Instrument Name	Nutrient Autoanalyzer
Dataset-specific Description	Modified Technicon Autoanalyzer and an Alpkem Flow Solution IV
Generic Instrument Description	Nutrient Autoanalyzer is a generic term used when specific type, make and model were not specified. In general, a Nutrient Autoanalyzer is an automated flow-thru system for doing nutrient analysis (nitrate, ammonium, orthophosphate, and silicate) on seawater samples.

Dataset-specific Instrument Name	Plankton Net
Generic Instrument Name	Plankton Net
Dataset-specific Description	Handheld 100 µm net
Generic Instrument Description	A Plankton Net is a generic term for a sampling net that is used to collect plankton. It is used only when detailed instrument documentation is not available.

[[table of contents](#) | [back to top](#)]

Deployments

AE1319

Website	https://www.bco-dmo.org/deployment/537979
Platform	R/V Atlantic Explorer
Report	http://dmoserv3.whoi.edu/data_docs/Bio_CNP_Ratios/AE1319_Cruise_Report_09182013_reduced2.pdf
Start Date	2013-08-14
End Date	2013-09-11
Description	Cruise for project 'Dimensions of Biodiversity: Biological Controls on the Ocean C:N:P ratios'.

X0606

Website	https://www.bco-dmo.org/deployment/58060
Platform	R/V Atlantic Explorer
Start Date	2006-05-19
End Date	2006-05-27
Description	One in a series of transect cruises to study the biological and biogeochemical aspects of the marine phosphorus cycle.

BVAL39

Website	https://www.bco-dmo.org/deployment/538365
Platform	R/V Atlantic Explorer
Start Date	2007-10-21
End Date	2007-10-29
Description	BATS Validation stations

AE0810

Website	https://www.bco-dmo.org/deployment/58062
Platform	R/V Atlantic Explorer
Start Date	2008-05-03
End Date	2008-05-25
Description	One in a series of transect cruises to study the biological and biogeochemical aspects of the marine phosphorus cycle. Note the cruise identifiers for the Atlantic Explorer were originally formatted as XYY## (e.g. X0806 was the 6th cruise in 2008). The data files include cruise IDs of this type. The vessel operator changed the cruise ID syntax several years after the cruise and the official cruise ID syntax was changed to AEYY##. For example, AE0810 should be the same cruise as X0810. One exception for this dataset is that X0804 is cruise ID AE0810 (unclear how the cruise numbering scheme got so confused). Database validation showed that AE0804 was not the correct cruiseid based on information at R2R. The cruiseid was then updated to reflect the corrected information (the May 2008 cruise was AE0810. Additional Information from R2R Site

BVAL37

Website	https://www.bco-dmo.org/deployment/540202
Platform	R/V Atlantic Explorer
Start Date	2006-10-18
End Date	2006-10-29
Description	BATS Validation stations

X0619

Website	https://www.bco-dmo.org/deployment/540209
Platform	R/V Atlantic Explorer
Start Date	2006-11-21
End Date	2006-11-22

X0705

Website	https://www.bco-dmo.org/deployment/58061
Platform	R/V Atlantic Explorer
Start Date	2007-06-02
End Date	2007-06-14
Description	One in a series of transect cruises to study the biological and biogeochemical aspects of the marine phosphorus cycle.

[[table of contents](#) | [back to top](#)]

Project Information

Biological Controls on the Ocean C:N:P ratios (Biological C:N:P ratios)

Coverage: western North Atlantic; 60N to 20N along 66W longitude; 20N to 15S in the tropical Pacific

One of the fundamental patterns of ocean biogeochemistry is the Redfield ratio, linking the stoichiometry of surface plankton with the chemistry of the deep ocean. There is no obvious mechanism for the globally consistent C:N:P ratio of 106:16:1 (Redfield ratio), especially as there is substantial elemental variation among plankton

communities in different ocean regions. Thus, knowing how biodiversity regulates the elemental composition of the ocean is important for understanding the ocean and climate as a whole -- now and in the future.

The conceptual hypotheses for this study are as follows: 1. The C:N:P ratio of a cell is constrained by its broad taxonomic group, which determines, for example, whether it has an outer shell, its size, functional metabolism, membrane lipid composition. 2. Within a taxon, there is high genetic diversity. Some of this genetic diversity is potentially laterally transferred, or can be lost within taxa, and confers various functional abilities (organic phosphate assimilation, nitrate assimilation, photoheterotrophy, etc.). Functional diversity provides the cell with further flexibility, such as the ability to respond to varying nutrient supply rates/ratios, and affects a cell's C:N:P ratio within the range specified by the taxon. 3. Given these taxonomic and genetic constraints, a cell is physiologically plastic and modifies how it allocates cellular resources in response to nutrient supply rates/ratios in the environment. 4. The microbial diversity (taxonomic, genetic, and functional) of the surface ocean varies over time and space, driven by many factors in addition to nutrients. The sum of this mixture composes the ecosystem C:N:P, the ratio that Redfield described.

Based on this framework, the CoPIs will make field observations of taxon-specific stoichiometry and growth rates, genomic analyses, and conduct laboratory chemostat experiments to improve understanding of how ocean taxonomic, genetic, and functional biodiversity control the stoichiometry of the surface ocean plankton. Their analyses of these data would lead to a mechanistic understanding of variations in the Redfield ratio, both spatially and temporally.

This study will greatly expand knowledge of the genomic diversity among ocean microbes and how this diversity affects biogeochemistry. The stoichiometry of the ocean's microbes is a parameter that nearly every chemical or biological oceanographer uses, from converting measurements made in one element to another, to estimating regional and global nitrogen budgets. The research also has important implications for the global carbon budget and any changes that might result from climate change.

To understand mechanistically temporal and spatial variability of the plankton C:N:P ratio, biodiversity must be studied not only at the traditional taxonomic level, but at the genetic and functional levels which dictate organism response to their environment. Data will be integrated into a combined ocean ecological, evolutionary, and biogeochemical model, with flexible stoichiometry, including cellular biochemical allocations. Seeding a coupled physical-biological model of the oceans with multiple competing genotypes enables the exploration of ecological and evolutionary patterns of resource acquisition and C:N:P ratios. Developing a more mechanistic examination of the course of ecology and evolution, in which laboratory and field data define tradeoffs between different growth and nutrient acquisition strategies, would establish the framework of adaptive dynamics for determining "evolutionarily convergence". Finally, model outcomes will be evaluated against field data.

The field work planned for this project includes several cruises: BV46 (September/October 2011), BV48 (September 2012), a June 2013 cruise from Bermuda to the Labrador Sea, and a cruise from Hawaii to Tahiti (May 2014). Additionally, samples will be acquired during cruises of opportunity.

[[table of contents](#) | [back to top](#)]

Program Information

Dimensions of Biodiversity (Dimensions of Biodiversity)

Website: http://www.nsf.gov/funding/pgm_summ.jsp?pims_id=503446

Coverage: global

(adapted from the NSF Synopsis of Program)

Dimensions of Biodiversity is a program solicitation from the NSF Directorate for Biological Sciences. FY 2010 was year one of the program. [\[MORE from NSF\]](#)

The NSF Dimensions of Biodiversity program seeks to characterize biodiversity on Earth by using integrative, innovative approaches to fill rapidly the most substantial gaps in our understanding. The program will take a broad view of biodiversity, and in its initial phase will focus on the integration of genetic, taxonomic, and functional dimensions of biodiversity. Project investigators are encouraged to integrate these three dimensions to understand the interactions and feedbacks among them. While this focus complements several core NSF programs, it differs by requiring that multiple dimensions of biodiversity be addressed simultaneously, to

understand the roles of biodiversity in critical ecological and evolutionary processes.

Ocean Carbon and Biogeochemistry (OCB)

Website: <http://us-ocb.org/>

Coverage: Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO₂ and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1046001
NSF Division of Ocean Sciences (NSF OCE)	OCE-1046368
NSF Division of Ocean Sciences (NSF OCE)	OCE-1046297
NSF Division of Ocean Sciences (NSF OCE)	OCE-1045966

[[table of contents](#) | [back to top](#)]