CTD data from gliders on the R/V Savannah in the South Atlantic Bight (SAB) continental shelf off Long Bay (-79W, 32N; -77W, 34 N) collected from January to April 2012 (Long Bay Wintertime Bloom project)

Website: https://www.bco-dmo.org/dataset/544615 Data Type: Cruise Results Version: 1 Version Date: 2014-12-18

Project

» <u>Mechanisms of nutrient input at the shelf margin supporting persistent winter phytoplankton blooms</u> <u>downstream of the Charleston Bump</u> (Long Bay Wintertime Bloom)

Contributors	Affiliation	Role
<u>Nelson, James</u>	Skidaway Institute of Oceanography (SkIO)	Principal Investigator
Edwards, Catherine	Skidaway Institute of Oceanography (SkIO)	Co-Principal Investigator
<u>Seim, Harvey E.</u>	University of North Carolina at Chapel Hill (UNC-Chapel Hill)	Co-Principal Investigator
<u>Haines, Sara</u>	University of North Carolina at Chapel Hill (UNC-Chapel Hill)	Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Two Webb Slocum gliders were used to measure salinity, temperature, and depth. Glider 'Pelagia' was deployed in Long Bay, S. Carolina, South Atlantic Bight and glider 'Ramses' was deployed along the upper slope of S. Carolina. in 2012. Data were collected in a time series from January to April, 2012.

Table of Contents

- <u>Coverage</u>
- Dataset Description
- <u>Data Processing Description</u>
- Data Files
- <u>Related Publications</u>
- <u>Parameters</u>
- Instruments
- Deployments
- <u>Project Information</u>
- <u>Funding</u>

Coverage

Spatial Extent: N:32.95739 E:-78.05085 S:32.84347 W:-78.14255 Temporal Extent: 2012-01-25

Data Processing Description

Data was read from glider stored memory after recovery.

Quality control (QC) steps applied to CTD data (bad flag means values were replaced with NaN in the matlab files):

1. data when glider was in testing modes or at the surface were bad flagged.

- 2. obvious spikes in temperature and conductivity were bad flagged based on a difference threshold
- 3. salinity errors were significant due to the unpumped conductivity cell and were corrected following Garau et

al., 2011. Matlab scripts available at http:// <u>www.socib.es/;glider/doco/gliderToolbox/ctdTools/</u> thermalLagTools) were used with modified parameters as specified in the data file under

config.thermal_lag_correction_parameters to create salinCorrected and densCorrected. 4. poor flushing of the conductivity cell and thus invalid salinity data were found to occur when the glider forward speed fell below 10 cm/s or the pitch was between +5 and +15 degrees (on an upcast). All CTD output values were bad flagged during these times.

Salinity (salin) was calculated from conductivity using PSS-78, as well as water density (dens) using the seawater routines.

Normally, density should not be archived with data submission to national repository. However since this parameter is integral to the main study, density is being reported with other measured parameters but with caveat that it is derived based on older conventions.

BCO-DMO Processing:

- -extracted data from MatLab .mat files
- -added conventional header with dataset name, PI name, version date, reference information
- -renamed parameters to BCO-DMO standard
- -added yrday_utc and ISO_DateTime_UTC to served view
- -reduced number of significant digits

[table of contents | back to top]

Data Files

File

glider_CTD_jgof.csv(Comma Separated Values (.csv), 418.52 MB) MD5:3ba63383fbc333c3313fcb5b3f59ca1c

Primary data file for dataset ID 544615

[table of contents | back to top]

Related Publications

Garau, B., Ruiz, S., Zhang, W. G., Pascual, A., Heslop, E., Kerfoot, J., & Tintoré, J. (2011). Thermal Lag Correction on Slocum CTD Glider Data. Journal of Atmospheric and Oceanic Technology, 28(9), 1065–1071. doi:10.1175/jtech-d-10-05030.1 <u>https://doi.org/10.1175/JTECH-D-10-05030.1</u> *Methods*

[table of contents | back to top]

Parameters

Parameter	Description	Units
glider_id	glider identification	unitless
year	year	уууу
month_utc	UTC month	1-12
day_utc	UTC day	1-31
yrday_utc	UTC day and decimal time: e.g. 326.5 for the 326th day of the year or November 22 at 1200 hours (noon)	unitless
hour	UTC hour	НН
min	UTC minute	ММ
sec	UTC second	SS.fraction of second
lat	latitude; north is positive	decimal degrees
lon	longitude; east is positive	decimal degrees
depth	depth	meters
press	pressure	decibars
temp	temperature	degrees Celsius
dens_corr	corrected density	kilograms/meter^3
sal_corr	corrected salinity	PSU
ISO_DateTime_UTC	Date/Time (UTC) ISO formatted	yyyy-mm- ddTHH:MM:SS[.xx]Z
lat_start	latitude at start of glider deployment.	decimal degrees
lon_start	longitude at start of glider deployment; east is positive.	decimal degrees

Instruments

Dataset- specific Instrument Name	CTD Sea-Bird
Generic Instrument Name	CTD Sea-Bird
Dataset- specific Description	The CTD is a component of the Slocum glider used for this dataset.
Generic Instrument Description	Conductivity, Temperature, Depth (CTD) sensor package from SeaBird Electronics, no specific unit identified. This instrument designation is used when specific make and model are not known. See also other SeaBird instruments listed under CTD. More information from Sea-Bird Electronics.

Dataset- specific Instrument Name	GPS
Generic Instrument Name	Global Positioning System Receiver
Dataset- specific Description	GPS is a component of the Slocum glider used for this dataset.
Generic Instrument Description	The Global Positioning System (GPS) is a U.S. space-based radionavigation system that provides reliable positioning, navigation, and timing services to civilian users on a continuous worldwide basis. The U.S. Air Force develops, maintains, and operates the space and control segments of the NAVSTAR GPS transmitter system. Ships use a variety of receivers (e.g. Trimble and Ashtech) to interpret the GPS signal and determine accurate latitude and longitude.

[table of contents | back to top]

Deployments

SAV-12-03

Website	https://www.bco-dmo.org/deployment/58863
Platform	R/V Savannah
Start Date	2012-01-24
End Date	2012-02-02
Description	Glider deployments (Ramses and Pelagia), glider monitoring; Acrobat (towed package) surveys of 3 cross-shelf sections (25 km along-shelf spacing, mid-shelf to upper slope); CTD profiles for hydrography, bio-optical, oxygen; station sampling for chlorophyll, nutrients, flow cytometry; Deployment of 2 moorings off Georgia on return leg to SkIO (75 m and 30 m). Cruise information and original data are available from the NSF R2R data catalog.

LB_2012_Pelagia-1

Website	https://www.bco-dmo.org/deployment/544628
Platform	AUV Slocum Glider Pelagia
Start Date	2012-01-26
End Date	2012-02-14
Description	CTD measurements along cross-shelf transect in Long Bay off South Carolina, USA. Glider Pelagia deployed on 2012-01-26 on SAV-12-03 and recovered on 2012-02-14 on SAV-12-05.

LB_2012_Pelagia-2

Website	https://www.bco-dmo.org/deployment/544629
Platform	AUV Slocum Glider Pelagia
Start Date	2012-02-16
End Date	2012-03-08
Description	CTD measurements along undulating transect. Cross-shelf transect in Long Bay off South Carolina, USA 78 20.007 W, 33 10.172 N. Deployed on 2012-02-16 on SAV-12-05 and recovered on 2012-03-08 on SAV-12-11.

LB_2012_Pelagia-3

Website	https://www.bco-dmo.org/deployment/544635
Platform	AUV Slocum Glider Pelagia
Start Date	2012-03-16
End Date	2012-04-04
Description	Cross-shelf transect in Long Bay off South Carolina, USA. Glider Pelagia was deployed on 2012-03-16 on SAV-12-11, and recovered on 2012-04-04 on SAV-12-14.

LB_2012_Ramses-1

Website	https://www.bco-dmo.org/deployment/544639
Platform	AUV Slocum Glider Ramses
Start Date	2012-01-26
End Date	2012-02-14
Description	CTD measurements along upper slope off South Carolina, USA. Glider Ramses was deployed on 2012-01-26 on SAV-12-03, and recovered on 2012-02-14 on SAV-12-05.

LB_2012_Ramses-2

Website	https://www.bco-dmo.org/deployment/544643
Platform	AUV Slocum Glider Ramses
Start Date	2012-02-15
End Date	2012-03-14
Description	CTD measurements along upper slope off South Carolina, USA. Glider Ramses was deployed on 2012-02-05 on SAV-02-15, and recovered on 2012-03-14 on SAV-12-11.

Website	https://www.bco-dmo.org/deployment/544647
Platform	AUV Slocum Glider Ramses
Start Date	2012-03-16
End Date	2012-04-03
Description	CTD measurements along upper slope off South Carolina, USA. Glider Ramses was deployed on 2012-03-16 on SAV-12-11, and recovered on 2012-04-03 on SAV-12-14.

SAV-12-05

Website	https://www.bco-dmo.org/deployment/58864
Platform	R/V Savannah
Start Date	2012-02-13
End Date	2012-02-24
Description	Glider recoveries (Ramses, Pelagia), shipboard replacement of batteries, ballast adjustments, then both units were redeployed. Survey work was conducted using the towed Acrobat package and CTD station section from upper slope to shelf. Deck incubation experiment for primary production, station sampling for various properties. Due to gale-force winds, the ship ran into Wilmington (docked at Cape Fear Community College) on 18 February. When heading back out on 20 February for an Acrobat survey and station work, a problem with a shaft coupling forced return to Wilmington on 21 February for repair work. When repairs were completed, strong SW winds had developed and were forecast to continue for several days, so offshore work was not possible the ship returned to SkIO along the coast. Cruise information and original data are available from the NSF R2R data catalog.

SAV-12-11

Website	https://www.bco-dmo.org/deployment/58865	
Platform	R/V Savannah	
Start Date	2012-03-13	
End Date	2012-03-22	
Description	Glider recoveries (Ramses, Pelagia), then battery replacement, reballasting, redeployment (at LB2). Acrobat surveys, upper slope to outer shelf CTD surveys, station sampling for water samples and primary productivity experiments (deck incubations and 'photosynthetron'). Attempted to survey a subsurface bloom in the mid-shelf (apparently Phaeocystis). During an Acrobat survey, the tow cable failed over upper slope. The Acrobat package was located with an acoustic range-finder but was too deep for a grapple attempt. With the very mild winter, shelf conditions were already post-winter (warm water across outer to mid-shelf) and not favorable for the winter bloom formation. Given the conditions the cruise was cut short by a few days. Cruise information and original data are available from the NSF R2R data catalog.	

SAV-12-14

Website	https://www.bco-dmo.org/deployment/58866	
Platform	R/V Savannah	
Start Date	2012-04-02	
End Date	2012-04-05	
Description	Recovery of moored instrument packages at LB3 (150 kHz ADCP in large TRBM pod/frame, MicroCAT, ECO FLNTU fluorometer/turbidity), LB2 (ADCP, MicroCAT, ECOFLNTU), LB1 (two moorings – ADCP; Taut-line package with HOBO T-chain [two with C], 3 ECO FLNTU units); Recovery of Ramses and Pelagia gliders; CTD section from upper slope to outer shelf (stations at 5 km intervals, LB+60 to LB+30, also at LB2 site), with samples for nutrients at selected stations/depths, some chlorophyll samples; Inter-calibration of the ECO FLNTU fluorometer/turbidity instruments after recovery of moorings (mounted on CTD/carousel for co-located sampling at two depths. This is the last CTD station). Cruise information and original data are available from the NSF R2R data catalog.	

[table of contents | back to top]

Project Information

Mechanisms of nutrient input at the shelf margin supporting persistent winter phytoplankton blooms downstream of the Charleston Bump (Long Bay Wintertime Bloom)

Website: http://nccoos.org/projects/long-bay-wintertime-blooms/

Coverage: outer South Atlantic Bight (SAB) continental shelf off Long Bay

NSF Project Title: Mechanisms of nutrient input at the shelf margin supporting persistent winter phytoplankton blooms downstream of the Charleston Bump

Sustained phytoplankton blooms along the outer South Atlantic Bight (SAB) continental shelf off Long Bay are observed in winter in multi-year satellite chlorophyll imagery. This section of the shelf lies north of the "Charleston Bump" (between 32.5-33.5°N), where the Gulf Stream is often strongly deflected offshore. Due to this offshore deflection, this is not an area where nutrient input to the shelf would be enhanced by upwelling associated with Gulf Stream frontal eddies, a major mechanism of nutrient input in other parts of the SAB shelf (Lee et al., 1991). Yet prior in situ observations suggest that there is recurring input of nutrients from the upper slope to the outer shelf off Long Bay from winter to early spring. This project will investigate a fundamental aspect of physical-biological coupling in the outer shelf to upper slope region. The PIs will test the hypotheses that: 1) the persistence of winter blooms on the outer shelf off Long Bay results from repeated episodes of nutrient input and mixing which maintains nutrient-sufficient conditions for extended periods; 2) several physical mechanisms are involved, including enhanced mixing energy from the internal tide along this section of the upper slope/shelf break; 3) the relatively high nutrient, intermittently turbulent environment will favor larger bloom-forming phytoplankton. The latter could have important implications for higher trophic levels, including early life history strategies of fish that spawn along the shelf margin off Long Bay in winter to early spring.

This project will combine several maturing observational technologies to address the following:

 What is the frequency and magnitude on on-shelf transport of nitrate from the upper slope?
What are the mechanisms of nutrient delivery from the upper slope to the outer continental shelf zone that are operating off Long Bay under the range of hydrographic and forcing conditions encountered in winter?
What is the 3-D structure of outer shelf hydrography and associated winter bloom features and how do these evolve through multiple nutrient input/mixing events?

4. What are the rates of nitrate utilization and primary production associated with the winter blooms?

5. Does the winter regime consistently favor a bloom assemblage dominated by larger diatom forms?

Near-continuous cross-shelf and upper slope observations will be obtained with two autonomous gliders, timeseries measurements on the outer shelf and slope from a set of moored instruments (including a moored profiling system at the shelf break), and repeated cross- and along-shelf ship surveys using a towed, undulating package. Ship station work will include measurements of primary production and on-board analyses of key functional characteristics of the phytoplankton assemblage (cell forms, abundance, size and bio-volume distributions) using a microfluidics/imaging system. In combination, these systems will provide a level of spatial and temporal resolution of physical, nutrient and biological fields that could not be achieved in earlier, stationbased field studies and the basis for improved understanding of physical mechanisms of recurring nutrient input to the shelf, and how the nutrient, mixing, and circulation regime in winter structures the phytoplankton community. Coastal naturalists will be engaged through a seabird survey component of the field program that will augment existing information on pelagic seabirds in winter and define their association with oceanographic features on the central South Atlantic Bight shelf and slope.

This project will provide a deeper understanding of shelf/slope exchange processes and how these influence shelf ecosystems, generating information that will contribute to implementation of ecosystem-based management in the region.

References:

Lee, T. N., J. A. Yoder, and L. P. Atkinson, 1991: Gulf Stream frontal eddy influence on productivity of the southeast U.S. continental shelf. J. Geophys. Res, 96, 22191-22205.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1032285</u>
NSF Division of Ocean Sciences (NSF OCE)	OCE-1032276

[table of contents | back to top]