

GenBank BLAST results of copepod nauplii gut contents analyzed at San Francisco State University in 2013 (Food Limitation in Copepod nauplii project)

Website: <https://www.bco-dmo.org/dataset/546671>

Data Type: experimental

Version: 2015-01-15

Project

» [Feeding and food limitation in copepod nauplii, the neglected life stage](#) (food limitation in copepod nauplii)

Contributors	Affiliation	Role
Kimmerer, William	San Francisco State University (SFSU)	Principal Investigator
Cohen, Sarah	San Francisco State University (SFSU)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Field collected copepod nauplii gut content sequences were identified using the BLASTN algorithm on GenBank or by comparison to a sequence library of local copepod species constructed for this project. Blast searches of the GenBank database were conducted on May 16, 2013. BLASTed sequences were identified to the taxonomic level at which there was consensus among matches within 10% of the best score.

Related Reference:

* Craig, Carrie, Wim J. Kimmerer, and C. Sarah Cohen. 2014. A DNA-based method for investigating feeding by copepod nauplii. *Journal of Plankton Research* 36 (1): 271-275

Vogt, R.A., T.R. Ignoffo, L.J. Sullivan, J. Herndon, J.H. Stillman, and W. Kimmerer. 2013. Feeding capabilities and limitations in the nauplii of two pelagic estuarine copepods, *Pseudodiaptomus marinus* and *Oithona davisae*. *Limnology and Oceanography* 58: 2145-2157.

Methods & Sampling

See [Methodology](#) (pdf).

Data Processing Description

BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date, reference information
- renamed parameters to BCO-DMO standard

- replaced blanks with underscores
- moved sequence id's to separate column
- replaced '?' with '_maybe'
- changed Pseudodiaptomus to Pseudodiaptomus

[[table of contents](#) | [back to top](#)]

Data Files

File
naup_blast.csv (Comma Separated Values (.csv), 2.12 MB) MD5:507a8ae381af8b3e6367d14fdf0e1c65
Primary data file for dataset ID 546671

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
sequence_id	sequence identification: sample name followed by sequence number. TDF=adult; TDN = nauplii	unitless
GenBank_id	GenBank accession number of BLAST hit	unitless
BLAST_score	BLAST score: statistical significance of sequence matches to GenBank database	unitless
domain	taxonomic domain: Bacteria or Eukaryota	unitless
tax_rank_1	highest taxonomic name	unitless
tax_rank_2	taxonomic name; more specific than preceding	unitless
tax_rank_3	taxonomic name; more specific than preceding	unitless
tax_rank_4	taxonomic name; more specific than preceding	unitless
tax_rank_5	taxonomic name; more specific than preceding	unitless
tax_rank_6	taxonomic name; more specific than preceding	unitless
tax_rank_7	taxonomic name; more specific than preceding	unitless
tax_rank_8	taxonomic name; more specific than preceding	unitless
tax_rank_9	taxonomic name; more specific than preceding	unitless
tax_rank_10	taxonomic name; more specific than preceding	unitless
tax_rank_11	taxonomic name; more specific than preceding	unitless
tax_rank_12	taxonomic name; more specific than preceding	unitless
tax_rank_13	taxonomic name; more specific than preceding	unitless
tax_rank_14	taxonomic name; more specific than preceding	unitless
tax_rank_15	taxonomic name; more specific than preceding	unitless
tax_rank_16	taxonomic name; more specific than preceding	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Automated Sequencer
Generic Instrument Name	Automated DNA Sequencer
Dataset-specific Description	ABI 3130 Genetic Analyzer
Generic Instrument Description	General term for a laboratory instrument used for deciphering the order of bases in a strand of DNA. Sanger sequencers detect fluorescence from different dyes that are used to identify the A, C, G, and T extension reactions. Contemporary or Pyrosequencer methods are based on detecting the activity of DNA polymerase (a DNA synthesizing enzyme) with another chemoluminescent enzyme. Essentially, the method allows sequencing of a single strand of DNA by synthesizing the complementary strand along it, one base pair at a time, and detecting which base was actually added at each step.

Dataset-specific Instrument Name	HPLC
Generic Instrument Name	High-Performance Liquid Chromatograph
Generic Instrument Description	A High-performance liquid chromatograph (HPLC) is a type of liquid chromatography used to separate compounds that are dissolved in solution. HPLC instruments consist of a reservoir of the mobile phase, a pump, an injector, a separation column, and a detector. Compounds are separated by high pressure pumping of the sample mixture onto a column packed with microspheres coated with the stationary phase. The different components in the mixture pass through the column at different rates due to differences in their partitioning behavior between the mobile liquid phase and the stationary phase.

Dataset-specific Instrument Name	Plankton Net
Generic Instrument Name	Plankton Net
Dataset-specific Description	Mesh size 100 microns or 150 microns
Generic Instrument Description	A Plankton Net is a generic term for a sampling net that is used to collect plankton. It is used only when detailed instrument documentation is not available.

Dataset-specific Instrument Name	Thermal Cyclers
Generic Instrument Name	Thermal Cyclers
Generic Instrument Description	A thermal cycler or "thermocycler" is a general term for a type of laboratory apparatus, commonly used for performing polymerase chain reaction (PCR), that is capable of repeatedly altering and maintaining specific temperatures for defined periods of time. The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. They can also be used to facilitate other temperature-sensitive reactions, including restriction enzyme digestion or rapid diagnostics. (adapted from http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html)

[[table of contents](#) | [back to top](#)]

Deployments

Kimmerer_2013

Website	https://www.bco-dmo.org/deployment/546436
Platform	SFSU RTC
Start Date	2009-09-01
End Date	2014-08-31
Description	Copepod feeding studies

[[table of contents](#) | [back to top](#)]

Project Information

Feeding and food limitation in copepod nauplii, the neglected life stage (food limitation in copepod nauplii)

Coverage: San Francisco Estuary

This project will investigate feeding by copepod nauplius larvae, the most abundant metazoans in the sea. It will answer three questions: 1) How does food selection by adults and nauplii differ when they are fed multiple prey species in the laboratory? 2) How does food selection by adults and nauplii differ when they are feeding on natural prey assemblages? and, 3) How do growth, development, and survival differ between copepodites and nauplii when their growth is food limited? Comparative experiments and field-based measurements will contrast the food consumed, and the effects of food limitation, between nauplii and later life stages. This contrast will include attributes of food such as size, taxon, and motility, and will include experiments with cultured prey offered singly or in a mixture, and natural prey, and apply genetic techniques to determine prey consumption by a predatory copepod. Copepods will be collected from the San Francisco Estuary, with four species selected for experiments to span taxonomic groups, sizes, salinity ranges, and general feeding behavior. A variety of techniques will be applied to account for the inevitable biases and limitations of each; all but one have previously been applied in our laboratories. These will include laboratory feeding experiments using cultured prey individually and in mixtures, and experiments using natural prey. Consumption of prey in experimental bottles will be measured as chlorophyll concentration and through particle counts by microscopy and flow cytometry. Radioactively labeled prey will be used in short incubations to determine feeding on particular prey types. Samples from the field will be examined for gut fluorescence. Separate experiments will determine how nauplii and copepodites survive and grow at different concentrations of food. Investigations of feeding by a

predatory copepod (*Tortanus dextrilobatus*) will use molecular techniques to identify mitochondrial and nuclear DNA from diverse suspected prey species. Specific primers will be developed for common zooplankton species consumed by *T. dextrilobatus* in the laboratory. General primers and screening protocols developed here will be useful for identifying food web interactions in other estuarine communities.

Copepod nauplii are important both in their diverse trophic roles in ocean foodwebs and in the population dynamics of copepods. Nauplii have a completely different feeding apparatus from later stages, and the first feeding stage can be very sensitive to starvation, making these life stages critical to population dynamics. Yet extant copepod population models treat nauplii as miniature adults. This work will provide valuable input to the growing efforts at modeling ocean ecosystems.

[Further details from final report \(pdf\)](#)

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0929075

[[table of contents](#) | [back to top](#)]