

Population analysis and metagenomics of Japanese tsunami marine debris mussels along the Hawaii, Washington and Oregon coasts in 2012 (JTMD-BF project)

Website: <https://www.bco-dmo.org/dataset/552349>

Version: 2015-02-27

Project

» [Testing the Invasion Process: Survival, Dispersal, Genetic Characterization and Attenuation of Marine Biota on the 2011 Japanese Tsunami Marine Debris Field.](#) (JTMD-BF)

| Contributors | Affiliation | Role |
|----------------------------------|---|---------------------------|
| Geller, Jonathan | Moss Landing Marine Laboratories (MLML) | Co-Principal Investigator |
| Copley, Nancy | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

A major objective of the JTMD project was to characterize the biodiversity of the arriving and landed JTMD fauna and flora. One constituent of the landed biota were mussels in the genus *Mytilus*, known to be member of a species complex (including species native to western North America). We therefore undertook genetic analysis of *Mytilus* contained on JTMD for purposes of species identification. To augment morphological assessment of diversity of JTMD fouling organisms, we performed metabarcoding, in which bulk samples were extracted for DNA and DNA barcoding markers sequenced in parallel.

Access to this dataset is temporarily RESTRICTED until publication. Please contact the PI for further information

Methods & Sampling

Mytilus specimens were collected from Japanese tsunami marine debris (JTMD) from 2012-2014 at coastal sites in Washington, Oregon and Hawaii.

Population level analysis of focal JTMD species. *Mytilus* spp. were one of the most common organisms on JTMD. Cytochrome c oxidase subunit III was chosen for species identification of mussels because available female-specific primers allowed a simpler approach to species discrimination than the standard COI primers, which amplify both male and female sequences and complicate analyses. The proposed method of pooling tissues for bulk DNA extraction and post-sequencing analysis of haplotype variation was set aside due to presumed sequence error observed when sequencing individual mussel amplicons. Instead, 695 mussels were extracted for DNA and sequenced individually on the Ion Torrent platform. Addition of index nucleotides to PCR products allowed post-sequencing separation of sequences by source mussel. Bins of reads from individual mussels were mapped to reference sequences of female *M. galloprovincialis*, *M. trossulus*, and *M. coruscus*. Known distribution of *Mytilus* suggested that mussels in Japan would be *M. galloprovincialis*. A sample of mussels for which Ion Torrent sequencing resulted in identification as *M. trossulus* or *M. coruscus* were reanalyzed by preparation of new PCR product and Sanger sequencing.

Metagenomic analysis of JTMD assemblages. Four 15 cm² samples were scraped from the sides and top of the floating dock "Misawa 1" that beached near Newport Oregon in June 2012. Samples were preserved in ethanol and shipped to Moss Landing Marine Laboratories. Entire samples were decanted of ethanol and homogenized in an IKE Basic Mill. Ten grams of each sample were extracted with the Mo Bio Powersoil Mega DNA extraction kit (Mo Bio, Carlsbad, CA). Cytochrome c oxidase subunit I was PCR amplified, and prepared for sequencing on the Ion Torrent PGM. Resulting reads were pooled, quality and size filtered, and assembled into operational taxonomic units (OTUs) using the Velvet assembler. Gene OTUs were compared to eukaryotic COI sequences culled from Genbank by using variants of the search term for COI (COI, COX1, Cytochrome oxidase subunit I, Cytochrome c oxidase subunit I, etc...) and excluding prokaryota. We also specifically searched for COI sequences associated with of morphologically identified JTMD specie and congeners, as well as con-familial when morphological identification was limited to the family level.

Data Processing Description

BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date
- renamed parameters to BCO-DMO standard
- replaced blanks with underscores

[[table of contents](#) | [back to top](#)]

Related Publications

Carlton, J. T., Chapman, J. W., Geller, J. B., Miller, J. A., Carlton, D. A., McCuller, M. I., ... & Ruiz, G. M. (2017). Tsunami-driven rafting: Transoceanic species dispersal and implications for marine biogeography. *Science*, 357(6358), 1402-1406. <https://doi.org/10.1126/science.aao1498>
Results

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|------------------|---|----------|
| taxon | phylum and more specific taxonomic group | unitless |
| species | genus and species name | unitless |
| accession_number | Genbank record that matched the recovered sequeunce | unitless |
| pairwise_id | Pairwise Identity is the percent similarity between recovered sequence and the Genbank record | unitless |

[[table of contents](#) | [back to top](#)]

Instruments

| | |
|---|--|
| Dataset-specific Instrument Name | DNA sequencer |
| Generic Instrument Name | Automated DNA Sequencer |
| Dataset-specific Description | Ion Torrent platform |
| Generic Instrument Description | General term for a laboratory instrument used for deciphering the order of bases in a strand of DNA. Sanger sequencers detect fluorescence from different dyes that are used to identify the A, C, G, and T extension reactions. Contemporary or Pyrosequencer methods are based on detecting the activity of DNA polymerase (a DNA synthesizing enzyme) with another chemoluminescent enzyme. Essentially, the method allows sequencing of a single strand of DNA by synthesizing the complementary strand along it, one base pair at a time, and detecting which base was actually added at each step. |

| | |
|---|--|
| Dataset-specific Instrument Name | Homogenizer |
| Generic Instrument Name | Homogenizer |
| Dataset-specific Description | IKA Basic Mill |
| Generic Instrument Description | A homogenizer is a piece of laboratory equipment used for the homogenization of various types of material, such as tissue, plant, food, soil, and many others. |

| | |
|---|--|
| Dataset-specific Instrument Name | Thermal Cycler |
| Generic Instrument Name | Thermal Cycler |
| Generic Instrument Description | A thermal cycler or "thermocycler" is a general term for a type of laboratory apparatus, commonly used for performing polymerase chain reaction (PCR), that is capable of repeatedly altering and maintaining specific temperatures for defined periods of time. The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. They can also be used to facilitate other temperature-sensitive reactions, including restriction enzyme digestion or rapid diagnostics. (adapted from http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html) |

[[table of contents](#) | [back to top](#)]

Deployments

JTMD_2012

| | |
|--------------------|---|
| Website | https://www.bco-dmo.org/deployment/552342 |
| Platform | Carlton_shore |
| Start Date | 2012-12-01 |
| End Date | 2014-11-30 |
| Description | Japanese tsunami marine debris collection |

[[table of contents](#) | [back to top](#)]

Project Information

Testing the Invasion Process: Survival, Dispersal, Genetic Characterization and Attenuation of Marine Biota on the 2011 Japanese Tsunami Marine Debris Field. (JTMD-BF)

Coverage: North Pacific Ocean (W and E)

I. Biodiversity; Population and Food Web Analysis; Viability and Reproductive Condition; Dispersal Track and Growth History; Shellfish Pathogens/Parasites

This project seeks to document the biodiversity of Japanese species on arriving tsunami-generated debris, through morphological and genetic identification (including massively parallel DNA sequencing of whole community samples) and through quantitative replicate samples to determine numerical abundance, density, frequency, and biomass. In addition, species accumulation and rarefaction curves will be determined to estimate total inbound diversity.

Focuses include:

- Population structure of selected taxa, based on size/age class distributions.
- Viability and reproductive condition of selected taxa, based on fecundity, gonadal indices, and/or spore production, upon arrival.
- Food web analyses based upon tissue stable isotope ratios ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$).
- Dispersal track and growth history of selected taxa based on oxygen isotopic and elemental composition of shell calcite.
- Identity and prevalence of parasites and pathogens in oysters (*Crassostrea gigas*) and mussels (*Mytilus galloprovincialis*).

II. Biotic Attrition Over Time

Comparison of dead species assemblages on JTMD to live assemblages to assess the fate and alteration of debris communities over time.

III. Genetic Matching of Novel Invasions With JTMD Biota

Genetically characterize populations of target species so that if and when new invasions are detected, or when previously established invasions appear to be newly expanding or appearing in new locations, genetic studies can be undertaken to determine if these events are related to the JTMD phenomenon.

This is a Rapid Response Grant.

2020-09-30: Final data was not submitted for this project. The data for this research are available at the Dryad data depository (<http://dx.doi.org/10.5061/dryad.rh01m>). Contact Dr. Carlton for more information.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|-----------------------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1266234 |

[[table of contents](#) | [back to top](#)]