

Data on number of dives observed and total number of larval tracks across all turbulence regimes collected at Wood Hole Oceanographic Institution in 2011 (Larvae in turbulence project)

Website: <https://www.bco-dmo.org/dataset/561205>

Data Type: experimental

Version: 2015-06-08

Project

» [Larval Response to Turbulence During Dispersal and Settlement](#) (Larvae in turbulence)

Contributors	Affiliation	Role
Mullineaux, Lauren	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
Helfrich, Karl R.	Woods Hole Oceanographic Institution (WHOI)	Co-Principal Investigator
Wheeler, Jeanette	Woods Hole Oceanographic Institution (WHOI)	Student, Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Count of number of dives observed in each dataset of four trials (T1-T4) of increasing turbulence (s0-s7), corresponding to tank forcing frequencies of 0-1.75 Hz. Dives are identified as per the method described in Wheeler et al. (2013). PIV recordings were performed by J. Wheeler in Summer 2011 and 2012.

Relevant Reference:

Wheeler J.D., Helfrich K.R., Anderson E.J., McGann B., Staats P., Wargula A.E., Wilt K., Mullineaux L.S. (2013) Upward swimming of competent oyster larvae *Crassostrea virginica* persists in highly turbulent flow as detected by PIV flow subtraction. *Mar Ecol Prog Ser* 488, 171-185.

Wheeler J.D., Helfrich K.R., Anderson E.J., Mullineaux L.S. (2015) Isolating the hydrodynamic triggers of the dive response in eastern oyster larvae. *Limnol Oceanogr*, (doi:10.1002/lno.10098)

[[table of contents](#) | [back to top](#)]

Data Files

File
dive_v_turb.csv (Comma Separated Values (.csv), 4.97 KB) MD5:a8181da1d4f7d150db4f625f9ec9dc09
Primary data file for dataset ID 561205

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
exp_id	experiment identification	unitless
turb	turbulence machine dial setting	unitless
freq	oscillation frequency of double grid at amplitude 5 cm	Hz
trial	trial identification	unitless
dataset	id of 45 second video of larvae in tank	unitless
num_dives	number of dives observed in dataset	dive
num_tracks	number of tracks	tracks
prob_dive	Probability of dive (frequency of dives divided by number of larval trajectories) with respect to energy dissipation rate for each data set	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	Camera
Dataset-specific Description	high-speed monochrome camera (Photron Fastcam SA3) and a pulsed near-infrared laser (Oxford Lasers, Firefly 300 W, 1000 Hz, 808 nm)
Generic Instrument Description	All types of photographic equipment including stills, video, film and digital systems.

[[table of contents](#) | [back to top](#)]

Deployments

lab_Mullineaux_2011

Website	https://www.bco-dmo.org/deployment/561100
Platform	WHOI
Start Date	2011-06-01
End Date	2012-08-31
Description	Larval oysters in turbulence experiments

[[table of contents](#) | [back to top](#)]

Project Information

Larval Response to Turbulence During Dispersal and Settlement (Larvae in turbulence)

Coverage: Laboratory studies at Woods Hole Oceanographic Institution

Description from NSF award abstract:

The planktonic larval stage of benthic marine invertebrates provides a mechanism for exchange of individuals between remote populations. Dispersal is affected by swimming behaviors, particularly those that alter the larva's vertical position in the water. Larvae of some species change their vertical positions in response to turbulence by ceasing to swim and sinking downward (diving). By doing so, they can alter their horizontal transport in currents and increase their supply to the seafloor. The main objectives of this study are to investigate behavioral responses of oyster (*Crassostrea virginica*) larvae to turbulence in the water column and at the seafloor, and to determine how these behaviors affect settlement. The investigators hypothesize that diving behavior enhances settlement into suitable habitat, even where mean bed shear stress is high. They expect that once larvae approach the bottom, they can take advantage of temporal and spatial refuges (such as turbulent lulls in the lee of roughness elements) to settle in otherwise harsh conditions. Investigating larval responses to turbulence is a challenge because it requires simultaneous measurement of time-variant flows and larval behaviors. The investigators will modify a conventional particle image velocimetry (PIV) approach so it can be used to track larval motions and fluid velocities simultaneously. PIV provides information on flow kinematics (e.g., rotation and strain rate) in the immediate vicinity of a larva, as well as bulk dissipation rates and measures of Taylor and integral length scales that likely influence larval acceleration. When these measurements are coupled with a larval trajectory, they provide a history of the fluid environment a larva experiences, and can be used to determine what characteristic of turbulence triggers the diving behavior. They also make it possible to calculate the bottom shear stress an individual larva experiences when it encounters the bottom and attempts to settle. The investigators will examine turbulence effects on larval behaviors in the water column using a grid-stirred tank. They will use a racetrack flume to test the hypothesis that larval settlement success depends on the frequency of lulls of sufficient duration for larval attachment.

Laboratory experiments will provide a mechanistic understanding of larval behavior that can be used in general theoretical models exploring how behavior influences dispersal and population connectivity. The quantified swimming responses of oysters are critical input for coupled bio-physical models of dispersal in the field. An understanding of larval behavior contributes to our ability to predict the effects of natural and anthropogenic perturbations (some of which are linked to global climate change) on benthic communities in coastal ecosystems where turbulence and habitat suitability vary spatially. This information is critical for informed decision making on shellfish management and design of marine reserves. The technique developed for simultaneous PIV and larval tracking will open new questions in larval ecology and be broadly applicable to studies of plankton interactions with turbulence.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0850419

[[table of contents](#) | [back to top](#)]