

# Seawater concentrations of dissolved and total Th-232 and Th-230 collected from the R/V Atlantic Explorer (AE1410) cruise from Barbados to Bermuda during 2014

**Website:** <https://www.bco-dmo.org/dataset/639016>

**Data Type:** Cruise Results

**Version:** 2016-02-22

## Project

» [Chief Scientist Training Cruise \(Barbados-Bermuda, May 2014\)](#) (NSF Chief Sci Training Barbados-Bermuda)

Contributors	Affiliation	Role
<a href="#">Hayes, Christopher T.</a>	University of Southern Mississippi (USM)	Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Seawater concentrations of dissolved and total Th-232 and Th-230 were collected from the R/V Atlantic Explorer (AE1410) cruise from Barbados to Bermuda during 2014. This dataset was collected as part of a Chief Scientist Training Cruise, May-June 2014. The focus of this research was 'Tracking Saharan dust deposition in the Northwest Atlantic using Th isotopes'. This dataset is GEOTRACES-compliant.

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Dataset Description

This dataset was collected as part of a Chief Scientist Training Cruise, May-June 2014. The focus of this research was 'Tracking Saharan dust deposition in the Northwest Atlantic using Th isotopes'. This dataset is GEOTRACES-compliant.

## Methods & Sampling

Water sampling followed GEOTRACES protocols from a standard Niskin Bottle CTD/Rosette (Anderson et al. 2012). 4-5 L water samples were drained or filtered directly from Niskin bottles using Teflon-lined tygon tubing, acidified at sea with ultra-pure 6 molar hydrochloric acid, and analyzed at on-shore at MIT. Analysis involved Fe hydroxide co-precipitation, acid digestion, anion exchange chromatography and isotope dilution inductively-coupled plasma mass spectrometry. Laboratory methods were slightly modified from Anderson et al. (2012) as described by Hayes et al (2015).

## References:

Anderson, R. F., M. Q. Fleisher, L. F. Robinson, R. L. Edwards, J. Hoff, S. B. Moran, M. M. Rutgers van der Loeff, A. L. Thomas, M. Roy-Barman, and R. François (2012), GEOTRACES intercalibration of  $^{230}\text{Th}$ ,  $^{232}\text{Th}$ ,  $^{231}\text{Pa}$ , and prospects for  $^{10}\text{Be}$ , *Limnol. Oceanogr. Methods*, 10, 179-213.

Hayes, C. T., R. F. Anderson, S. L. Jaccard, R. François, M. Q. Fleisher, M. Soon, and R. Gersonde (2013), A new perspective on boundary scavenging in the North Pacific Ocean, *Earth Planet. Sci. Lett.*, 369-370(0), 86-

97, doi:<http://dx.doi.org/10.1016/j.epsl.2013.03.008>.

Owens, S. A., K. O. Buesseler, and K. W. W. Sims. "Re-evaluating the  $^{238}\text{U}$ -salinity relationship in seawater: implications for the  $^{238}\text{U}$ - $^{234}\text{Th}$  disequilibrium method." *Marine Chemistry* 127.1 (2011): 31-39.

## Data Processing Description

Th isotope concentrations are derived from measured Th-232/Th-229 and Th-230/Th-229 ratios as determined by mass spectrometry. The Th-229 spike at MIT was calibrated against a gravimetric Th standard as well as an artificial seawater standard called SWSTD 2010-1, described by Anderson et al. (2012). Corrections for ingrowth of Th-230 from dissolved U-234 during sample storage were made based on measured salinity and the U-salinity relationship of Owens et al. (2011).

The dissolved "excess" Th-230 (Th\_230\_D\_XS) is derived by subtracting from the measured dissolved Th-230 the component sourced from the partial dissolution of lithogenic material based on measured dissolved Th-232 and a lithogenic Th-230/232 atom ratio of  $4 \times 10^{-6}$ , as described by Hayes et al. (2013).

### BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date
- renamed parameters to BCO-DMO standard
- changed longitude from positive to negative values

Cruise AE1410 was supported by UNOLS and NSF. The sample analysis was supported by the W.O. Crosby post-doctoral fellowship from MIT to Christopher Hayes.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Th_AE1410.csv</b> (Comma Separated Values (.csv), 4.04 KB) MD5:6e2eb699b0386b96200041d8a9fcf6c9
Primary data file for dataset ID 639016

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
cruise_id	cruise identification	unitless
station	station number	unitless
sample	sample number derived from cast and bottle numbers	unitless
date_start	date at start of samping	yyyymmdd
lat_start	starting latitude; north is positive	decimal degrees
lon_start	starting longitude; east is positive	decimal degrees
press	pressure	decibars
depth	depth	meters
temp_ctd	temperature from CTD	degrees Celsius
sal_ctd	salinity from CTD	unitless
O2_ctd	dissolved oxygen from CTD	umol/kg
BAC	Beam Attenuation Coefficient from the transmissometer	per meter
fluor	fluorescence	arbitrary units
Th_230_D_CONC_BOTTLE	Dissolved (	uBq/kg
Th_230_D_CONC_BOTTLE_ERR	Dissolved (	uBq/kg
Th_230_D_CONC_BOTTLE_FLAG	quality flag: 1 = good; 2 = questionable; 3 = rejected	unitless
Th_230_TD_CONC_BOTTLE	Total dissolvable (from unfiltered water) Th-230 concentration	uBq/kg
Th_230_TD_CONC_BOTTLE_ERR	Total dissolvable (from unfiltered water) Th-230 concentration error	uBq/kg
Th_230_TD_CONC_BOTTLE_FLAG	quality flag: 1 = good; 2 = questionable; 3 = rejected	unitless
Th_232_D_CONC_BOTTLE	Dissolved ((	pmol/kg
Th_232_D_CONC_BOTTLE_ERR	Dissolved ((	pmol/kg
Th_232_D_CONC_BOTTLE_FLAG	quality flag: 1 = good; 2 = questionable; 3 = rejected	unitless
Th_232_TD_CONC_BOTTLE	Total dissolvable (from unfiltered water) Th-232 concentration	pmol/kg
Th_232_TD_CONC_BOTTLE_ERR	Total dissolvable (from unfiltered water) Th-232 concentration error	pmol/kg
Th_232_TD_CONC_BOTTLE_FLAG	quality flag: 1 = good; 2 = questionable; 3 = rejected	unitless
Th_230_D_XS_CONC_BOTTLE	Dissolved (	uBq/kg
Th_230_D_XS_CONC_BOTTLE_ERR	Dissolved (	uBq/kg
Th_230_D_XS_CONC_BOTTLE_ERR_FLAG	quality flag: 1 = good; 2 = questionable; 3 = rejected	unitless
date_U_separated	The date that uranium was chemically separated from thorium in the sample analysis; used in the ingrowth calculation	yyyymmdd

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Aerosol Sampler
<b>Dataset-specific Description</b>	Tisch Environmental PM10 aerosol sampler (TE-6070), using 20 cm x 25 cm Whatman41 filter sheets.
<b>Generic Instrument Description</b>	A device that collects a sample of aerosol (dry particles or liquid droplets) from the atmosphere.

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	GO-FLO Bottle
<b>Dataset-specific Description</b>	5-L Go-Flo bottle, attached to the ship's Kevlar line
<b>Generic Instrument Description</b>	GO-FLO bottle cast used to collect water samples for pigment, nutrient, plankton, etc. The GO-FLO sampling bottle is specially designed to avoid sample contamination at the surface, internal spring contamination, loss of sample on deck (internal seals), and exchange of water from different depths.

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Niskin bottle
<b>Dataset-specific Description</b>	ship's CTD with conventional Niskin bottles
<b>Generic Instrument Description</b>	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Trace element sampler
<b>Dataset-specific Description</b>	MITESS unit
<b>Generic Instrument Description</b>	Automated trace element sampler (MITESS or ATE unit). Bell, J., J. Betts, and E. Boyle (2002) MITESS: A Moored In-situ Trace Element Serial Sampler for Deep-Sea Moorings, Deep-Sea Research I: 49:2103-2118 (pdf) More description: <a href="http://boyle.mit.edu/~ed/MITESS/MITESShomepage.html">http://boyle.mit.edu/~ed/MITESS/MITESShomepage.html</a>

---

## Deployments

### AE1410

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/638976">https://www.bco-dmo.org/deployment/638976</a>
<b>Platform</b>	R/V Atlantic Explorer
<b>Report</b>	<a href="http://dmoserv3.bco-dmo.org/data_docs/GEOTRACES/AE1410/2014_ChfSci_Training_final_cruise_rpt.pdf">http://dmoserv3.bco-dmo.org/data_docs/GEOTRACES/AE1410/2014_ChfSci_Training_final_cruise_rpt.pdf</a>
<b>Start Date</b>	2014-05-30
<b>End Date</b>	2014-06-12
<b>Description</b>	UNOLS Chief Scientist training cruise. The original data from this cruise are available from the NSF R2R data catalog.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Chief Scientist Training Cruise (Barbados-Bermuda, May 2014) (NSF Chief Sci Training Barbados-Bermuda)

#### *NSF Award Abstract:*

##### Overview:

This grant will support a CST cruise on the R/V Atlantic Explorer from Barbados to Bermuda in May and June 2014. Chief Scientist Training (CST) cruises are part of an educational initiative designed for young US oceanographers, from senior graduate students to junior faculty, to increase their participation in seagoing research by educating them about the capabilities and opportunities provided by the UNOLS research fleet. In addition to providing sampling opportunities for individual research, a CST cruise is specifically organized to teach cruise participants how to successfully plan and execute a research cruise, from proposal to post-cruise reporting. Cruise participants will be selected based on the feasibility of their proposed Intellectual Merit: The ship track covers a oceanographically important transitional area in the western subtropical Atlantic, and thus the participants will be able to obtain scientifically valuable data in addition to gaining valuable skills and training that will help them to plan and execute successful cruises in their future careers.

##### Broader Impacts:

The CST cruise will help develop future expertise in the US oceanographic community by reducing the barriers to participation of young scientists in seagoing research, particularly those scientists from non ship-operator institutions. The activities proposed will help to disseminate information about the research opportunities that are provided by NSF and the UNOLS research fleet. The training and workshop activities are projected to increase the number and breadth of future proposals requesting ship support as well as to improve the quality of execution and success of funded cruises. The CST cruise opportunity will also help to broaden the geographic representation of the research community that is involved in ship-based scientific studies as well as increase the diversity of this community.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1360666</a>

[ [table of contents](#) | [back to top](#) ]