

# Food consumption and body measurements in *Panopeus herbstii* collected near North Inlet Estuary, Georgetown, SC during 2012 (Variation in Metabolic Processes project)

Website: <https://www.bco-dmo.org/dataset/648309>

Version:

Version Date: 2016-06-06

## Project

» [Linking Variation in Metabolic Processes as a Key to Prediction](#) (Variation in Metabolic Processes)

Contributors	Affiliation	Role
<a href="#">Griffen, Blaine D.</a>	University of South Carolina	Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Temporal Extent:** 2014-05-01

## Dataset Description

### Related Reference:

Griffen BD, Norelli AP (2015) Spatially variable habitat quality contributes to within-population variation in reproductive success. *Ecology and Evolution* 5:1474-1483

### Related Datasets:

[P. herbstii diet: sampling site characteristics \(Eco-Evo 2015\)](#)

[P. herbstii diet: body measurements \(Eco-Evo 2015\)](#)

## Methods & Sampling

We conducted a laboratory study in which we quantified consumption and reproductive effort. We collected 40 female mud crabs (*Panopeus herbstii*) in May from North Inlet. These were returned to the University of South Carolina in Columbia where they were placed into a recirculating aquarium held at 20°C on a 16 h:8 h light:dark cycle. Each crab was housed in an individual 1-L chamber, individually plumbed to ensure constant flow at 3 L/h. Crabs were fed twice per week (Monday and Thursday) and were given 48 h to consume their food before uneaten food was removed. The food was dried for 72 h at 70°C, and weighed to the nearest 0.01 mg. Crabs were fed one of 20 experimental diets that varied the total amount of food present (0.3, 0.6, 1.2, 3% of body weight at each feeding) and the proportion of that food that was animal tissue or algae (0.0:1.0, 0.25:0.75, 0.5:0.5, 0.75:0.25, 1.0:0.0 animal:algal). Two crabs were fed each of these 20 experimental diets. However, food consumed could not be directly controlled (food offered served as an upper limit to consumption, but

individual crabs could always choose to consume less on any given day). Each crab therefore had a different diet over the course of the experiment that reflected its aggregate daily diet decisions. We determined the average animal and algal consumption across feeding periods for each crab over the course of the experiment. Algal food was *Ulva lactuca*, animal food was tilapia filets. This experiment continued as described above for 10 weeks, after which the crabs were dissected and ovaries, hepatopancreas, and the rest of the body were dried separately at 70°C for 72 h and were then weighed to the nearest 0.01 mg.

## Data Processing Description

Raw data are presented.

### BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date, reference information
- renamed parameters to BCO-DMO standard

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Pherb_diet_repro.csv</b> (Comma Separated Values (.csv), 2.92 KB) MD5:0aef36a6fb664d904791098b905dcc8c
Primary data file for dataset ID 648309

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
crab	unique identification number for each crab	unitless
algae_offered	average wet mass of algae <i>Ulva lactuca</i> offered at each feeding	grams
fish_offered	average wet mass of tilapia filet offered at each feeding	grams
algae_consumed	average dry mass of algae consumed at each feeding	grams
fish_consumed	average dry mass of fish consumed at each feeding	grams
absorption	average proportion of consumed food that was absorbed at each feeding (calculated from consumed - feces produced at each feeding)	grams
carap_width	carapace width of crab	millimeters
body_wgt_tot	dry mass of crab	grams
hepato_wgt	dry mass of hepatopancreas	grams
ovary_wgt	dry mass of ovary	grams
egg_wgt	dry mass of egg clutches that were produced during the experiment	grams
vitellogenic	whether the crab was vitellogenic: Y=1; N=0	unitless
molted	whether crab molted during the experiment: Y=1; N=0	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	scale
<b>Generic Instrument Description</b>	An instrument used to measure weight or mass.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### Griffen\_lab

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/638572">https://www.bco-dmo.org/deployment/638572</a>
<b>Platform</b>	Univ_S_Carolina
<b>Start Date</b>	2012-01-01
<b>End Date</b>	2016-12-31

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Linking Variation in Metabolic Processes as a Key to Prediction (Variation in Metabolic Processes)

#### *Description from NSF award abstract:*

A major goal of biological and ecological sciences is to understand natural systems well enough to predict how species and populations will respond to a rapidly changing world (i.e., climate change, habitat loss, etc.). A population under any conditions will grow, shrink, or disappear altogether depending on how efficiently individuals consume resources (food), utilize that food metabolically, and eventually reproduce. However, making accurate predictions based on these metabolic processes is complicated by the realities that each species has different resource requirements and that no two individuals within a species are exactly alike. Rather, individuals vary and this variation, both within and across species, is central to many ecological and evolutionary processes. Developing the ability to predict responses of biological systems to a changing world therefore requires a mechanistic understanding of variation. The goal of this project is to improve this mechanistic understanding by examining variation within a metabolic context across a range of species that have a spectrum of commonly-seen resource requirements. Further, the work capitalizes on a unique biological characteristic of this group of species that allows control and manipulation of individual reproduction, facilitating experimental study of the mechanistic links between variation in individual consumption, metabolism, and reproduction. The foundation this research is a combination of field measurements and laboratory experiments using both well-established and newly-developed techniques to quantify these links. The result will be a quantitative framework to predict how individuals will respond reproductively to changes in resource use. Because of the close link between individual reproduction and population dynamics, this research will contribute substantially to predictions in population dynamics under realistic conditions where individuals use more than a single resource, and improve the prediction of responses to current and future ecological changes.

#### **The following publications and data resulted from this project:**

Belgrad, B. and B. Griffen. 2016. Predator-prey interactions mediated by prey personality and predator identity. *Proc. Roy. Soc. B*: In Review. [2016-01-20]

[P. herbstii mortality data](#): Mortality of crabs when exposed to either a single blue crab, toadfish, or no predator for a week

[P. herbstii personality data](#): Refuge use of crabs when exposed to predator odor cues from either blue crabs, toadfish, or control of no cue

[P. herbstii predator behavior data](#): Refuge use and mobility of blue crabs and toadfish while in mesocosms for a week - behavior measured during two days.

Belgrad, B. and B. Griffen. 2016. The influence of dietary shifts on fitness of the blue crab, *Callinectes sapidus*. *PLoS One*. DOI: [10.1371/journal.pone.0145481](https://doi.org/10.1371/journal.pone.0145481).

[Blue crab activity](#): Activity of crabs fed different diets over a summer

[Blue crab egg size](#): Volume of eggs for crabs fed different diets

[Blue crab hepatopancreas index \(HSI\)](#): Weight of hepatopancreas for crabs fed different diets

[Blue crab hepatopancreas lipid content](#): Hepatopancreas lipid content of crabs fed different diets

[Blue crab reproductive tissue analysis \(GSI\)](#): Gonadosomatic index of blue crabs on various diets

[Blue crab survival](#): Blue crab survival data during the dietary study

Knotts ER, Griffen BD. 2016. Individual movement rates are sufficient to determine and maintain dynamic spatial positioning within *Uca pugilator* herds. *Behavioral Ecology and Sociobiology* 70:639-646

[Uca pugilator: behavior change with carapace marking](#): Search space behavior due to carapace treatment (control, nail polish, and food dye)

[Uca pugilator: field spatial position](#): Assessment of individual's position within a herd at 3 min. intervals; for proportion of time found at edge of herd

[Uca pugilator: herd position proportion](#): Individual's proportion of time spent in an edge/alone position among a herd

[Uca pugilator: search space distribution](#): Search space that crabs traveled; to evaluate the sample's distribution of exploratory behavior

Belgrad, B. and B. Griffen. 2015. Rhizocephalan infection modifies host food consumption by reducing host activity levels. *Journal of Experimental Marine Biology and Ecology*. 466: 70-75.

[E. depressus digestion time](#) : Time taken for food to pass through gut of flat-backed mud crabs infected by a parasite

[E. depressus metabolism](#): Respiration rate of infected/uninfected flat-backed mud crabs

[E. depressus reaction time to prey](#): Time taken for infected/uninfected flat-backed mud crabs to react to the presence of prey

Blakeslee, A.M., C.L. Keogh, A.E. Fowler, B. Griffen. 2015. Assessing the effects of trematode infection on invasive green crabs in eastern North America. *PLOS One* 10(6): e0128674.(pdf)

[Carcinus: hemocyte density](#): Counts of circulating hemocyte density in *Carcinus maenas*

[Carcinus: parasites physiology behavior](#): Behavior and physiology of *Carcinus maenas* infected with trematode parasite

Griffen BD, Norelli AP (2015) Spatially variable habitat quality contributes to within-population variation in reproductive success. *Ecology and Evolution* 5:1474-1483.

[P. herbstii diet: sampling site characteristics \(Eco-Evo 2015\)](#)

[P. herbstii diet: body measurements \(Eco-Evo 2015\)](#)

[P. herbstii diet & reproduction \(Eco-Evo 2015\)](#)

[P. herbstii: collection sites \(Eco-Evo 2015\)](#)

Griffen BD, Riley ME (2015) Potential impacts of invasive crabs on one life history strategy of native rock crabs in the Gulf of Maine. *Biological Invasions* 17:2533-2544.

[Cancer consumption and reproduction \(Bio.Inv. 2015\)](#): Lab experiment linking dietary consumption and reproduction

Griffen BD, Vogel M, Goulding L, Hartman R (2015) Energetic effects of diet choice by invasive Asian shore crabs: implications for persistence when prey are scarce. *Marine Ecology Progress Series* 522:181-192.

[Hemigrapsus diet 1 \(MEPS 2015\)](#)

[Hemigrapsus diet 2 \(MEPS 2015\)](#)

Hogan and Griffen (2014). The Dietary And Reproductive Consequences Of Fishery-Related Claw Removal For The Stone Crab *Menippe* Spp. *Journal of Shellfish Research*, Vol. 33, No. 3, 795-804.

[Stone crab: 052012-DietChoiceExp1](#): Prey choice for 2-clawed and 1-clawed Stone Crabs (*Menippe* spp.)

[Stone crab: 052012-LongTermConsumption](#): Long-term consumption for 2-clawed and 1-clawed Stone Crabs (*Menippe* spp.), summer of 2012

[Stone crab: 062013-DietChoiceExp2](#): Prey choice for 2-clawed and 1-clawed Stone Crabs (*Menippe* spp.)

[Stone crab: 062013-PreySizeSelection](#): Prey Size selection ranking for 2-clawed and 1-clawed Stone Crabs (*Menippe* spp.)

Riley M, Johnston CA, Feller IC, and Griffen B. 2014. Range expansion of *Aratus pisonii* (mangrove tree crab) into novel vegetative habitats. *Southeastern Naturalist* 13(4): 43-38

[A. pisonii: range expansion](#): *Aratus pisonii* survey in native mangrove and novel salt marsh habitats

Riley M, Vogel M, Griffen B. 2014. Fitness-associated consequences of an omnivorous diet for the mangrove tree crab *Aratus pisonii*. *Aquatic Biology* 20:35-43, DOI: 10.3354/ab00543

[A. pisonii: fitness and diet](#): Impact of diet variation on physiological and reproductive condition of *A. pisonii*

Toscano BJ, Newsome B, Griffen BD (2014) Parasite modification of predator functional response. *Oecologia* 175:345-352b

[E. depressus - parasite and feeding \(Oecologia, 2014\)](#): Feeding with and without parasitic barnacle infection

[E. depressus - parasite and prey handling \(Oecologia, 2014\)](#): Food handling with and without parasitic barnacle infection

[E. depressus - parasite study - field survey \(Oecologia, 2014\)](#): Parasitised field survey

Toscano BJ, Griffen BD (2014) Trait-mediated functional responses: predator behavioural type mediates prey consumption. *Journal of Animal Ecology* 83:1469-1477

[P. herbstii - activity and feeding \(JAE, 2014\)](#): Activity level and feeding with and without predator cue

Toscano BJ, Gatto J, Griffen BD (2014) Effects of predation threat on repeatability of individual crab behavior revealed by mark recapture. *Behavioral Ecology and Sociobiology* 68:519-527

[P. herbstii - recapture behavior \(BESB, 2014\)](#): Mud crabs refuge use and activity level - initial measurements

[P. herbstii - refuge use \(BESB, 2014\)](#): Effect of predation threat on repeatability of individual crab behavior revealed by mark-recapture

Griffen BD, Altman I, Bess BM, Hurley J, Penfield A (2012) The role of foraging in the success of invasive species. *Biological Invasions*. 14:2545-2558

[Hemigrapsus seasonal diet \(Bio.Inv. 2012\)](#): Percent herbivory and gut fullness for *Hemigrapsus sanguineus* at different times of year

Griffen BD, Toscano B, Gatto J (2012) The role of intraspecific trait variation in mediating indirect interactions. *Ecology* 93:1935-1943

[P. herbstii refuge use \(Ecology, 2012\)](#): Proportion of time that *Panopeus herbstii* spent using refuge habitats in a lab experiment

[P. herbstii: Field personality distribution \(Ecology, 2012\)](#): Field distribution of personality types in the mud crab *Panopeus herbstii* relative to tidal height

[P. herbstii: Trait mediated indirect effect \(Ecology, 2012\)](#): Influence of refuge use by the mud crab *Panopeus herbstii* on consumption of bivalves

Riley ME, Griffen BD (2017) Habitat-specific differences alter traditional biogeographic patterns of life history in a climate-change induced range expansion. *PLOS One* 12(5):e0176263

[A. pisonii: egg size](#): Comparing egg size in *Aratus pisonii* populations from mangrove and salt marsh habitats

[A. pisonii: fecundity](#): Determining fecundity of *Aratus pisonii* populations in mangrove and salt marsh habitats

[A. pisonii: larval starvation resistance](#): Comparing larval quality in *Aratus pisonii* populations from mangrove and salt marsh habitats

[A. pisonii: latitudinal body size](#): Survey examining latitudinal body size patterns in *Aratus pisonii*

[A. pisonii: predation](#): Comparing predation pressure on *Aratus pisonii* in mangrove and salt marsh habitats

[A. pisonii: reproductive effort](#): Survey comparing *Aratus pisonii* reproductive effort in native and novel habitats

[A. pisonii: herbivory](#): Relationship between leaf herbivory, tree characteristics, and refuge availability

[A. pisonii: mangrove tree survey](#): Mangrove tree distribution and characteristics in a dwarf mangrove system

Cannizzo ZJ, Dixon SR & Griffen BD (2018). An anthropogenic habitat within a suboptimal colonized ecosystem provides improved conditions for a range-shifting species. *Ecology and Evolution*, 8(3):1524-1533.

[A. pisonii: behavior](#): Proportion of time the mangrove tree crab *Aratus pisonii* spent in different behaviors related to diet and energy storage

[A. pisonii: dock-marsh thermal](#): Thermal readings from under a dock and in a nearby salt marsh

[A. pisonii: sun-shade](#): Proportion of time that mangrove tree crab *Aratus pisonii* spent in sun and shade in three habitats, 2015-2016.

[A. pisonii: thermal picture](#): Thermal condition of *A. pisonii* in three habitats: under dock, mangroves, saltmarsh

[ [table of contents](#) | [back to top](#) ]

---

## Funding

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1129166</a>

[ [table of contents](#) | [back to top](#) ]