

Velocity measurements from the turbulence generator taken with an Acoustic Doppler Velocimeter on the Friday Harbor dock and laboratory in 2012 (Jellyfish predation in turbulence project)

Website: <https://www.bco-dmo.org/dataset/649971>

Data Type: Other Field Results

Version:

Version Date: 2016-07-06

Project

» [Influence of organism-scale turbulence on the predatory impacts of a suite of cnidarian medusae](#) (jellyfish predation in turbulence)

Contributors	Affiliation	Role
Sutherland, Kelly Rakow	University of Oregon (OIMB)	Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	Guest, BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Related Dataset:

[ADV_field](#)

Methods & Sampling

Current velocity measurements were taken from the collection site at the surface using a Vectrino Acoustic Doppler Velocimeter (ADV; Nortek, Oslo, Norway), at depth between 0.5 and 5 m using in situ PIV (Katija and Dabiri, 2008). The ADV measurements were collected at a sampling rate of 100 Hz over 1-5 min intervals.

Data Processing Description

The TKE dissipation rate at the surface was calculated from the ADV-produced velocities in the horizontal, x , direction, u , following (Sanford, 1997) $\text{Epsilon(ADV)} = (u_{\text{rms}})^3 / Z \kappa$ where Z is the total water column depth (14 m) and κ is von Karman's constant (0.4).

BCO-DMO Processing:

- added conventional header with dataset name, PI name, version date
- renamed parameters to BCO-DMO and BODC standards

- combined tank and field tables
- added site column
- reduced digits to right of decimal from 13 to 2
- changed 'betwn 2 and 18 ft. (0.6 and 5 m)' to '0.6-5m'

[[table of contents](#) | [back to top](#)]

Data Files

File
tank_field_turb.csv (Comma Separated Values (.csv), 530 bytes) MD5:d51e15cec0cf9213c4471c438af83b80
Primary data file for dataset ID 649971

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
site	experimental setting: tank or field	unitless
location	location within tank or depth off dock	unitless
U_rms	root mean square of velocity	cm/sec
dissip_rate	dissipation rate of turbulence kinetic energy	m ² /s ³

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	
Generic Instrument Name	Acoustic Doppler Velocimeter
Dataset-specific Description	ADV; Nortek, Oslo, Norway
Generic Instrument Description	ADV is the acronym for acoustic doppler velocimeter. The ADV is a remote-sensing, three-dimensional velocity sensor. Its operation is based on the Doppler shift effect. The sensor can be deployed either as a moored instrument or attached to a still structure near the seabed. Reference: G. Voulgaris and J. H. Trowbridge, 1998. Evaluation of the Acoustic Doppler Velocimeter (ADV) for Turbulence Measurements. J. Atmos. Oceanic Technol., 15, 272-289. doi: http://dx.doi.org/10.1175/1520-0426(1998)0152.0.CO;2

[[table of contents](#) | [back to top](#)]

Deployments

FHL_Sutherland

Website	https://www.bco-dmo.org/deployment/649916
Platform	Friday_Harbor
Start Date	2012-06-01
End Date	2016-06-30

[[table of contents](#) | [back to top](#)]

Project Information

Influence of organism-scale turbulence on the predatory impacts of a suite of cnidarian medusae (jellyfish predation in turbulence)

Coverage: Friday Harbor Labs, WA

Bloom-forming jellyfish are increasing in number, frequency and magnitude, in part due to anthropogenic impacts, underscoring a need for enhanced understanding of trophic exchanges in jellyfish-dominated ecosystems. Interactions between jellyfish and their prey are driven by morphology, behavior, and unique fluid signatures that result in species-specific prey selection patterns. Fluid signatures generated by predators entrain prey, and motile prey organisms have evolved to sense and respond to these stereotyped fluid signatures. The shape and coherence of these unique fluid signatures are strongly mediated by turbulence, which is ubiquitous in the ocean. Yet, the effects of turbulence are almost always neglected in feeding studies. This three-year project will investigate the influence of turbulence on predator-prey interactions using a suite of cnidarian hydromedusae with unique morphologies, fluid signatures and prey selection patterns collected in the region of Friday Harbor Laboratory, WA.

This project seeks to establish a detailed, mechanistic understanding of the effects of turbulence on organism-scale predator-prey interactions using gelatinous zooplankton predators with contrasting predation modes. The PI will investigate prey selection under varying levels of turbulence by studying swimming behavior, wake structure, and predator-prey interactions in a laboratory turbulence generator designed for fragile plankton. The PI will also make in situ measurements of turbulence and observations of organism behavior using a Self-contained Underwater Velocimetry Apparatus (SCUVA). This is a fully submersible instrument for flow visualization, and its use will provide a cross-calibration of field and laboratory rates and behaviors. The influence of turbulence on trophic position among the different species of hydromedusae will be quantified through field studies of prey selection patterns. The proposed comparative approach using species with distinct predation modes will provide insights applicable to other planktonic predators that can be similarly grouped.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1155084

[[table of contents](#) | [back to top](#)]