

Isolation, culturing, and sequencing of bacteria and viruses collected in Canoe Cove, Nahant, MA during 2010 (Marine Bacterial Viruses project)

Website: <https://www.bco-dmo.org/dataset/658497>

Data Type: experimental

Version: 1

Version Date: 2016-09-09

Project

» [How can bacterial viruses succeed in the marine environment?](#) (Marine Bacterial Viruses)

| Contributors | Affiliation | Role |
|--------------------------------|---|---------------------------|
| Kelly, Libusha | Yeshiva University | Principal Investigator |
| Polz, Martin | Massachusetts Institute of Technology (MIT) | Co-Principal Investigator |
| Ake, Hannah | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

Isolation, culturing, and sequencing of bacteria and viruses collected in Canoe Cove, Nahant, MA during 2010 (Marine Bacterial Viruses project)

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: Lat:42.419 Lon:-70.906

Temporal Extent: 2010-08-10 - 2010-10-13

Dataset Description

This data contains isolation, culturing, and sequencing of bacteria and viruses from the Nahant Vibrio and Phage Genome Collection. Vibrio and associated phage genomes isolated off the coast of Nahant, MA, USA.

Related Datasets:

NCBI BioSample accessions for viruses and microbes: <http://www.bco-dmo.org/dataset/658586>

Methods & Sampling

Bacteria and viruses were collected from the littoral marine zone at Canoe Cove, Nahant, MA, USA, on August 22 [ordinal day 222], September 18 [261], and October 13 [286], 2010.

Bacteria were collected using a previously described size-fractionation method[1]. Bacterial strain naming convention is described using the example of 10N.286.54.E5: the first position (here "10N") indicates the year

(2010) and location (Nahant) of isolation, the second position (here “286”) indicates the ordinal day of isolation, the third position (here “54”) is a code representing the size-fraction of origin (0.2um: 45,46,47; 1um: 48,49,50; 5um: 51,52,53; 63um: 54,55,56), and the fourth position is the storage plate well identifier. Multiple codes within the size-fraction identifier reflect independent water samples for the 63um fraction, and independent water sample fractionation series for the other size classes (water sample A: 45,51,54; sample B: 46, 52, 55; sample C: 47, 53, 56).

Bacterial genome libraries were prepared for sequencing using a tagmentation-based approach and 1-2ng input DNA per isolate, as previously described[2]. Genomes were sequenced in multiplexed pools of 50-60 samples per Illumina HiSeq lane. Accession numbers for all bacterial genomes associated with this study are provided in Supplementary Table S1.

Bacterial phylogenetic relationships were determined by extracting ribosomal proteins from 278 genomes with hmmsearch[3] and aligning with MAFFT[4] as described in Hehemann et al. (2016)[5]. These strains were added to the Vibrionaceae ribosomal phylogeny used in Hehemann et al., 2016 and taxonomy was assigned using manual inspection. Full-length *hsp60* sequences were also extracted from these genomes using hmmsearch with default parameters and the Cpn60 hmm (PF00118) from Pfam[6]. The *hsp60* sequences were aligned using the mafft-fftinsi algorithm. Sanger-sequenced *hsp60* fragments from 40 strains lacking genome sequences were added to this alignment using the mafft-fftinsi algorithm with the --addfragments option. The *hsp60* alignment was concatenated to the ribosomal protein alignment and used to create a phylogeny using RAxML under a partitioned general time reversible (GTR) model (options: -q, -m GTRGAMMAX [7]). SH-like supports were calculated using RAxML.

Viruses were collected using a previously described iron flocculation approach[8], using 4L sample volumes, 0.2um pre-filtration to remove bacteria, 0.2um filters for floc-capture, and oxalate solution for resuspension to maintain virus viability. Isolation of viruses was performed as follows. Iron-oxalate concentrate volumes equivalent to 15mL of seawater were mixed into agar overlays of 1,334 potential host *Vibrio*. The agar overlays were performed by combining 150uL overnight host culture and virus concentrate directly on a bottom agar (1% agar, 5% glycerol, 125mL/L of chitin supplement [40g/L coarsely ground chitin, autoclaved, 0.2um filtered] in 2216MB), directly pipetting 2mL of molten top agar (52 degrees C, 0.4% agar, 5% glycerol, in 2216MB) onto the bottom agar, and rapidly swirling to mix. Plates were incubated for 2 weeks, and plaques were archived for later purification. Sequencing and genome analysis of viruses is described briefly, as follows. High titer lysates of serially purified viruses were concentrated using 30kD centrifugal filter units (Millipore, Amicon Ultra Centrifugal Filters, Ultracel 30K, UFC903024) and washed with 1:100 Marine Broth 2216 to reduce salts for nuclease treatment. Concentrates were brought to approximately 500uL using 1:100 diluted 2216MB and then treated with DNase I and RNase A for 65 min at 37 degrees C to digest unencapsidated nucleic acids. Nuclease treated viral lysates were extracted by addition of 1:10 final volume of SDS mix (0.25M EDTA, 0.5M Tris-HCl (pH9.0), 2.5% sodium dodecyl sulfate), 30 min incubation at 65C; addition of 0.125 volumes 8M potassium acetate, 60 min incubation on ice; addition of 0.5 volumes phenol-chloroform; recovery of nucleic acids from aqueous phase by isopropanol and ethanol precipitation. Genomes were sequenced in multiplexed pools using Illumina MiSeq and HiSeq technologies, assembled using CLC assembly cell, and manually curated to standardize genome start positions for the Caudovirales.

Viral strain naming convention is described using the example of 1.008.O_10N.286.54.E5, with specific identifiers separated by a period. The first position (here “1”) represents a unique identifier for each independent plaque isolated from a given host from the initial exposure of a given host to an environmental virus concentrate. The second position (here “008”) represents a unique working ID for a host strain. The third position (here “O”) indicates a unique sublineage generated from a single plaque during viral serial purification, for example due to the emergence of multiple plaque morphologies. Following the underscore is the full strain ID of the host of isolation, as described above.

Accession numbers for all viral genomes associated with this study are included under NCBI BioProject PRJNA328102.

Data Processing Description

These are sequenced, assembled, annotated genomes for 282 viruses and 301 *Vibrio* strains.

Data Management Office Notes:

- Data was initially separated into two files for microbes and viruses. These files were combined.
- All columns which contained no data for either viruses or microbes were removed.

- "nd" was entered into columns with no recorded data.
- Column names were reformatted to comply with BCO-DMO naming standards.

[[table of contents](#) | [back to top](#)]

Data Files

| File |
|---|
| biosample_submissions.csv (Comma Separated Values (.csv), 153.71 KB) MD5:a2f8dae1ee7439697c8a9b33bbcad1f5 |
| Primary data file for dataset ID 658497 |

[[table of contents](#) | [back to top](#)]

Related Publications

Baym, M., Kryazhimskiy, S., Lieberman, T. D., Chung, H., Desai, M. M., & Kishony, R. (2015). Inexpensive Multiplexed Library Preparation for Megabase-Sized Genomes. *PLOS ONE*, 10(5), e0128036.

doi:[10.1371/journal.pone.0128036](https://doi.org/10.1371/journal.pone.0128036)

Methods

Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLoS Computational Biology*, 7(10), e1002195.

doi:[10.1371/journal.pcbi.1002195](https://doi.org/10.1371/journal.pcbi.1002195)

Methods

Finn, R. D., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Mistry, J., Mitchell, A. L., ... Bateman, A. (2015). The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Research*, 44(D1), D279–D285.

doi:[10.1093/nar/gkv1344](https://doi.org/10.1093/nar/gkv1344)

Methods

Hehemann, J.-H., Arevalo, P., Datta, M. S., Yu, X., Corzett, C. H., Henschel, A., ... Polz, M. F. (2016). Adaptive radiation by waves of gene transfer leads to fine-scale resource partitioning in marine microbes. *Nature Communications*, 7(1). doi:[10.1038/ncomms12860](https://doi.org/10.1038/ncomms12860)

Methods

Hunt, D. E., David, L. A., Gevers, D., Preheim, S. P., Alm, E. J., & Polz, M. F. (2008). Resource Partitioning and Sympatric Differentiation Among Closely Related Bacterioplankton. *Science*, 320(5879), 1081–1085.

doi:[10.1126/science.1157890](https://doi.org/10.1126/science.1157890)

Methods

John, S. G., Mendez, C. B., Deng, L., Poulos, B., Kauffman, A. K. M., Kern, S., ... Sullivan, M. B. (2010). A simple and efficient method for concentration of ocean viruses by chemical flocculation. *Environmental Microbiology Reports*, 3(2), 195–202. doi:[10.1111/j.1758-2229.2010.00208.x](https://doi.org/10.1111/j.1758-2229.2010.00208.x)

Methods

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30(4), 772–780. doi:[10.1093/molbev/mst010](https://doi.org/10.1093/molbev/mst010)

Methods

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30(9), 1312–1313. doi:[10.1093/bioinformatics/btu033](https://doi.org/10.1093/bioinformatics/btu033)

Methods

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|--------------------------|--|-----------------|
| organism_type | Type of organism described | unitless |
| sample_name | Sample name in source database | unitless |
| bioproject_accession | The accession number of the BioProject(s) to which the BioSample belongs | unitless |
| organism | Organism associated with sample. Identified to species when possible. | unitless |
| strain | Microbial or eukaryotic strain name | unitless |
| isolate | Identification or description of the specific individual from which this sample was obtained | unitless |
| host | The natural (as opposed to laboratory) host to the organism from which the sample was obtained. | unitless |
| lab_host | Scientific name and description of the laboratory host used to propagate the source organism or material from which the sample was obtained. | unitless |
| isolation_source | Describes the physical, environmental and/or local geographical source of the biological sample from which the sample was derived. | unitless |
| collection_date | Date of sampling; mm/dd/yy | unitless |
| geo_loc_name | Geographical origin of the sample. | unitless |
| sample_type | Sample type, such as cell culture, mixed culture, tissue sample, whole organism, single cell, metagenomic assembly | unitless |
| env_biome | Descriptor of the broad ecological context of a sample. | unitless |
| lat | Latitude | decimal degrees |
| lon | Longitude | decimal degrees |
| temp | Temperature of the sample at time of sampling. | degrees celsius |
| ordinal_day_of_isolation | Day of year sample was isolated. | unitless |
| description | Description of the sample. | unitless |

[[table of contents](#) | [back to top](#)]

Deployments

CanoeCove 2010

| | |
|-------------------|---|
| Website | https://www.bco-dmo.org/deployment/658581 |
| Platform | shoreside Massachusetts |
| Start Date | 2010-08-22 |
| End Date | 2010-10-13 |

[[table of contents](#) | [back to top](#)]

Project Information

How can bacterial viruses succeed in the marine environment? (Marine Bacterial Viruses)

Coverage: Coastal waters off Nahant, MA

Description from NSF award abstract:

Microbes make up the majority of the biomass in the ocean and viral mortality is one of the main ecological factors determining the diversity, abundance and turnover of microbial taxa. Yet, in spite of the known overall importance of viruses, the dynamics of their interactions with their specific microbial hosts remain poorly understood. This project will characterize viral strategies for survival and interaction with their hosts in the ocean, with the goal of enabling a better understanding of the conditions under which viruses can effectively control bacterial populations. The work will generate and integrate diverse data types, ranging from quantification of specific interactions, environmental dynamics of microbial hosts and their viruses, and comparative genome analysis. While the project focuses on the coastal ocean of New England, the approaches and findings will be applicable to the larger field of marine microbial ecology, to other virus/host systems in nature and to engineered systems. This project will fill a gap in current microbial ecology curricula by creating a bioinformatics module to provide training in large-scale sequence data collection and analysis. The module will be refined through testing during an annual course in Nicaragua and will be broadly accessible in the US and internationally. The close collaboration, throughout this project and its associated outreach, between two laboratories with complementary research strengths will provide highly interdisciplinary training for undergraduate students as well for two graduate students.

Viruses and their microbial hosts have co-evolved over billions of year and shape the ecology of the ocean in many ways. Broadly, understanding the mechanisms and emergent properties of virus-host interactions will allow for better understanding and modeling of biogeochemical cycles and the diversity of microbes at the population and genomic level. The guiding hypothesis of this project is that the prevalence of each of different viral strategies is probabilistic and linked to host availability, environmental parameters, and frequency-dependent competition with other virus strains for available hosts. This research will address four aims that characterize how viruses interact with their hosts in the dilute ocean environment by:

- (1) quantification of ecological tradeoffs between specialist and generalist viral strategies,
- (2) estimation of the prevalence of dual lytic/lysogenic strategy in marine viruses,
- (3) identification of host surface receptors of particular viruses and examination of genetic signatures of distinct viral strategies in virus and host genomes, and
- (4) identification of genetic and metabolic interactions between virus and host genomes.

This study takes advantage of a model system with the largest available collection of viruses and hosts for which host range and genome sequences have been determined. This work will provide fine-scale analysis of host and phage genomic diversity and abundance in this model system, while at the same time estimating host-range and co-infection, all of which represent important, poorly constrained parameters in virus-host interactions. Finally, this project complements the large number of studies that have looked at single host-virus interactions, metagenome sequencing, and assessment of viral impact on microbial production.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|-----------------------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1435868 |

[[table of contents](#) | [back to top](#)]