

# Grazing experiment 4: Cell size measurements of low-high pCO<sub>2</sub> acclimated Rhodomonas (E Hux Response to pCO<sub>2</sub> project)

**Website:** <https://www.bco-dmo.org/dataset/668794>

**Data Type:** experimental

**Version:**

**Version Date:** 2016-12-06

## Project

» [Planktonic interactions in a changing ocean: Biological responses of \*Emiliana huxleyi\* to elevated pCO<sub>2</sub> and their effects on microzooplankton](#) (E Hux Response to pCO<sub>2</sub>)

Contributors	Affiliation	Role
<a href="#">Olson, M Brady</a>	Western Washington University (WWU)	Principal Investigator
<a href="#">Love, Brooke</a>	Western Washington University (WWU)	Co-Principal Investigator
<a href="#">Strom, Suzanne</a>	Western Washington University (WWU)	Co-Principal Investigator
<a href="#">Still, Kelly Ann</a>	Western Washington University (WWU)	Student
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Dataset Description

### Related Reference:

Still, Kelly Ann, Microzooplankton grazing, growth and gross growth efficiency are affected by pCO<sub>2</sub> induced changes in phytoplankton biology. (Masters Thesis) Western Washington University. <http://cedar.wvu.edu/cgi/viewcontent.cgi?article=1490&context=wwuet>

### Methods & Sampling

The phytoplankton *Rhodomonas* sp. CCMP 755 was grown semi-continuously in atmosphere controlled chambers at three different CO<sub>2</sub> treatment concentrations; Ambient (400ppmv), Moderate (750ppmv), and High (1000ppmv). Cultures were diluted daily starting day 4 with pre-equilibrated media containing f/50 nutrients. On day 10, after *Rhodomonas* cells from the treatments were mounted live on a microscope slide and 50 cells from each treatment replicate were imaged using RSImage software under 400X magnification on an Olympus CHA microscope. ImageJ software was used to measure *Rhodomonas* length and width. *Rhodomonas* cells are described as having a prolate spheroid shape. The volume was calculated using:  $V_{prolate}(\mu m^3) = (4/3) \pi a^2 b$ . Where  $a=1/2$  width and  $b=1/2$  length of the *Rhodomonas* cell

### Data Processing Description

These data are unprocessed cell sizes as calculated above.

### BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- nd (no data) was entered into all blank cells
- added column for treatment to allow leveling of data

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>expt4_cell_size_day10.csv</b> (Comma Separated Values (.csv), 20.80 KB) MD5:c10b1047172cdb036fe4eab2330a4b5a
Primary data file for dataset ID 668794

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
treatment	treatment identifier: ambient, moderate, or high	unitless
Exp_day_treatment_rep_cell_number	sample identifier: individual cell measured: experiment day_pCO2 level_replicate_cell number	unitless
length	cell length	micrometers
width	cell width	micrometers
volume	cell volume	micrometers

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Olympus CHA microscope
<b>Generic Instrument Name</b>	Microscope - Optical
<b>Dataset-specific Description</b>	Used to measure Rhodomonas cells
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

## Lab\_Olson\_B

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/521277">https://www.bco-dmo.org/deployment/521277</a>
<b>Platform</b>	WWU
<b>Start Date</b>	2011-03-31
<b>End Date</b>	2016-09-15
<b>Description</b>	laboratory experiments

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Planktonic interactions in a changing ocean: Biological responses of *Emiliana huxleyi* to elevated pCO<sub>2</sub> and their effects on microzooplankton (E Hux Response to pCO<sub>2</sub>)

#### **Description from NSF award abstract:**

The calcifying Haptophyte *Emiliana huxleyi* appears to be acutely sensitive to the rising concentration of ocean pCO<sub>2</sub>. Documented responses by *E. huxleyi* to elevated pCO<sub>2</sub> include modifications to their calcification rate and cell size, malformation of coccoliths, elevated growth rates, increased organic carbon production, lowering of PIC:POC ratios, and elevated production of the active climate gas DMS. Changes in these parameters are mechanisms known to elicit alterations in grazing behavior by microzooplankton, the oceans dominant grazer functional group. The investigators hypothesize that modifications to the physiology and biochemistry of calcifying and non-calcifying Haptophyte *Emiliana huxleyi* in response to elevated pCO<sub>2</sub> will precipitate alterations in microzooplankton grazing dynamics. To test this hypothesis, they will conduct controlled laboratory experiments where several strains of *E. huxleyi* are grown at several CO<sub>2</sub> concentrations. After careful characterization of the biochemical and physiological responses of the *E. huxleyi* strains to elevated pCO<sub>2</sub>, they will provide these strains as food to several ecologically-important microzooplankton and document grazing dynamics. *E. huxleyi* is an ideal organism for the study of phytoplankton and microzooplankton responses to rising anthropogenic CO<sub>2</sub>, the effects of which in the marine environment are called ocean acidification; *E. huxleyi* is biogeochemically important, is well studied, numerous strains are in culture that exhibit variation in the parameters described above, and they are readily fed upon by ecologically important microzooplankton.

The implications of changes in microzooplankton grazing for carbon cycling, specifically CaCO<sub>3</sub> export, DMS production, nutrient regeneration in surface waters, and carbon transfer between trophic levels are profound, as this grazing, to a large degree, regulates all these processes. *E. huxleyi* is a model prey organism because it is one of the most biogeochemically influential global phytoplankton. It forms massive seasonal blooms, contributes significantly to marine inorganic and organic carbon cycles, is a large producer of the climatically active gas DMS, and is a source of organic matter for trophic levels both above and below itself. The planned controlled study will increase our knowledge of the mechanisms that drive patterns of change between trophic levels, thus providing a wider array of tools necessary to understand the complex nature of ocean acidification field studies, where competing variables can confound precise interpretation.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-0961229</a>

[ [table of contents](#) | [back to top](#) ]