

Grazing experiments 4: Daily cell count data for low-high pCO₂ acclimated *Rhodomonas* sp. (E Hux Response to pCO₂ project)

Website: <https://www.bco-dmo.org/dataset/668811>

Data Type: experimental

Version:

Version Date: 2016-12-06

Project

» [Planktonic interactions in a changing ocean: Biological responses of *Emiliana huxleyi* to elevated pCO₂ and their effects on microzooplankton](#) (E Hux Response to pCO₂)

Contributors	Affiliation	Role
Olson, M Brady	Western Washington University (WWU)	Principal Investigator
Love, Brooke	Western Washington University (WWU)	Co-Principal Investigator
Strom, Suzanne	Western Washington University (WWU)	Co-Principal Investigator
Still, Kelly Ann	Western Washington University (WWU)	Student
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Dataset Description

Related Reference:

Still, Kelly Ann, Microzooplankton grazing, growth and gross growth efficiency are affected by pCO₂ induced changes in phytoplankton biology. (Masters Thesis) Western Washington University. <http://cedar.wwu.edu/cgi/viewcontent.cgi?article=1490&context=wwuet>

Methods & Sampling

The phytoplankton *Rhodomonas* sp. CCMP 755 was grown semi-continuously in atmosphere controlled chambers at three different CO₂ treatment concentrations; Ambient (400ppmv), Moderate (750ppmv), and High (1000ppmv). Cultures were started and allowed to grow four days to reach a density of approximately 50,000 cells per ml. Cultures were then diluted daily with pre-equilibrated media containing f/50 nutrients. Each morning of dilution the cultures were gently mixed prior to a small sample being taken for cell counts. Cells were counted live on a Z2 Coulter Particle Counter. The dilution volume was then calculated to achieve a cell density of approximately 25,000 cells per ml which was the density determined in preliminary experiments to be adequate to maintain pCO₂ near target concentrations.

Data Processing Description

These data are unprocessed cell counts.

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- nd (no data) was entered into all blank cells

[[table of contents](#) | [back to top](#)]

Data Files

File
expt4_cell_count_daily.csv (Comma Separated Values (.csv), 894 bytes) MD5:cd3ab8405b4a5380600e7cabbd9582c3
Primary data file for dataset ID 668811

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
treatment_replicate	sample identifier: individual grazer analyzed	unitless
Day_1	Count for day 1 morning cell count	cells/milliliter
day_2	Count for day 2 morning cell count	cells/milliliter
day_2_post_dilution	Count for day 2 post-dilution; for the predicted cell count after dilution	cells/milliliter
day_3	Count for day 3 morning cell count	cells/milliliter
day_3_post_dilution	Count for day 3 post-dilution; for the predicted cell count after dilution	cells/milliliter
day_4	Count for day 4 morning cell count	cells/milliliter
day_4_post_dilution	Count for day 4 post-dilution; for the predicted cell count after dilution	cells/milliliter
day_5	Count for day 5 morning cell count	cells/milliliter
day_5_post_dilution	Count for day 5 post-dilution; for the predicted cell count after dilution	cells/milliliter
day_6	Count for day 6 morning cell count	cells/milliliter
day_6_post_dilution	Count for day 6 post-dilution; for the predicted cell count after dilution	cells/milliliter
day_7	Count for day 7 morning cell count	cells/milliliter

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Z2 Coulter Particle Counter
Generic Instrument Name	Coulter Counter
Dataset-specific Description	Used to count cells
Generic Instrument Description	An apparatus for counting and sizing particles suspended in electrolytes. It is used for cells, bacteria, prokaryotic cells and virus particles. A typical Coulter counter has one or more microchannels that separate two chambers containing electrolyte solutions. from https://en.wikipedia.org/wiki/Coulter_counter

[[table of contents](#) | [back to top](#)]

Deployments

Lab Olson B

Website	https://www.bco-dmo.org/deployment/521277
Platform	WWU
Start Date	2011-03-31
End Date	2016-09-15
Description	laboratory experiments

[[table of contents](#) | [back to top](#)]

Project Information

Planktonic interactions in a changing ocean: Biological responses of *Emiliana huxleyi* to elevated pCO₂ and their effects on microzooplankton (E Hux Response to pCO₂)

Description from NSF award abstract:

The calcifying Haptophyte *Emiliana huxleyi* appears to be acutely sensitive to the rising concentration of ocean pCO₂. Documented responses by *E. huxleyi* to elevated pCO₂ include modifications to their calcification rate and cell size, malformation of coccoliths, elevated growth rates, increased organic carbon production, lowering of PIC:POC ratios, and elevated production of the active climate gas DMS. Changes in these parameters are mechanisms known to elicit alterations in grazing behavior by microzooplankton, the oceans dominant grazer functional group. The investigators hypothesize that modifications to the physiology and biochemistry of calcifying and non-calcifying Haptophyte *Emiliana huxleyi* in response to elevated pCO₂ will precipitate alterations in microzooplankton grazing dynamics. To test this hypothesis, they will conduct controlled laboratory experiments where several strains of *E. huxleyi* are grown at several CO₂ concentrations. After careful characterization of the biochemical and physiological responses of the *E. huxleyi* strains to elevated pCO₂, they will provide these strains as food to several ecologically-important microzooplankton and document grazing dynamics. *E. huxleyi* is an ideal organism for the study of phytoplankton and microzooplankton responses to rising anthropogenic CO₂, the effects of which in the marine environment are called ocean acidification; *E. huxleyi* is biogeochemically important, is well studied, numerous strains are in culture that exhibit variation in the parameters described above, and they are readily fed upon by ecologically important microzooplankton.

The implications of changes in microzooplankton grazing for carbon cycling, specifically CaCO₃ export, DMS

production, nutrient regeneration in surface waters, and carbon transfer between trophic levels are profound, as this grazing, to a large degree, regulates all these processes. *E. huxleyi* is a model prey organism because it is one of the most biogeochemically influential global phytoplankton. It forms massive seasonal blooms, contributes significantly to marine inorganic and organic carbon cycles, is a large producer of the climatically active gas DMS, and is a source of organic matter for trophic levels both above and below itself. The planned controlled study will increase our knowledge of the mechanisms that drive patterns of change between trophic levels, thus providing a wider array of tools necessary to understand the complex nature of ocean acidification field studies, where competing variables can confound precise interpretation.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0961229

[[table of contents](#) | [back to top](#)]