16S rRNA gene sequences accessions from iron mats sampled during the R/V Thomas G. Thompson cruise TN293 in the Loihi Seamount, Hawaii from March to April 2013 (Ecology of Vent Mats project)

Website: https://www.bco-dmo.org/dataset/707550

Data Type: experimental

Version:

Version Date: 2017-06-30

Project

» Ecology of microbial mats at seamount associated Fe-rich hydrothermal vent systems (Ecology of Vent Mats)

| Contributors | Affiliation | Role |
|----------------|---|---------------------------------|
| Emerson, David | Bigelow Laboratory for Ocean Sciences | Principal Investigator, Contact |
| York, Amber D. | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Table of Contents

- Coverage
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Related Publications
- Parameters
- Instruments
- Deployments
- Project Information
- <u>Funding</u>

Coverage

Spatial Extent: Lat:18.92 Lon:-155.27

Dataset Description

This dataset contains accession numbers and links to NCBI GenBank for 16S rRNA gene sequences from iron mats for BioProject PRJEB10821. Samples were collected during the R/V Thomas G. Thompson cruise TN293 and used the ROV Jason 2 (dives J2-671 - J2-677) at Loihi Seamount, summit area (1300m) to base (5000m), lat/lon (18.9200, -155.2700).

These accessions are also available at the European Nucleotide Archive under the sample accessions numbers ERS1089449–ERS1089534, study accession number PRIEB10821.

Methods & Sampling

Microbial mat samples acquired with sub-centimeter sampling resolution from the seafloor using a novel microbial mat sampling device (see Breier, et al. 2012. for details) operated by the ROV Jason 2. Microbial mat samples were frozen on board the ship and returned to the laboratory for DNA extraction and analysis. The V4-V5 region of the 16S rRNA gene was sequenced using pyrosequencing (titanium chemistry) as described in Scott et al. 2015.

Data Processing Description

Raw data was processed using mothur and oligotyping/minimum entropy decomposition.

[table of contents | back to top]

Data Files

File

Fe_mat_accessions.csv(Comma Separated Values (.csv), 9.96 KB)

MD5:284b818c43c6d487cbb53ff1c6b13f32

Primary data file for dataset ID 707550

[table of contents | back to top]

Related Publications

Breier, J. A., Gomez-Ibanez, D., Reddington, E., Huber, J. A., & Emerson, D. (2012). A precision multi-sampler for deep-sea hydrothermal microbial mat studies. Deep Sea Research Part I: Oceanographic Research Papers, 70, 83–90. doi:10.1016/j.dsr.2012.10.006

Methods

Scott, J. J., Breier, J. A., Luther, G. W., & Emerson, D. (2015). Microbial Iron Mats at the Mid-Atlantic Ridge and Evidence that Zetaproteobacteria May Be Restricted to Iron-Oxidizing Marine Systems. PLOS ONE, 10(3), e0119284. doi:10.1371/journal.pone.0119284

Methods

[table of contents | back to top]

Parameters

| Parameter | Description | Units |
|----------------|---------------------------|----------|
| accession_id | Accession number at NCBI | unitless |
| accession_link | Link to accession at NCBI | unitless |

[table of contents | back to top]

Instruments

| Dataset- specific Instrument Name | ROV Jason 2 |
|--|---|
| Generic Instrument Name | ROV Jason |
| Dataset- specific Description | Dives J2-671 - J2-677 |
| | The Remotely Operated Vehicle (ROV) Jason is operated by the Deep Submergence Laboratory (DSL) at Woods Hole Oceanographic Institution (WHOI). WHOI engineers and scientists designed and built the ROV Jason to give scientists access to the seafloor that didn't require them leaving the deck of the ship. Jason is a two-body ROV system. A 10-kilometer (6-mile) fiber-optic cable delivers electrical power and commands from the ship through Medea and down to Jason, which then returns data and live video imagery. Medea serves as a shock absorber, buffering Jason from the movements of the ship, while providing lighting and a bird's eye view of the ROV during seafloor operations. During each dive (deployment of the ROV), Jason pilots and scientists work from a control room on the ship to monitor Jason's instruments and video while maneuvering the vehicle and optionally performing a variety of sampling activities. Jason is equipped with sonar imagers, water samplers, video and still cameras, and lighting gear. Jason's manipulator arms collect samples of rock, sediment, or marine life and place them in the vehicle's basket or on "elevator" platforms that float heavier loads to the surface. More information is available from the operator site at URL. |

[table of contents | back to top]

Deployments

TN293

| Website | https://www.bco-dmo.org/deployment/659705 |
|-------------|--|
| Platform | R/V Thomas G. Thompson |
| Start Date | 2013-03-16 |
| End Date | 2013-04-01 |
| Description | For Jason data/information from this cruise, please visit the Jason Virtual Control Van: http://4dgeo.whoi.edu/webdata/virtualvan/html/VV-tn293/index.html (This link includes all navigational data, images, and sample collection metadata and comments for the cruise. Data can be downloaded with export viewer.) Additional cruise information and original data are available from the NSF R2R data catalog. |

[table of contents | back to top]

Project Information

Ecology of microbial mats at seamount associated Fe-rich hydrothermal vent systems (Ecology of Vent Mats)

Website: http://oceanexplorer.noaa.gov/explorations/14fire/welcome.html

Coverage: Loihi Seamount, Hawaii; and Mariana Arc and Backarc Hydrothermal Systems

Description from NSF award abstract:

A grand challenge in microbial ecology is to understand what drives the structure of microbial communities. A recently discovered novel class of Proteobacteria, the Zetaproteobacteria, are associated with microbial mats at iron rich hydrothermal vents at submarine volcanoes deep in the ocean. These bacteria only grow using iron as an energy source and fix carbon dioxide. Within iron rich microbial mats, Zetaproteobacteria are the dominant bacterial population; however they are rare in most other deep-sea or marine habitats, suggesting they may be restricted to specific niches characterized by gradients of oxygen and iron. Recent discoveries have expanded their range to fluids collected from deep ocean crust boreholes, iron deposits in coastal saltmarshes, and with steel associated bio-corrosion, demonstrating that marine Zetaproteobacteria are cosmopolitan. A unique property of these marine iron oxidizing bacteria is that they produce morphologically distinct iron oxide structures in the form of filamentous sheaths or stalk-like structures. These structures are easily recognized by light microscopy, and electron microscopy is beginning to reveal subtle differences among them that may be diagnostic of different populations of iron oxidizing bacteria. Another unusual aspect of iron oxidizing bacteria is that they produce large quantities of oxides with relatively little bacterial biomass. As a result, the oxides form a matrix that influences water and nutrient flow in the microbial mats where they grow, and in turn, may influence the growth of other groups of bacteria and archaea that live in the mats. In an ecological context, the PIs believe this makes them a keystone species that form the predominant structural matrix of the mat, and engineer an environment conducive for growth of specific bacterial populations within the mat ecosystem. The PIs propose to use high resolution mat sampling techniques to investigate the architecture of mat ecosystems and couple these with modern molecular methods (i.e., single-cell metagenomics) and geochemical measurements of the vent fluid to couple morphological and functional diversity to phylogenetic and physiological diversity. Because the Zetaproteobacteria are ancient, have unique metabolic and morphological attributes, and appear to be restricted to a well-defined habitat, they offer an interesting model for understanding fundamental ecological concepts that drive microbial diversity and evolution.

A better understanding of iron oxidizing bacteria that include Zetaproteobacteria is of fundamental interest to scientists interested in areas of earth science and oceanography because they illustrate how microbes can fundamentally influence geochemical cycling and mineral deposition. Furthermore, morphological structures similar to those produced by Zetaproteobacteria can still be identified hundreds of millions (and possibly billions) of years back in the geological record, making them of paleontological, and potentially of exobiological, interest. As knowledge of extant populations grow, it is possible they will also help to inform us of environmental change in past Earth history. A wealth of educational and outreach opportunities will be made possible by this work, including graduate and postdoctoral education, research experiences for undergraduates, and teacher training. In addition the participating scientists are involved in a number of programs to make the general public aware of the process of how scientific research is conducted, and how discoveries of a fundamental nature can ultimately benefit humankind.

Additional information/resources: TN293 (Loihi 2013)

Loihi Summit Map (PDF)

Cruise blog: https://zetahunters.wordpress.com/

Jason Virtual Control Van: http://4dgeo.whoi.edu/webdata/virtualvan/html/VV-tn293/index.html

Related Publications:

Fullerton, H., K. W. Hager, S. M. McAllister, and C. L. Moyer. 2017. Hidden diversity revealed by genome-resolved metagenomics of iron-oxidizing microbial mats from Lō'ihi Seamount, Hawai'i. ISMEJ 11:1900–1914. doi:10.1038/ismej.2017.40

Emerson, D., J. J. Scott, A. Leavitt, E. Fleming, and C. L. Moyer. 2016. In situ estimates of iron-oxidation and accretion rates for iron-oxidizing bacterial mats at Loihi Seamount. bioRxiv 095414. doi:10.1101/095414 **Scott**, J. J., B. T. Glazer, and D. Emerson. 2017. Bringing microbial diversity into focus: high-resolution analysis of iron mats from the Lō'ihi Seamount. Environmental Microbiology 19:301–316. doi:10.1111/1462-2920.13607

Chan, C.S., S.M. McAllister, A.H. Leavitt, B.T. Glazer, S.T. Krepski, and D. Emerson. 2016. The architecture of iron microbial mats reflects the adaptation of chemolithotrophic iron oxidation in freshwater and marine environments. Frontiers in Microbiology 7:796. doi:10.3389/fmicb.2016.00796

Fullerton, H., K. W. Hager, and C. L. Moyer. 2015. Draft genome sequence of Mariprofundus ferrooxydans strain JV-1, isolated from Loihi Seamount, Hawaii. Genome announcements 3:e01118-15. doi:10.1128/genomeA.01118-15

Field, E.K., A. Sczyrba, A.E. Lyman, C.C. Harris, T. Woyke, R. Stepanauskas, and D. Emerson. 2015. Genomic insights into the uncultivated marine Zetaproteobacteria at Loihi Seamount. ISMEJ 9:857–870. doi:10.1038/ismej.2014.183

Jesser, KJ, Fullerton H, Hager KW, Moyer CL. 2015. Quantitative PCR analysis of functional genes in iron-rich microbial mats at an active hydrothermal vent system (Lō'ihi Seamount, Hawai'i). Appl Environ Microbiol

81:2976-2984. doi:10.1128/AEM.03608-14. (PDF)

RR1413 (Mariana 2014)

RR1413 Cruise Report (5.2 MB PDF)

<u>Urushima to Rota Map (PDF)</u>

Jason Virtual Control Van website: http://4dgeo.whoi.edu/webdata/virtualvan/html/VV-rr1413/index.html

Related Publications:

Hager, K. W., H. Fullerton, D. A. Butterfield, and C. L. Moyer. 2017. Community structure of lithotrophically-driven hydrothermal microbial mats from the Mariana Arc and Back-Arc. Frontiers in Microbiology 8:1578. doi:10.3389/fmicb.2017.01578

[table of contents | back to top]

Funding

| Funding Source | Award |
|--|-------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-1155754 |

[table of contents | back to top]