

Survival, length, and growth responses of *M. menidia* offspring from different females exposed to contrasting CO2 environments.

Website: <https://www.bco-dmo.org/dataset/719420>

Data Type: experimental

Version: Final

Version Date: 2017-11-14

Project

» [Collaborative research: Understanding the effects of acidification and hypoxia within and across generations in a coastal marine fish](#) (HYPOA)

Contributors	Affiliation	Role
Baumann, Hannes	University of Connecticut (UConn)	Principal Investigator, Contact
Nye, Janet	Stony Brook University - SoMAS (SUNY-SB SoMAS)	Co-Principal Investigator
Ake, Hannah	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: Lat:41.3213389 Lon:-72.0148361

Dataset Description

Wild-caught *M. menidia* adults were spawned to test whether offspring from different mothers differ in their average survival and size responses to elevated CO2 conditions. The experiment quantified three related survival and three size traits for each replicate, female, and CO2 treatment: embryo survival (fertilization to 1 dph), larval survival (1 to 16 dph), and overall survival (fertilization to 16 dph); and size (SL) at hatch (1 dph), SL at 16 dph, and larval growth rate ($GR = (SL_{16dph} - SL_{1dph})/15$).

These data are associated with the corresponding paper:

[Snyder, J.T.*, Murray, C.S.*, and Baumann, H. \(2017\) Potential for maternal effects on offspring CO2-sensitivities in a coastal marine fish. *Journal of Experimental Marine Biology and Ecology* \(in press\).](#)

Other datasets related to this paper:

[Fatty acid profiles of *M. menidia* females and their unfertilized eggs.](#)

Methods & Sampling

Methodology from **Snyder, J.T.*, Murray, C.S.*, and Baumann, H. (2017) Potential for maternal**

effects on offspring CO₂-sensitivities in a coastal marine fish. *Journal of Experimental Marine Biology and Ecology* (in press).

Five randomly selected females were strip-spawned onto cutout sections of window screen (1-mm mesh) that were placed into separate seawater-filled spawning dishes (Murray et al., 2014). To ensure full fertilization success and randomize potential paternal effects, eggs were fertilized with a mixture of milt from 22 males, thus producing full-sib and maternal half-sib embryos from each female. Adults were measured for total length (TL; mean TL_{male} = 9.14 cm, mean TL_{female} = 10.4 cm) and frozen for later analysis of FA. Mesh screens with attached embryos were subsequently cut into smaller sections to allow precise enumeration, and within 2-hr post-fertilization 100 embryos were placed into each of three replicate rearing containers (20 L) per female and CO₂ treatment (i.e., 600 embryos for each of five females, 3 × 100 in ambient and 3 × 100 in acidified treatments). Rearing containers were filled with 1-um filtered, UV-sterilized seawater (~30 psu) from Long Island Sound and placed in temperature-controlled water baths set to 24 deg C, the known thermal optimum for survival and growth in this species (Middaugh et al., 1987). Offspring were reared for 24 d post fertilization under a 15h light:9h dark lighting regime. After hatch, larvae were fed ad libitum rations of newly hatched brine shrimp nauplii *Artemia salina* (brineshrimpdirect.com), and 50% of water was replaced every 5 d to ensure safe ammonia levels (< 0.25 ppm). Hatched larvae were counted and subsampled (n = 10 per replicate) at 1 d post hatch (dph) by gently scooping them into identical 20 L containers, and final samples were taken at 16 dph. All samples were preserved in 5% buffered formalin for later measurements of larval standard length (SL, 0.01 mm) via calibrated digital images (ImagePro Premier, MediaCybernetics). The experiment thus quantified three related survival and three size traits for each replicate, female, and CO₂ treatment: embryo survival (fertilization to 1 dph), larval survival (1 to 16 dph), overall survival (fertilization to 16 dph), size (SL) at hatch (1 dph), SL at 16 dph, and larval growth rate (GR = (SL_{16dph} - SL_{1dph})/15).

CO₂ regime:

Offspring were reared at ambient (~ 400 uatm, pH_{NBS} = 8.18) and acidified CO₂ conditions (~2,300 uatm, pH_{NBS} = 7.50). The higher value was set to a level commonly used in OA research (consistent with projections of future pCO₂ values for open oceans over in the next 200 yr (IPCC, 2007)) and represents current conditions experienced during seasonal extremes by this species in nature (Murray et al., 2014). Ambient conditions were achieved by bubbling partially CO₂-stripped air into each rearing container, thereby offsetting metabolic CO₂ accumulation. Acidified conditions were achieved via gas proportioners (Cole Parmer®) that mixed CO₂ stripped air with 100% bone-dry CO₂ delivered to the bottom of each rearing container via air stones. Target pH and temperature were monitored daily via a handheld pH probe (Hach® HQ40d portable meter with a PHC201 standard pH-probe) calibrated regularly via two-point National Bureau of Standards (NBS) pH buffers (electronic supplementary material, Fig.S1). To characterize actual pCO₂ levels and related water chemistry parameters, water was sampled from four randomly chosen rearing containers per treatment three times over the course of the experiment and immediately measured for total alkalinity (AT) via endpoint titration (Mettler Toledo™ G20 Potentiometric Titrator). The instrument has previously been shown to quantify AT in Dr. Andrew Dickson's reference material (batch 147, AT= 2231.39 umol kg seawater⁻¹) with an average error of 0.6%. Actual levels of total dissolved inorganic carbon (CT), partial pressure of CO₂ (pCO₂), fugacity of CO₂ (fCO₂), and carbonate ion concentration were calculated in CO₂SYS (<http://cdiac.ornl.gov/ftp/co2sys>) based on measured AT, pH (NBS), temperature, and salinity using K1 and K2 constants from Mehrbach et al. (1973) refit by Dickson and Millero (1987) and Dickson (1990) for KHSO₄ (Table 1).

Data Processing Description

BCO-DMO Data Processing Notes:

- added underscores to column headers
- replaced blank cells with nd
- added underscores to site name
- removed commas from data
- changed date format to yyyy/mm/dd

[[table of contents](#) | [back to top](#)]

Data Files

File

survival_size.csv(Comma Separated Values (.csv), 3.30 KB)
MD5:b3b943d7b26084a0b4423aeb3f3fa219

Primary data file for dataset ID 719420

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Species	Atlantic silverside; Menidia menidia	unitless
Collection_site	Mumford Cove Connecticut USA	unitless
Lat	Latitude of field collection site	decimal degrees
Lon	Longitude of field collection site	decimal degrees
Collection_date	Date of field collection for the spawners used in the experiment; YYYY/MM/DD	unitless
Female	Five females denoted by letter A B C D E	unitless
Total_length	Fish were measured to the lower half centimeter; lower 0.5 centimeters	centimeters
pH	pH units reported on the NIST scale	NIST units
pCO2	Calculated CO2 partial pressure	uatm
Replicate	There were 3 replicates labled 1 2 3	unitless
Embryo_survival	Relative survival of offspring from fertilization to 1 day post hatch	percent
Larval_survival	Relative survival of offspring from data 1 to day 16 post hatch	percent
Overall_survival	Relative survival of offspring from fertilization to day 16 post hatch	percent
Hatch_length	Mean length of newly hatched larvae per replicate	millimeters
Larval_length	Mean length of larvae at the end of the experiment per replicate	millimeters
Growth	Average growth of larvae from d1 to d16 per replicate	millimeters per day

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Mettler Toledo™ G20 Potentiometric Titrator
Generic Instrument Name	Automatic titrator
Dataset-specific Description	Used to measure total alkalinity
Generic Instrument Description	Instruments that incrementally add quantified aliquots of a reagent to a sample until the end-point of a chemical reaction is reached.

Dataset-specific Instrument Name	Hach® HQ40d portable meter with a PHC201 standard pH-probe
Generic Instrument Name	pH Sensor
Dataset-specific Description	handheld pH probe
Generic Instrument Description	An instrument that measures the hydrogen ion activity in solutions. The overall concentration of hydrogen ions is inversely related to its pH. The pH scale ranges from 0 to 14 and indicates whether acidic (more H+) or basic (less H+).

[[table of contents](#) | [back to top](#)]

Deployments

AP_Rankin

Website	https://www.bco-dmo.org/deployment/651907
Platform	Avery_Point
Start Date	2015-05-03
End Date	2015-09-15
Description	This was where the Long-term Menidia menidia growth experiments took place. The samples were collected from offshore in Mumford Cove.

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative research: Understanding the effects of acidification and hypoxia within and across generations in a coastal marine fish (HYPOA)

Coverage: Eastern Long Island Sound, CT, USA

Description from NSF award abstract:

Coastal marine ecosystems provide a number of important services and resources for humans, and at the same time, coastal waters are subject to environmental stressors such as increases in ocean acidification and reductions in dissolved oxygen. The effects of these stressors on coastal marine organisms remain poorly understood because most research to date has examined the sensitivity of species to one factor, but not to more than one in combination. This project will determine how a model fish species, the Atlantic silverside, will respond to observed and predicted levels of dissolved carbon dioxide (CO₂) and oxygen (O₂). Shorter-term experiments will measure embryo and larval survival, growth, and metabolism, and determine whether parents experiencing stressful conditions produce more robust offspring. Longer-term experiments will study the consequences of ocean acidification over the entire life span by quantifying the effects of high-CO₂ conditions on the ratio of males to females, lifetime growth, and reproductive investment. These studies will provide a more comprehensive view of how multiple stressors may impact populations of Atlantic silversides and potentially other important forage fish species. This collaborative project will support and train three graduate students at the University of Connecticut and the Stony Brook University (NY), two institutions that attract students from minority groups. It will also provide a variety of opportunities for undergraduates to participate in research and the public to learn about the study, through summer research projects, incorporation in the "Women in Science and Engineering" program, and interactive displays of environmental data from monitoring

buoys. The two early-career investigators are committed to increasing ocean literacy and awareness of NSF-funded research through public talks and presentations.

This project responds to the recognized need for multi-stressor assessments of species sensitivities to anthropogenic environmental change. It will combine environmental monitoring with advanced experimental approaches to characterize early and whole life consequences of acidification and hypoxia in the Atlantic silverside (*Menidia menidia*), a valued model species and important forage fish along most of the US east coast. Experiments will employ a newly constructed, computer-controlled fish rearing system to allow independent and combined manipulation of seawater pCO₂ and dissolved oxygen (DO) content and the application of static and fluctuating pCO₂ and DO levels that were chosen to represent contemporary and potential future scenarios in productive coastal habitats. First CO₂, DO, and CO₂ × DO dependent reaction norms will be quantified for fitness-relevant early life history (ELH) traits including pre- and post-hatch survival, time to hatch, post-hatch growth, by rearing offspring collected from wild adults from fertilization to 20 days post hatch (dph) using a full factorial design of 3 CO₂ × 3 DO levels. Second, the effects of tidal and diel CO₂ × DO fluctuations of different amplitudes on silverside ELH traits will be quantified. To address knowledge gaps regarding the CO₂-sensitivity in this species, laboratory manipulations of adult spawner environments and reciprocal offspring exposure experiments will elucidate the role of transgenerational plasticity as a potential short-term mechanism to cope with changing environments. To better understand the mechanisms of fish early life CO₂-sensitivity, the effects of temperature × CO₂ on pre- and post-hatch metabolism will be robustly quantified. The final objective is to rear silversides from fertilization to maturity under different CO₂ levels and assess potential CO₂-effects on sex ratio and whole life growth and fecundity.

Related references:

Gobler, C.J. and Baumann, H. (2016) Hypoxia and acidification in ocean ecosystems: Coupled dynamics and effects on marine life. *Biology Letters* 12:20150976. doi:[10.1098/rsbl.2015.0976](https://doi.org/10.1098/rsbl.2015.0976)

Baumann, H. (2016) Combined effects of ocean acidification, warming, and hypoxia on marine organisms. *Limnology and Oceanography e-Lectures* 6:1-43. doi:[10.1002/loe2.10002](https://doi.org/10.1002/loe2.10002)

Depasquale, E., Baumann, H., and Gobler, C.J. (2015) Variation in early life stage vulnerability among Northwest Atlantic estuarine forage fish to ocean acidification and low oxygen *Marine Ecology Progress Series* 523: 145-156. doi:[10.3354/meps11142](https://doi.org/10.3354/meps11142)

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1536336

[[table of contents](#) | [back to top](#)]