

# Porewater sulfate from samples collected by pushcore from Guaymas Basin hydrothermal sediments on R/V Atlantis cruise AT37-06 in the Guaymas Basin in December 2016

**Website:** <https://www.bco-dmo.org/dataset/720669>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2017-12-08

## Project

» [Collaborative Research: Microbial Carbon cycling and its interactions with Sulfur and Nitrogen transformations in Guaymas Basin hydrothermal sediments](#) (Guaymas Basin Interactions)

Contributors	Affiliation	Role
<a href="#">Teske, Andreas</a>	University of North Carolina at Chapel Hill (UNC-Chapel Hill)	Principal Investigator
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Porewater sulfate from samples collected by pushcore from Guaymas Basin hydrothermal sediments on cruise AT37-06 in December 2016.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** Lat:27 Lon:-111

---

## Dataset Description

Porewater sulfate from samples collected by pushcore from Guaymas Basin hydrothermal sediments on cruise AT37-06 in December 2016.

Publications associated with this project are as follows:

Schutte, C.A., A. Teske, B.J. MacGregor, V. Salman-Carvalho, G. Lavik, and D. de Beer. Filamentous giant *Beggiatoaceae* from Guaymas Basin are capable of both denitrification and dissimilatory nitrate reduction to ammonium (DNRA). For submission.

## Methods & Sampling

Sampling and Analytical Methodology: Freshly collected sediment cores were sliced on the ship into 3 cm layers; porewater was obtained by gently centrifuging freshly collected sediment in 50 ml conical Falcon tubes for ca. 5 to 10 minutes until the sediment had settled; one Falcon tube produced ca. 8 to 10 ml of porewater. For porewater sulfate measurements, 1 ml subsamples of the overlying porewater were drawn into syringes and injected through 0.45 µm filters into screw cap Eppendorf vials, each acidified with 50 µl of 6N HCl, and

then gently bubbled with nitrogen for 4 min to remove sulfide; the samples were then stored at 4°C before shipping and analysis.

Geochemical Analyses: SO<sub>4</sub> was analyzed on a Dionex ICS-1000 using a RFIC IonPac AS22 4 X 250mm column and a RFIC IonPac AG22 Guard column 4 X 50 mm. The IC was calibrated using a 5 point std curve and a minimum r<sup>2</sup> value of 0.999. The MDL measured at 0.10 ppm SO<sub>4</sub>. A check standard from a 1000 ppm was made from a Dionex 7 anion standard to a concentration of 15 ppm. An external Standard from HACH 100 ppm SO<sub>4</sub> was used to make a 20 ppm QC standard. In general, QCs were analyzed for 10% of the samples.

Data Processing: The porewater data of Guaymas Basin sand Sonora Margin piston cores were tabulated in Excel sheets.

Quality Control: Note that Sample\_IDs "D7A\_10X\_CKS" are internal quality control standards (15.5 mg/L) and Sample\_IDs "DI\_SPK" are external quality control standards (20.0 mg/L).

## Data Processing Description

BCO-DMO Processing:

- modified parameter names (replaced spaces with underscores);
- replaced commas in Sample\_ID column with semi-colons.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>AT37-06_SO4.csv</b> (Comma Separated Values (.csv), 1.01 KB) MD5:84e75fdeeb5c3db8415383b0e980f481
Primary data file for dataset ID 720669

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Sample_ID	Sample identification number	unitless
Sulfate_Concentration	Sulfate concentration	millimoles per liter (mMol/L)

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Alvin tube core
<b>Generic Instrument Description</b>	A plastic tube, about 40 cm (16 inches) long, is pushed into the sediment by Alvin's manipulator arm to collect a sediment core.

<b>Dataset-specific Instrument Name</b>	Dionex ICS-1000
<b>Generic Instrument Name</b>	Ion Chromatograph
<b>Dataset-specific Description</b>	SO4 was analyzed on a Dionex ICS-1000 using a RFIC IonPac AS22 4 X 250mm column and a RFIC IonPac AG22 Guard column 4 X 50 mm.
<b>Generic Instrument Description</b>	Ion chromatography is a form of liquid chromatography that measures concentrations of ionic species by separating them based on their interaction with a resin. Ionic species separate differently depending on species type and size. Ion chromatographs are able to measure concentrations of major anions, such as fluoride, chloride, nitrate, nitrite, and sulfate, as well as major cations such as lithium, sodium, ammonium, potassium, calcium, and magnesium in the parts-per-billion (ppb) range. (from <a href="http://serc.carleton.edu/microbelife/research_methods/biogeochemical/ic....">http://serc.carleton.edu/microbelife/research_methods/biogeochemical/ic....</a> )

[ [table of contents](#) | [back to top](#) ]

## Deployments

### AT37-06

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/720354">https://www.bco-dmo.org/deployment/720354</a>
<b>Platform</b>	R/V Atlantis
<b>Report</b>	<a href="https://datadocs.bco-dmo.org/d3/data_docs/GuaymasBasin_Interactions/AT37-06_CruiseReport.pdf">https://datadocs.bco-dmo.org/d3/data_docs/GuaymasBasin_Interactions/AT37-06_CruiseReport.pdf</a>
<b>Start Date</b>	2016-12-09
<b>End Date</b>	2016-12-27

[ [table of contents](#) | [back to top](#) ]

## Project Information

### **Collaborative Research: Microbial Carbon cycling and its interactions with Sulfur and Nitrogen transformations in Guaymas Basin hydrothermal sediments (Guaymas Basin Interactions)**

**Coverage:** Guaymas Basin, Gulf of California, 27.00 N, 111.00W

#### *Description from NSF award abstract:*

Hydrothermally active sediments in the Guaymas Basin are dominated by novel microbial communities that catalyze important biogeochemical processes in these seafloor ecosystems. This project will investigate genomic potential, physiological capabilities and biogeochemical roles of key uncultured organisms from Guaymas sediments, especially the high-temperature anaerobic methane oxidizers that occur specifically in hydrothermally active sediments (ANME-1Guaymas). The study will focus on their role in carbon transformations, but also explore their potential involvement in sulfur and nitrogen transformations. First-order research topics include quantifying anaerobic methane oxidation under high temperature, in situ concentrations of phosphorus and methane, and with alternate electron acceptors; sulfate and sulfur-dependent microbial pathways and isotopic signatures under these conditions; and nitrogen transformations in methane-oxidizing microbial communities, hydrothermal mats and sediments.

This integrated biogeochemical and microbiological research will explore the pathways of and environmental controls on the consumption and production of methane, other alkanes, inorganic carbon, organic acids and organic matter that fuel the Guaymas sedimentary microbial ecosystem. The hydrothermal sediments of Guaymas Basin provide a spatially compact, high-activity location for investigating novel modes of methane cycling and carbon assimilation into microbial biomass. In the case of anaerobic methane oxidation, the high temperature and pressure tolerance of Guaymas Basin methane-oxidizing microbial communities, and their potential to uncouple from the dominant electron acceptor sulfate, vastly increase the predicted subsurface habitat space and biogeochemical role for anaerobic microbial methanotrophy in global deep subsurface diagenesis. Further, microbial methane production and oxidation interlocks with sulfur and nitrogen transformations, which will be explored at the organism and process level in hydrothermal sediment microbial communities and mats of Guaymas Basin. In general, first-order research tasks (rate measurements, radiotracer incorporation studies, genomes, in situ microgradients) define the key microbial capabilities, pathways and processes that mediate chemical exchange between the subsurface hydrothermal/seeps and deep ocean waters.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1357238</a>

[ [table of contents](#) | [back to top](#) ]