Initial and final measured buoyant weights, final measured dry weight, and the initial dry weight (estimated from buoyant-dry weight regression) of the urchins reared under the experimental conditions.

Website: https://www.bco-dmo.org/dataset/721312 Version: 1 Version Date: 2017-12-19

Project

» <u>A combined boron isotope, pH microelectrode and pH-sensitive dye approach to constraining acid/base</u> <u>chemistry in the calcifying fluids of corals</u> (CoralCalcifyFluid_pH)

Contributors	Affiliation	Role
<u>Ries, Justin B.</u>	Northeastern University	Principal Investigator
<u>Switzer, Megan</u>	Woods Hole Oceanographic Institution (WHOI)	BCO-DMO Data Manager

Table of Contents

- <u>Dataset Description</u>
- Data Files
- <u>Parameters</u>
- Project Information
- <u>Funding</u>

Dataset Description

Experimental data for echinoid calcification under different temperature and pCO2 conditions. This dataset includes measurements for buoyant weights of echinoids at the beginning and end of the experiment used to estimate calcification rates.

See also datasets for seawater chemistry and stable isotopes. Echinoid seawater chemistry Echinoid stable isotopes

Data are published in:

Courtney, T., Ries, J. B., 2015, Impact of atmospheric pCO_2 , seawater temperature, and calcification rate on the $\delta^{18}O$ and $\delta^{13}C$ composition of echinoid calcite (*Echinometra viridis*). Chemical Geology 411: 228-239. DOI: <u>j.chemgeo.2015.06.030</u>

Please see manuscript for complete methodology.

[table of contents | back to top]

Data Files

File	
721312.csv (Comma Separated Values (.csv), 1.24 KB) MD5:172177720b853b055dc33b68f51aaeba	

```
Primary data file for dataset ID 721312
```

```
[ table of contents | back to top ]
```

Parameters

Parameter	Description	Units
Treatment	Temperature and CO2 concentration for the indicated experiment (experimental conditions)	no units
Initial_Buoyant_Weight	Initial buoyant weight	mg
Final_Buoyant_Weight	Final Buoyant weight	mg
Percent_Change_Buoyant_Weight	Percent change buoyant weight	percent
Initial_Dry_Weight	Initial dry weight	mg
Final_Dry_Weight	Final dry weight	mg

[table of contents | back to top]

Project Information

A combined boron isotope, pH microelectrode and pH-sensitive dye approach to constraining acid/base chemistry in the calcifying fluids of corals (CoralCalcifyFluid_pH)

Website: http://nuweb2.neu.edu/rieslab/

Coverage: Marine Science Center, Northeastern University

Description from NSF award abstract:

The anthropogenic elevation of atmospheric CO2 is causing the oceans to become more acidic, which may make it more challenging for corals to build their skeletons and, ultimately, entire reef structures. How corals respond to future ocean acidification will largely depend on how the pH of the internal fluid from which they produce their skeletons-their so-called calcifying fluid-is impacted by the surrounding seawater. It is therefore essential that current methods are refined to accurately measure the pH of corals' calcifying fluids in order to understand and, ideally, predict their responses to CO2-induced ocean acidification. In this project, a three-pronged approach to measure calcifying fluid pH within three species of reef-forming corals will be used to assess how their calcifying fluid pH responds to experimentally induced ocean acidification. This research will improve our understanding of corals' responses to ocean acidification and thus has the potential to inform the decisions of policy makers and legislators seeking to mitigate the deleterious effects of rising atmospheric CO2 on marine ecosystems. The work will support the development of three early career scientists, a postdoctoral fellow, graduate students, and undergraduate researcher assistants-several of whom are from underrepresented groups in the earth and ocean sciences. Results will be widely disseminated through publications, conference presentations, the PIs' websites, an educational film, coursework, and outreach activities at area schools, museums, and science centers.

Corals and other types of marine calcifiers are thought to begin the mineralization of their calcium carbonate skeletons by actively elevating pH of their calcifying fluid, thereby converting bicarbonate ions (comprising ~90% of seawater dissolved inorganic carbon) to carbonate ions, the form of carbon used in calcification. This project will compare the combined boron isotope, pH microelectrode, and pH-sensitive dye approach to measure the calcifying fluid pH of three species of scleractinian corals, and to assess how their calcifying fluid pH (a primary factor controlling their calcification) responds to experimentally induced ocean acidification. As a result this multi-pronged approach to measuring calcifying fluid pH of the same coral species under equivalent

culturing conditions will permit the first systematic cross-examination of the validity of these independent approaches. The combined approach will also yield values of calcifying fluid pH with uncertainties that can be quantified via inter-comparison and statistical treatment of these independent measurements. Importantly, this multi-pronged approach will be used on three coral species that due to differences in the carbonate chemistry of their native waters possess differing capacities for proton regulation at their site of calcification; a deep, cold-water coral (strong proton-pumper); a shallow, temperate coral (moderate proton-pumper); and a shallow, tropical coral (weak proton-pumper). Target outcomes of this research include (1) cross-examination of the validity of three independent approaches to estimating coral calcifying fluid pH, (2) quantification of uncertainty associated with the three approaches to estimating coral calcifying fluid pH, (3) advancement of our mechanistic understanding of coral calcification, (4) exploration of the mechanism by which ocean acidification impacts coral calcification, (5) elucidation why corals exhibit such varied responses to ocean acidification, (6) identification of coral types most vulnerable to ocean acidification, (7) exploration of so-called "vital effects" that limit the use of corals in paleoceanographic reconstructions, and (8) quantitative constraint of existing models of coral biomineralization.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1357665</u>
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1437371</u>
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1459706</u>
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1429373</u>

[table of contents | back to top]