

# Dissolved oxygen from 4 field sites in Bogue Sound, North Carolina from 2014 to 2015.

**Website:** <https://www.bco-dmo.org/dataset/721344>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2017-12-14

## Project

» [Microbial Regulation of Greenhouse Gas N2O Emission from Intertidal Oyster Reefs](#) (Oyster Reef N2O Emission)

Contributors	Affiliation	Role
<a href="#">Piehler, Michael F.</a>	University of North Carolina at Chapel Hill (UNC-Chapel Hill-IMS)	Principal Investigator
<a href="#">Brush, Mark J.</a>	Virginia Institute of Marine Science (VIMS)	Co-Principal Investigator
<a href="#">Song, Bongkeun</a>	Virginia Institute of Marine Science (VIMS)	Co-Principal Investigator
<a href="#">Tobias, Craig</a>	University of Connecticut (UConn - Avery Point)	Co-Principal Investigator
<a href="#">Ake, Hannah</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Dissolved oxygen from 4 field sites in Bogue Sound, North Carolina from 2014 to 2015.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** Lat:34.710129 Lon:-76.867882

## Dataset Description

Dissolved oxygen from 4 field sites in Bogue Sound, located in coastal North Carolina.

## Methods & Sampling

Water samples collected on-site. Dissolved oxygen concentrations were measured with Membrane Inlet Mass Spectrometer (MIMS). The 4 field sites are sites of oyster reef and salt marsh restoration. Identified in data as UNC Institute of Marine Sciences (IMS), Carrot Island (Carrot), NOAA Beaufort (NOAA), and Army Marsh (Army). All sites located in Bogue Sound near Morehead City, NC.

## Data Processing Description

Oxygen concentration data corrected using standard curves.

### BCO-DMO Data Processing Notes:

- sorted by season, site

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>O2_conc.csv</b> (Comma Separated Values (.csv), 37.66 KB) MD5:2a9eee3ee49542aaa4234e1182735ab1 Primary data file for dataset ID 721344

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Season	Season data were collected	unitless
Site	Site where data were collected	unitless
CoreName	Core name	unitless
CoreNo	Core ID number	unitless
TimePt	Time point of collection; 1 2 or 3	unitless
O2Flux	Oxygen concentration	milligrams per liter

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	Membrane inlet mass spectrometer
<b>Generic Instrument Name</b>	Membrane Inlet Mass Spectrometer
<b>Dataset-specific Description</b>	Used to measure oxygen concentrations
<b>Generic Instrument Description</b>	Membrane-introduction mass spectrometry (MIMS) is a method of introducing analytes into the mass spectrometer's vacuum chamber via a semipermeable membrane.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Microbial Regulation of Greenhouse Gas N<sub>2</sub>O Emission from Intertidal Oyster Reefs (Oyster Reef N<sub>2</sub>O Emission)

*Extracted from the NSF award abstract:*

Oyster reefs are biogeochemical hot spots and prominent estuarine habitats that provide disproportionate ecological function. Suspension-feeding eastern oysters, *Crassostrea virginica*, are capable of improving water quality and diminishing eutrophication by filtering nutrients and particles from the water and depositing them in the sediments. Remineralization of these deposits may enhance sedimentary denitrification that facilitates nitrogen removal in tidal estuaries. However, the scientific underpinning of oyster reef function has been challenged in various studies. In addition, recent studies of filter feeding invertebrates reported the production of nitrous oxide (N<sub>2</sub>O), a greenhouse gas, as an end product of incomplete denitrification by gut microbes. *C. virginica* could be another source of N<sub>2</sub>O flux from intertidal habitats. Preliminary work indicated substantial N<sub>2</sub>O production from individual oysters. The estimated N<sub>2</sub>O production from high density oyster reefs may exceed the N<sub>2</sub>O flux measured from some estuaries. With the new discovery of N<sub>2</sub>O emission and uncertainty regarding eutrophication control, the ecological value of oyster reef restoration may become equivocal.

This project will quantify N<sub>2</sub>O fluxes to understand the factors controlling N<sub>2</sub>O emission from oyster reefs. Sedimentary N processes will be examined to develop an oyster reef N model to estimate N<sub>2</sub>O emission from tidal creek estuaries relative to other N cycling processes. The PIs hypothesize that intertidal oyster reefs are a substantial source of N<sub>2</sub>O emission from estuarine ecosystems and the magnitude of emission may be linked to water quality. If substantial N<sub>2</sub>O flux from oyster reefs is validated, ecological benefits of oyster reef restoration should be reevaluated. This interdisciplinary research team includes a microbial ecologist, a biogeochemist, an ecologist and an ecosystem modeler. They will utilize stable isotope and molecular microbiological techniques to quantify oyster N<sub>2</sub>O production, elucidate microbial sources of N<sub>2</sub>O emission from oysters and sediments, and estimate seasonal variation of N<sub>2</sub>O fluxes from oyster reefs. Measurements from this study will be integrated into a coupled oyster bioenergetics-sediment biogeochemistry model to compare system level rates of N cycling on oyster reefs as a function of oyster density and water quality. Modeling results will be used to assess the relative trade-offs of oyster restoration associated with N cycling. They expect to deliver the following end products: 1) estimation of annual N<sub>2</sub>O flux from oyster reefs as an additional source of greenhouse gases from estuaries, 2) a better understanding of the environmental and microbial factors influencing N<sub>2</sub>O and N<sub>2</sub> fluxes in tidal estuaries, 3) transformative knowledge for the effect of oyster restoration on water quality enhancement and ecosystem function, 4) direct guidance for oyster restoration projects whose goals include water quality enhancement, and 5) a modeling tool for use in research and restoration planning.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1233327</a>

[ [table of contents](#) | [back to top](#) ]