Species composition from net tows and associated CTD data from a small boat in the Mid-Chesapeake Bay from August to October 2013 (CopesPopDynHypoZone project)

Website: https://www.bco-dmo.org/dataset/723801 Data Type: Cruise Results Version: Version Date: 2018-01-12

Project

» <u>Copepod Population Dynamics in Hypoxic Coastal Waters: Physical and Behavioral Regulation of Resupply</u> <u>and Advective Losses</u> (CopesPopDynHypoZone)

Contributors	Affiliation	Role
<u>Roman, Michael</u> <u>R.</u>	University of Maryland Center for Environmental Science (UMCES/HPL)	Principal Investigator
<u>Pierson, James J.</u>	University of Maryland Center for Environmental Science (UMCES/HPL)	Co-Principal Investigator, Contact
<u>Fitzgerald,</u> <u>Catherine</u>	University of Maryland Center for Environmental Science (UMCES/HPL)	Contact
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- <u>Coverage</u>
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Parameters
- Instruments
- Deployments
- Project Information
- Funding

Coverage

Spatial Extent: N:38.48245 **E**:-76.35638 **S**:38.45012 **W**:-76.43992 **Temporal Extent**: 2013-08-02 - 2013-10-04

Dataset Description

This dataset includes species composition data from plankton net samples and associated CTD information.

Related Datasets:

Acartia tonsa mortality with CTD data: 5 daytrips

Methods & Sampling

This abundance and diversity data was obtained from several ~8 hour small-boat cruises done about every other week from August to October. Vertical tow samples were taken from the mid-bay of the Chesapeake from several stations in a transect, corresponding to the mid-line transect sampled by the two week-long cruises (stations M1-M3, cruises 1301/1302). More stations along the same transect were included in the collection of this data set.

The plankton nets used for sampling were 0.5m in diameter and made of either $200\mu m$ or $64\mu m$ mesh. These were equipped to close with a mechanical messenger trigger and were deployed from the winch arm of the R/V Parker.

A CTD cast was done at each station prior to sampling with the plankton net. Sampling depths were determined based on the location of the pycnocline; the aim was to capture zooplankton below and above the pycnocline. For sampling below the pycnocline, the net was deployed below the pycnocline, towed up to the pycnocline, triggered to close, and returned to the surface. Sampling above the pycnocline was done by standard vertical tow: deploy net to desired depth and winch to the surface.

Once brought on board, the nets were rinsed down with to seawater to collect plankton in the codend. Codends were filtered onto 64µm sieves, then the samples were transferred to glass jars labeled on the lid with the date, time, station, and depth sampled. Sieves with 64µm mesh were selected to catch all life stages of the copepod *Acartia tonsa* since *Acartia tonsa* eggs are about 75µm in diameter and all subsequent life stages are larger. Buffered formalin was added to each jar to preserve the sample in a 4% solution.

After returning from the cruises, samples were stored indoors in climate-controlled laboratory space. To process the samples, the contents of the jars were filtered onto 25µm mesh (to avoid loss of organisms), resuspended, and a subsample taken with a stemple pipette was transferred to a counting wheel where it was checked for density and diluted if necessary, the goal being at least 200 individuals of *Acartia tonsa* present but less than 300.

The sample was then examined for species composition under dissecting microscope with darkfield illumination. All organisms were identified to lowest possible taxonomic level. When species composition analysis was complete, the sample was returned to the original glass jar and returned to storage.

Abundance data were entered into Excel spreadsheets and checked for transcription errors, then imported into MatLab for data analysis.

Data Processing Description

Zooplankton samples were sorted under a stereo dissecting microscope within one year of collection. Subsamples were taken with a pipet such that a minimum of 200 individuals were counted from each sample. Zooplankton were identified to lowest possible taxonomic level, to species where possible for copepods, and copepod adults were sexed.

BCO-DMO Data Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- replaced commas with semicolons
- changed blank cells and NaN's with nd for 'no data'
- converted longitudes to negative degree (west)
- split Net_mesh and Tow into separate columns

[table of contents | back to top]

Data Files

 File

 species_composition.csv(Comma Separated Values (.csv), 115.04 KB)

 MD5:b3192efbc24c03b932f2bbc8d95bf509

 Primary data file for dataset ID 723801

[table of contents | back to top]

Parameters

Parameter	Description	Units
cruise	Chronological cruise number used to refer to each day-long cruise	unitless
date	Gergorian calendar date as recorded in the cruise log formatted as mm/dd/yyyy	unitless
time	Time of sampling in EDT as recorded in the cruise log formatted as hhmm	unitless
EDT_DOY	Day of year calculation using cruise log EDT time	decimal day
Station	Station ID as recorded in the day-cruise log	unitless
Lateral_station	Station ID as given in the Lateral Copepods week-long cruises	unitless
Net_mesh_um	Net mesh size (64 or 200 micron)	microns
Tow	plankton tow cast identifier	unitless
Low_depth	The lowest depth of the range sampled	meters
High_depth	The highest depth of the range sampled	meters
Genus	Genus or least specific identifier of organism in sample	unitless
Species	Species or most specific identifier of organism in sample	unitless
Stage	Life stage of organism in sample	unitless
Num_in_aliquot	Taxon count per aliquot	individuals
Num_per_m3	Taxon abundance	individuals/meter^3 (#/m^3)
Abund_m2	Taxon abundance integrated of the depth column	individuals/meter^2 (#/m^2)
Notes	Notes from the technician made during sampling and sample processing	unitless
Latitude	Station latitude	decimal degrees
Longitude	Station longitude	decimal degrees
Year	Year as recorded by CTD as yyyy	unitless
Month	Month as recorded by CTD as mm	unitless
Day	Day as recorded by CTD as dd	unitless
GMT_Hour	Hour as recorded by CTD; GMT time as hh	unitless
Minute	Minute as recorded by CTD as mm	unitless
Second	Second as recorded by CTD as ss	unitless
GMT_DOY	Day of year calculation using CTD recorded GMT time for each cast	decimal day
prSM	Pressure as recorded by CTD sensor for each cast	decibars
t090C	Temperature recorded by CTD sensor for each cast	degrees Celsius
c0S_m	Condictivity recorded by CTD sensor for each cast	microSiemens/meter (uS/m)
sbeox0V	Raw oxygen measurements as recorded by CTD sensors for each cast	volts
wetStar	Fluorescence measure as recorded by WETStar fluorometer on CTD for each cast	volts?
obs	Optical backscatter as recorded by CTD sensor for each cast	volts?
bat	Beam attenuation as recorded by SBE software	per meter
xmiss	Beam transmission as recorded by SBE software	unitless
sbeox0Mg_L	Dissolved oxygen calculated in SEB post-processing from oxygen data recorded by CTD sensors for each cast	milligrams/liter (mg/L)

sbeox0PS	Dissolved oxygen pressure saturation calculated in SEB post- processing from oxygen data recorded by CTD sensors for each cast	unitless
sigma_t00	Density recorded by CTD sensors for each cast	?
depSM	Depth bins; one bin per half meter and then averaged over the depth range of each net cast	meters
sal00	Salinity recorded by CTD sensor for each cast	Practical Salinity Units (PSU)
nbin	Number of measures per depth bin; averaged per net depth range	unitless
flag	Flag for abberant data	unitless

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	SBE 9plus
Generic Instrument Name	CTD - profiler
Dataset- specific Description	Used for sampling; There were twelve Niskin bottles on the SBE 32 Carousel Water Sampler, deployed from the starboard winch of the RV Sharp, along with the SBE 9plus unit which was attached to the rosette. Attached to an SBE 11plus V2 Deck Unit
Generic Instrument Description	The Conductivity, Temperature, Depth (CTD) unit is an integrated instrument package designed to measure the conductivity, temperature, and pressure (depth) of the water column. The instrument is lowered via cable through the water column. It permits scientists to observe the physical properties in real-time via a conducting cable, which is typically connected to a CTD to a deck unit and computer on a ship. The CTD is often configured with additional optional sensors including fluorometers, transmissometers and/or radiometers. It is often combined with a Rosette of water sampling bottles (e.g. Niskin, GO-FLO) for collecting discrete water samples during the cast. This term applies to profiling CTDs. For fixed CTDs, see https://www.bco-dmo.org/instrument/869934 .

Dataset- specific Instrument Name	Dissecting microscope
Generic Instrument Name	Microscope - Optical
Dataset- specific Description	Used to count live and dead copepods.
Generic Instrument Description	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

Dataset- specific Instrument Name	0.5m plankton net
Generic Instrument Name	Plankton Net
Dataset- specific Description	Used to collect live zooplankton. The plankton net was 0.5m in diameter and made of 64μ m mesh, equipped to close with a mechanical messenger trigger, and was deployed from the winch arm of the R/V Parker.
Generic Instrument Description	A Plankton Net is a generic term for a sampling net that is used to collect plankton. It is used only when detailed instrument documentation is not available.

[table of contents | back to top]

Deployments

HPL_2013-08-02

Website	https://www.bco-dmo.org/deployment/723585
Platform	small boat: UMCES
Start Date	2013-08-02
End Date	2013-08-02
Description	Day trip to collect zooplankton

HPL_2013-08-16

Website	https://www.bco-dmo.org/deployment/723591
Platform	small boat: UMCES
Start Date	2013-08-16
End Date	2013-08-16
Description	Day trip to collect zooplankton.

HPL_2013-09-06

Website	https://www.bco-dmo.org/deployment/723594
Platform	small boat: UMCES
Start Date	2013-09-06
End Date	2013-09-06
Description	Day trip to collect zooplankton

HPL_2013-09-20

Website	https://www.bco-dmo.org/deployment/723602
Platform	small boat: UMCES
Start Date	2013-09-20
End Date	2013-09-20
Description	Day trip to collect zooplankton.

HPL_2013-10-04

Website	https://www.bco-dmo.org/deployment/723607
Platform	small boat: UMCES
Start Date	2013-10-04
End Date	2013-10-04
Description	Day trip to collect zooplankton.

[table of contents | back to top]

Project Information

Copepod Population Dynamics in Hypoxic Coastal Waters: Physical and Behavioral Regulation of Resupply and Advective Losses (CopesPopDynHypoZone)

Coverage: hypoxic zone of Chesapeake Bay

Description from NSF award abstract:

The PIs will develop a mechanistic understanding of how circulation interacts with hypoxia-induced behavioral and physiological changes to affect the population dynamics of coastal zooplankton. They will do this by assessing two potentially contrasting mechanisms influencing the dynamics of the copepod *Acartia tonsa* in the hypoxic zone of Chesapeake Bay. The first hypothesis is that maintenance of copepod populations in the hypoxic region requires replenishment by advection (immigration) of animals through wind-driven lateral transport processes. The second, counteractive, hypothesis is that bottom water hypoxia alters the vertical distribution of *A. tonsa*, thereby making them more susceptible to advective losses from the region (emigration) via surface water transport in the estuarine circulation. They will take advantage of a current NSF-funded physical oceanography research program in Chesapeake Bay that will comprehensively measure and model axial and lateral water exchanges in the mid-Bay region.

The present study will use the physical oceanography study site as a Controlled Volume (CV) in which the oceanographic exchanges of water and the driving mechanisms for those exchanges will be well defined. The PIs will conduct high-resolution spatial and temporal sampling of zooplankton and combine the data with measurements of copepod behavior, mortality and egg production in the hypoxic region. They will use an improved Individual-Based Model of the life history of *A. tonsa* coupled with the circulation to explore the combined effects of advection, behavior, egg production, and mortality on population dynamics. In addition to increasing our knowledge of the impacts of bottom water hypoxia on copepod populations in Chesapeake Bay, the study will improve our general understanding of the regulation of zooplankton populations by physical and biological processes and the impacts of hypoxia on secondary production and food webs in coastal waters.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1259691

[table of contents | back to top]