# Size distribution of Pocillopora colonies used in the experiment at Moorea, French Polynesia from April to June 2008 (CDD\_in\_Reef\_Fish project)

Website: https://www.bco-dmo.org/dataset/726929

**Data Type**: experimental

Version: 1

Version Date: 2017-10-05

#### **Proiect**

» <u>Cryptic density dependence: the effects of spatial, ontogenetic, and individual variation in reef fish</u> (CDD in Reef Fish)

Contributors	Affiliation	Role
Geange, Shane	Victoria University of Wellington	Principal Investigator, Contact
Shima, Jeffrey	Victoria University of Wellington	Co-Principal Investigator
Stier, Adrian	University of Washington (UW)	Co-Principal Investigator
Biddle, Mathew	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

#### **Abstract**

This dataset is from a manipulative experiment investigating the independent and combined effects of priority effects and habitat complexity on the strength of intraspecific competitive interactions among recently settled individuals of a coral reef fish.

# **Table of Contents**

- Coverage
- Dataset Description
  - Methods & Sampling
  - Data Processing Description
- Data Files
- Related Publications
- Parameters
- Deployments
- Project Information
- Funding

# Coverage

**Spatial Extent**: Lat:-17.5 Lon:-149.8333333 **Temporal Extent**: 2008-04 - 2008-06

# **Dataset Description**

This dataset is from a manipulative experiment investigating the independent and combined effects of priority effects and habitat complexity on the strength of intraspecific competitive interactions among recently settled individuals of a coral reef fish (Thalassoma quinquevittatum: Labridae). This research was conducted in 2008 at the Gump Biological Research Station, Moorea, French Polynesia in the northern lagoon (17 30' S, 149 50' W).

This file describes the size distribution of Pocillopora colonies used in the experiment. For additional data, please see files listed in Related Datasets.

#### **Related Datase**

- Geange and Stier 2010 Priority Effects: <a href="https://www.bco-dmo.org/dataset/726890">https://www.bco-dmo.org/dataset/726890</a>
- Geange\_and\_Stier\_2010 Priority Effects Area: <a href="https://www.bco-dmo.org/dataset/726929">https://www.bco-dmo.org/dataset/726929</a> (current page)

- Geange and Stier 2010 Priority Effects Length: <a href="https://www.bco-dmo.org/dataset/727007">https://www.bco-dmo.org/dataset/727007</a>
- Geange\_and\_Stier\_2010 Priority Effects Background Community: <a href="https://www.bco-dmo.org/dataset/726945">https://www.bco-dmo.org/dataset/726945</a>

# Methods & Sampling

This dataset describes the size distribution of *Pocillopora* colonies used in the experiment.

We examined the effects of habitat complexity and timing of arrival in the context of the survival of focal Thalassoma quinquevittatum settlers. The study was conducted in the northern lagoon of Moorea between April and June 2008. We used an array of 30 isolated live-coral patch reefs separated by \*10 m in water 2-4 m deep. Reefs were located within a sand-flat, separated from each other, and from nearby natural reefs, by a minimum of 15 m. We constructed reefs to minimize habitat variation by standardizing size, rugosity, and water depth. Each reef consisted of a base of live *Porites lobata* coral with an average area of 2.23 m<sup>2</sup> (SD = 0.56), and a mean height of 0.59 m (SD = 0.10). We controlled habitat complexity by manipulating the availability of the branching coral Pocillopora verrucosa. This was achieved by drilling holes into the upper surface of patch reefs. Into these holes, we inserted stainless steel pins attached to P. verrucosa colonies with Z-Spar Splash Zone Compound (Kopcoat, Pittsburgh, PA, USA). Mean colony surface area was 0.2 m<sup>2</sup> (SD = 0.07). We crossed the availability of *P. verrucosa* (two levels: two, or four colonies) with the presence of three tagged T. auinquevittatum competitors (three levels; absent, introduced simultaneously with (0 days), or 5 days earlier than the focal individuals). To each reef, we simulated settlement by introducing three tagged T. quinquevittatum focal individuals. Thus, our design had six treatments: (1) focal individuals without competitors, with two P. verrucosa colonies; (2) focal individuals and competitors introduced simultaneously, with two P. verrucosa colonies; (3) focal individuals with competitors introduced 5 days previously, with two P. verrucosa colonies; (4) focal individuals without competitors, with four P. verrucosa colonies; (5) focal individuals and competitors introduced simultaneously, with four P. verrucosa colonies; and (6) focal individuals with competitors introduced 5 days previously, with four P. verrucosa colonies. We ran the experiment in two temporal blocks (17-23 April and 1-7 May 2008), yielding ten replicates (five in each temporal block) for each of the six treatments. We surveyed reefs twice daily (approximately 0800 and 1600 hours) for 5 days after the introduction of focal individuals.

# **Data Processing Description**

poc\_area\_m2 was calculated as the total surface area of a rectangle using over\_top\_length, over\_top\_width, and max height.

## **BCO-DMO Processing:**

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions

[ table of contents | back to top ]

#### **Data Files**

#### File

**GeangeandStier\_2010\_PriorityEffects\_Area.csv**(Comma Separated Values (.csv), 2.13 KB)

MD5:9f69246d739678388d6eda5b7307ff21

Primary data file for dataset ID 726929

[ table of contents | back to top ]

## **Related Publications**

Geange, S. W., & Stier, A. C. (2010). Priority effects and habitat complexity affect the strength of competition.

# [ table of contents | back to top ]

### **Parameters**

Parameter	Description	Units
colony	unique identifier for each Pocillopora colony	unitless
over_top_length	distance in meters of the over the top length of the Pocillopora colony	meters (m)
over_top_width	distance in meters of the over the top width of the Pocillopora colony	meters (m)
max_height	distance in meters of the maximum height of the Pocillopora colony	meters (m)
poc_area_m2	surface area of the Pocillopora colony in square meters (calculated as the total surface area of a rectangle using over_top_length over_top_width and max_height.)	

[ table of contents | back to top ]

# **Deployments**

Osenberg\_et\_al\_Moorea

Website	Website https://www.bco-dmo.org/deployment/64475	
Platform	Osenberg et al Moorea	
Start Date	2003-05-19	
End Date	2015-07-12	

[ table of contents | back to top ]

# **Project Information**

Cryptic density dependence: the effects of spatial, ontogenetic, and individual variation in reef fish (CDD\_in\_Reef\_Fish)

Coverage: Moorea, French Polynesia (-17.48, -149.82)

# Description from NSF award abstract:

Ecologists have long been interested in the factors that drive spatial and temporal variability in population density and structure. In marine reef systems, attention has focused on the role of settlement-the transition of pelagic larvae to a benthic stage-and on density-dependent processes affecting recently settled juveniles. Recent data suggest that co-variance in settlement and subsequent density-dependent survival can obscure the patterns of density dependence at larger scales, a phenomenon called cryptic density dependence. This research will explore the mechanisms that underlie the spatial covariance of settlement and site quality - a process that has received little attention in the standard paradigm. These mechanistic studies of cryptic density dependence will facilitate the development of new frameworks for fish population dynamics that incorporate larval ecology, habitat quality, density dependence, life history, and the patterns and implications of spatial covariance among these factors. More generally, the work provides a specific empirical context, and a general theoretical treatment, of cryptic heterogeneity (hidden individual variation in demographic rates).

**Note:** Drs. Craig W. Osenberg and Ben Bolker were at the University of Florida at the time the NSF award was granted. Dr. Osenberg moved to the University of Georgia during the summer of 2014 (<u>current contact information</u>). Dr. Bolker moved to McMaster University in 2010 (<u>current contact information</u>).

# [ table of contents | back to top ]

# **Funding**

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0242312

[ table of contents | back to top ]