

# Lionfish and grouper density from surveys conducted at Lee Stocking Island, Bahamas from 2009-2015 (BiodiversityLossEffects\_lionfish project)

**Website:** <https://www.bco-dmo.org/dataset/728810>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2018-04-04

## Project

» [Mechanisms and Consequences of Fish Biodiversity Loss on Atlantic Coral Reefs Caused by Invasive Pacific Lionfish](#) (BiodiversityLossEffects\_lionfish)

Contributors	Affiliation	Role
<a href="#">Hixon, Mark</a>	University of Hawaii at Manoa (HIMB)	Principal Investigator
<a href="#">Benkwitt, Cassandra E.</a>	Oregon State University (OSU)	Contact
<a href="#">York, Amber D.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** Lat:23.7667 Lon:-76.099167

**Temporal Extent:** 2009 - 2015

## Dataset Description

Survey data for invasive lionfish and a comparison native predator (Nassau Grouper) on reefs around Lee Stocking Island, The Bahamas (23°46'0.12"N 76° 5'57.00"W). These data are compiled across multiple spatial scales and were collected from the time lionfish first arrived in the study area (2009) through 2015.

Species names and common names for the species codes used in this dataset can be found in the "Species Key" dataset <https://www.bco-dmo.org/dataset/655195>.

These data were utilized in the following publication (Benkwitt et al., 2017)

## Methods & Sampling

To determine patterns of lionfish (*Pterois volitans*) and Nassau Grouper (*Epinephelus striatus*) abundance and body size over the first decade of the lionfish invasion, we examined fish survey data collected in the Exuma Cays, The Bahamas, from when lionfish first arrived in the study area in 2005 through 2015. We surveyed reef-resident fishes at three spatial scales: (1) small patch reefs; (2) medium-sized patch reefs; and (3) large reefs. Small reefs included 32 coral patch reefs (ca. 6 m<sup>2</sup>) as well as 16 artificial reefs (ca. 1 m<sup>2</sup>) constructed

of concrete blocks. Surveys were conducted annually on these reefs from June to August during 2005–2015, with the exception of 2006 and 2012–2014. A pair of trained observers recorded the abundance and visually estimated body size (to nearest cm total length, TL) of all resident fishes on each reef. Censuses consisted of divers slowly circling the reefs at distances of ~3, ~1, and 0 m and then using flashlights to thoroughly search all holes and crevices. Medium-sized reefs comprised six natural coral patches (ca. 6 to 23 m<sup>2</sup>). Fish populations on these reefs were censused in 2006–2012 and 2015 following the same protocol as for the small reefs. At the largest scale, ten reefs (ca. 1400 to 4000 m<sup>2</sup>) were surveyed as part of a long-term lionfish manipulation experiment. All resident fishes in 400 m<sup>2</sup> survey areas (two 10 x 10 m plots and four 2 x 25 m transects) were censused on each reef. We examined fish surveys that were conducted in June 2009, before any lionfish manipulations, to surveys conducted in July/August 2015, 3 years after the experiment of Albins (2015) was completed. Lionfish were never manipulated on one of these large reefs, which was isolated from other reefs. We examined a complete time series of surveys on this particular reef, including censuses conducted in both winter (November–February) and summer (June–August) of 2009, 2010, 2011, and in summer of 2012 and 2015.

Additional methods for surveys of large reefs included in: Albins, MA. 2015.

## Data Processing Description

Several years are missing data, due to logistical constraints that prevented surveys. In 2015, only 3 out of 6 subsamples of large reefs were surveyed due to time constraints.

BCO-DMO Data Manager Processing Notes:

- \* added a conventional header with dataset name, PI name, version date
- \* modified parameter names to conform with BCO-DMO naming conventions
- \* changed column names 5,10,15,..65 to len\_5,len\_10..len\_15 etc. Data parameters in our system can't start with a number.
- \* column "NOTES" removed as it has nothing in it.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Lionfish_and_grouper_density.csv</b> (Comma Separated Values (.csv), 56.39 KB) MD5:1687e492617792bc351e9891b072ad62
Primary data file for dataset ID 728810

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Albins, M. (2015). Invasive Pacific lionfish *Pterois volitans* reduce abundance and species richness of native Bahamian coral-reef fishes. *Marine Ecology Progress Series*, 522, 231–243. doi:[10.3354/meps11159](https://doi.org/10.3354/meps11159)  
*Methods*

Benkwitt, C. E., Albins, M. A., Buch, K. L., Ingeman, K. E., Kindinger, T. L., Pusack, T. J., ... Hixon, M. A. (2017). Is the lionfish invasion waning? Evidence from The Bahamas. *Coral Reefs*, 36(4), 1255–1261. doi:[10.1007/s00338-017-1620-7](https://doi.org/10.1007/s00338-017-1620-7)  
*Results*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
YEAR	Sampling year in format yyyy	unitless
MONTH	Sampling month in format mm	unitless
DAY	Sampling day in format dd	unitless
OBSERVER	Initials of primary observer conducting census	unitless
REEF	Name of reef surveyed	unitless
SUBSAMPLE	For large reefs, number or letter denoting permanent subsample (1-2 = 10 x 10 m plot, A-D = 2 x 25 m transect). For all other reefs, the entire reef was surveyed so subsample is not applicable (n/a)	unitless
TYPE	Type of reef, either ART = artificial (cinderblocks), NAT = natural, or TRANS = transplanted coral heads	unitless
SCALE	Size of reef, either small (< 6 m <sup>2</sup> ), medium (6 - 23 m <sup>2</sup> ), or large (1400-4000 m <sup>2</sup> )	unitless
SPECIES	Either EPST = <i>Epiniphalus striatus</i> (Nassau Grouper) or PTVO = <i>Pterois volitans</i> (Red lionfish)	unitless
len_5	Number of individuals that were between 0-5 cm total length (visually estimated)	unitless
len_10	Number of individuals that were between 5.1-10 cm total length (visually estimated)	unitless
len_15	Number of individuals that were between 10.1-15 cm total length (visually estimated)	unitless
len_20	Number of individuals that were between 15.1-20 cm total length (visually estimated)	unitless
len_25	Number of individuals that were between 20.1-25 cm total length (visually estimated)	unitless
len_30	Number of individuals that were between 25.1-30 cm total length (visually estimated)	unitless
len_35	Number of individuals that were between 30.1-35 cm total length (visually estimated)	unitless
len_40	Number of individuals that were between 35.1-40 cm total length (visually estimated)	unitless
len_45	Number of individuals that were between 40.1-45 cm total length (visually estimated)	unitless
len_50	Number of individuals that were between 45.1-50 cm total length (visually estimated)	unitless
len_55	Number of individuals that were between 50.1-55 cm total length (visually estimated)	unitless
len_60	Number of individuals that were between 55.1-60 cm total length (visually estimated)	unitless
len_65	Number of individuals that were greater than 65 cm total length (visually estimated)	unitless
TOTAL	Total number of individuals (summed across all size classes)	unitless

[ [table of contents](#) | [back to top](#) ]

## Deployments

### PIMS\_Hixon

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/59038">https://www.bco-dmo.org/deployment/59038</a>
<b>Platform</b>	Tropical Marine Lab at Lee Stocking Island
<b>Start Date</b>	2009-05-30
<b>End Date</b>	2012-08-18
<b>Description</b>	Various lab experiments were conducted between 2009 and 2012 at the facilities at the Perry Institute for Marine Science Tropical Marine Lab (at Lee Stocking Island, Bahamas) for the project "Ecological Release and Resistance at Sea: Invasion of Atlantic Coral Reefs by Pacific Lionfish".

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Mechanisms and Consequences of Fish Biodiversity Loss on Atlantic Coral Reefs Caused by Invasive Pacific Lionfish (BiodiversityLossEffects\_lionfish)

**Website:** <http://hixon.science.oregonstate.edu/content/highlight-lionfish-invasion>

**Coverage:** Three Bahamian sites: 24.8318, -076.3299; 23.8562, -076.2250; 23.7727, -076.1071; Caribbean Netherlands: 12.1599, -068.2820

The Pacific red lionfish (*Pterois volitans*), a popular aquarium fish, was introduced to the Atlantic Ocean in the vicinity of Florida in the late 20th century. Voraciously consuming small native coral-reef fishes, including the juveniles of fisheries and ecologically important species, the invader has undergone a population explosion that now ranges from the U.S. southeastern seaboard to the Gulf of Mexico and across the greater Caribbean region. The PI's past research determined that invasive lionfish (1) have escaped their natural enemies in the Pacific (lionfish are much less abundant in their native range); (2) are not yet controlled by Atlantic predators, competitors, or parasites; (3) have strong negative effects on populations of native Atlantic fishes; and (4) locally reduce the diversity (number of species) of native fishes. The lionfish invasion has been recognized as one of the major conservation threats worldwide.

The Bahamas support the highest abundances of invasive lionfish globally. This system thus provides an unprecedented opportunity to understand the direct and indirect effects of a major invader on a diverse community, as well as the underlying causative mechanisms. The PI will focus on five related questions: (1) How does long-term predation by lionfish alter the structure of native reef-fish communities? (2) How does lionfish predation destabilize native prey population dynamics, possibly causing local extinctions? (3) Is there a lionfish-herbivore-seaweed trophic cascade on invaded reefs? (4) How do lionfish modify cleaning mutualisms on invaded reefs? (5) Are lionfish reaching densities where natural population limits are evident?

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1233027</a>

[ [table of contents](#) | [back to top](#) ]