

Summary of pCO₂ and temperature treatment combinations for each culture pot and experimental trial (Seaweed OA Resilience project)

Website: <https://www.bco-dmo.org/dataset/731452>

Data Type: experimental

Version: 1

Version Date: 2018-03-23

Project

» [Ocean Acidification: Scope for Resilience to Ocean Acidification in Macroalgae](#) (Seaweed OA Resilience)

Program

» [Science, Engineering and Education for Sustainability NSF-Wide Investment \(SEES\): Ocean Acidification \(formerly CRI-OA\)](#) (SEES-OA)

Contributors	Affiliation	Role
Kubler, Janet E.	California State University Northridge (CSUN)	Principal Investigator
Dudgeon, Steve	California State University Northridge (CSUN)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Summary of pCO₂ and temperature treatment combinations for each culture pot and experimental trial.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Supplemental Files](#)
 - [Parameters](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:34 E:-117 S:33 W:-118

Temporal Extent: 2014-06-04 - 2015-02-23

Dataset Description

This dataset includes a summary of the experimental pCO₂ and temperature treatments for *Plocamium cartilagineum* cultures.

Related Datasets:

[Plocamium carbon nitrogen and stable isotopes](#): Plocamium carbon and nitrogen content and stable isotope values, 2014-2015 (Seaweed OA Resilience project)

[Plocamium culture carbonate chemistry](#): Carbonate chemistry in experimental cultures of Plocamium cartilagineum cultured at different temperatures and pCO₂ levels (Seaweed OA Resilience project)

[Plocamium culture: seawater delta13C](#): Stable isotope ratio and concentration of carbon in seawater during Plocamium culture experiments, 2014-2015 (Seaweed OA Resilience project)

[Plocamium cultures pH and temperature](#): Plocamium culture pot pH and temperature time-series at 10 minute sampling intervals from 2014-2015 (Seaweed OA Resilience project)

[Plocamium growth and biomass](#): Experimental results of Plocamium cartilagineum growth and biomass as a function of pCO₂ and temperature (Seaweed OA Resilience project)

[Plocamium pigments](#): Photosynthetic pigment concentrations in Plocamium cartilagineum, trials 3-8, 2014-2015 (Seaweed OA Resilience project)

[Plocamium: pH drift](#): Carbonate chemistry over a time-course in pH drift experiments with Plocamium growth collected at Catalina Island, 2014-2015 (Seaweed OA Resilience project)

[Rapid Light Curves_PAM](#): Measurements of fluorescence of photosystem II in Plocamium cartilagineum under various and pCO₂ and temperature conditions

Methods & Sampling

Plocamium cartilagineum was collected from Catalina Island in June - Nov. 2014 and Jan. 2015.

Culture pots were placed in large thermally insulated coolers in a temperature-controlled water bath at either 15 or 20°C under saturating illumination of ~150 micro-moles photons/m²/s. pCO₂ treatments were supplied to closed culture pots by use of a gas mixing system combining Nitrogen, Oxygen and Carbon Dioxide to specific CO₂ partial pressures, 20.9% oxygen and the balance being Nitrogen.

Trial 1 was a pilot test of culture system and methodological procedures so was not used for data collection in the testing of hypotheses. In each culture pot, pCO₂ was set by the supply rate of CO₂ in the corresponding mass-flow controlled gas mixing system to be within a target range of either near ambient, moderately elevated or highly elevated in each trial. The near-ambient range was narrower than the other target ranges (set points typically ~380 - 390 µatm) as it served as the control range in each trial. However, actual pCO₂ in solution in each culture pot varied slightly from constant target values on a diurnal cycle associated with the metabolic activities of the algae contained within each pot. Unique average values of pCO₂ in each culture pot based on different set values for each mass-flow controlled mixer within the qualitative ranges of ambient, moderate and highly increased pCO₂ levels that were replicated in each trial of the experiment enabled a more powerful regression-type experimental design. With a regression type approach, we could estimate the functional relationship between response variables and pCO₂, which was not possible with a simple categorical treatment design.

See Supplementary File for a table of culture conditions for each of the 8 trials (pdf).

Data Processing Description

BCO-DMO Processing Notes:

- added a conventional header with dataset name and description, PI names, version date

[[table of contents](#) | [back to top](#)]

Data Files

File
CO2trt_x_pot_trial.csv (Comma Separated Values (.csv), 1.65 KB) MD5:3b3032d6c914bc5e87e5d1b3973fc7ce
Primary data file for dataset ID 731452

[[table of contents](#) | [back to top](#)]

Supplemental Files

File

Table of culture conditions for each of the 8 trials

filename: Dudgeon_seaweed_trial_meta_2018-01.pdf

(Portable Document Format (.pdf), 429.04 KB)
MD5:5a4a9cbd3cb5092a419b4883c96979b8

Notes: Trial 1 was a pilot test of culture system and methodological procedures so was not used for data collection in the testing of hypotheses. In each culture pot, pCO₂ was set by the supply rate of CO₂ in the corresponding mass-flow controlled gas mixing system to be within a target range of either near ambient, moderately elevated or highly elevated in each trial. The near-ambient range was narrower than the other target ranges (set points typically ~380 - 390 micro-atm) as it served as the control range in each trial. However, actual pCO₂ in solution in each culture pot varied slightly from constant target values on a diurnal cycle associated with the metabolic activities of the algae contained within each pot. Unique average values of pCO₂ in each culture pot based on different set values for each mass-flow controlled mixer within the qualitative ranges of ambient, moderate and highly increased pCO₂ levels that were replicated in each trial of the experiment enabled a more powerful regression-type experimental design. With a regression type approach we could estimate the functional relationship between response variables and pCO₂, which was not possible with a simple categorical treatment design.

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
pot	culture pot identifier	unitless
pCO ₂ _target_range	categorical descriptor of 3 target ranges in each trial	unitless
pCO ₂ _avg_meas	measured average partial pressure of CO ₂	microatmospheres (uatm)
Temp	temperature	degrees Celsius
Trial	experimental trial	unitless

[[table of contents](#) | [back to top](#)]

Project Information

Ocean Acidification: Scope for Resilience to Ocean Acidification in Macroalgae (Seaweed OA Resilience)

Coverage: Temperate coastal waters of the USA (30 - 45 N latitude, -66 to -88 W and -117 to -125 W longitude)

Benthic macroalgae contribute to intensely productive near shore ecosystems and little is known about the potential effects of ocean acidification on non-calcifying macroalgae. Kübler and Dudgeon will test hypotheses about two macroalgae, *Ulva* spp. and *Plocamium cartilagineum*, which, for different reasons, are hypothesized to be more productive and undergo ecological expansions under predicted changes in ocean chemistry. They have designed laboratory culture-based experiments to quantify the scope for response to ocean acidification in *Plocamium*, which relies solely on diffusive uptake of CO₂, and populations of *Ulva* spp., which have an inducible concentrating mechanism (CCM). The investigators will culture these algae in media equilibrated at 8 different pCO₂ levels ranging from 380 to 940 ppm to address three key hypotheses. The first is that macroalgae (such as *Plocamium cartilagineum*) that are not able to acquire inorganic carbon in changed form will benefit, in terms of photosynthetic and growth rates, from ocean acidification. There is little existing data to support this common assumption. The second hypothesis is that enhanced growth of *Ulva* sp. under OA will result from the energetic savings from down regulating the CCM, rather than from enhanced photosynthesis per se. Their approach will detect existing genetic variation for adaptive plasticity. The third key hypothesis to be addressed in short-term culture experiments is that there will be a significant interaction between ocean acidification and nitrogen limited growth of *Ulva* spp., which are indicator species of eutrophication. Kübler and Dudgeon will be able to quantify the individual effects of ocean acidification and nitrogenous nutrient addition on *Ulva* spp. and also, the synergistic effects, which will inevitably apply in many highly productive, shallow coastal areas. The three hypotheses being addressed have been broadly identified as urgent needs in our growing understanding of the impacts of ocean acidification.

Program Information

Science, Engineering and Education for Sustainability NSF-Wide Investment (SEES): Ocean Acidification (formerly CRI-OA) (SEES-OA)

Website: https://www.nsf.gov/funding/pgm_summ.jsp?pims_id=503477

Coverage: global

NSF Climate Research Investment (CRI) activities that were initiated in 2010 are now included under Science, Engineering and Education for Sustainability NSF-Wide Investment (SEES). SEES is a portfolio of activities that highlights NSF's unique role in helping society address the challenge(s) of achieving sustainability. Detailed information about the SEES program is available from NSF (https://www.nsf.gov/funding/pgm_summ.jsp?pims_id=504707).

In recognition of the need for basic research concerning the nature, extent and impact of ocean acidification on oceanic environments in the past, present and future, the goal of the SEES: OA program is to understand (a) the chemistry and physical chemistry of ocean acidification; (b) how ocean acidification interacts with processes at the organismal level; and (c) how the earth system history informs our understanding of the effects of ocean acidification on the present day and future ocean.

Solicitations issued under this program:

[NSF 10-530](#), FY 2010-FY2011

[NSF 12-500](#), FY 2012

[NSF 12-600](#), FY 2013

[NSF 13-586](#), FY 2014

NSF 13-586 was the final solicitation that will be released for this program.

PI Meetings:

[1st U.S. Ocean Acidification PI Meeting](#) (March 22-24, 2011, Woods Hole, MA)

[2nd U.S. Ocean Acidification PI Meeting](#) (Sept. 18-20, 2013, Washington, DC)

3rd U.S. Ocean Acidification PI Meeting (June 9-11, 2015, Woods Hole, MA – Tentative)

NSF media releases for the Ocean Acidification Program:

[Press Release 10-186 NSF Awards Grants to Study Effects of Ocean Acidification](#)

[Discovery Blue Mussels "Hang On" Along Rocky Shores: For How Long?](#)

[Discovery nsf.gov - National Science Foundation \(NSF\) Discoveries - Trouble in Paradise: Ocean Acidification This Way Comes - US National Science Foundation \(NSF\)](#)

[Press Release 12-179 nsf.gov - National Science Foundation \(NSF\) News - Ocean Acidification: Finding New Answers Through National Science Foundation Research Grants - US National Science Foundation \(NSF\)](#)

[Press Release 13-102 World Oceans Month Brings Mixed News for Oysters](#)

[Press Release 13-108 nsf.gov - National Science Foundation \(NSF\) News - Natural Underwater Springs Show How Coral Reefs Respond to Ocean Acidification - US National Science Foundation \(NSF\)](#)

[Press Release 13-148 Ocean acidification: Making new discoveries through National Science Foundation research grants](#)

[Press Release 13-148 - Video nsf.gov - News - Video - NSF Ocean Sciences Division Director David Conover answers questions about ocean acidification. - US National Science Foundation \(NSF\)](#)

[Press Release 14-010 nsf.gov - National Science Foundation \(NSF\) News - Palau's coral reefs surprisingly resistant to ocean acidification - US National Science Foundation \(NSF\)](#)

[Press Release 14-116 nsf.gov - National Science Foundation \(NSF\) News - Ocean Acidification: NSF awards](#)

[\\$11.4 million in new grants to study effects on marine ecosystems - US National Science Foundation \(NSF\)](#)

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1316198

[[table of contents](#) | [back to top](#)]