

[DEPRECATED] Dissolved mercury measurements from the Western Arctic Ocean, which were sampled as part of the 2015 US GEOTRACES Cruise (GN01; HLY1502)

Website: <https://www.bco-dmo.org/dataset/738136>

Data Type: Cruise Results

Version: 2

Version Date: 2018-06-22

Project

» [U.S. Arctic GEOTRACES Study \(GN01\)](#) (U.S. GEOTRACES Arctic)

» [Collaborative Research: GEOTRACES Arctic Section: Mercury Speciation and Cycling in the Arctic Ocean](#) (GEOTRACES Arctic Mercury)

Program

» [U.S. GEOTRACES](#) (U.S. GEOTRACES)

Contributors	Affiliation	Role
Lamborg, Carl	University of California-Santa Cruz (UCSC)	Principal Investigator
Hammerschmidt, Chad	Wright State University	Co-Principal Investigator
Agather, Alison	Wright State University	Contact
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:89.995 E:11.9102 S:60.2579 W:-179.5926

Temporal Extent: 2015-08-03 - 2015-10-08

Dataset Description

This dataset includes dissolved mercury measurements from the Western Arctic Ocean, which were sampled as part of the 2015 US GEOTRACES Cruise (GN01; HLY1502). This dataset contains four species of dissolved mercury: total mercury, elemental mercury, monomethylmercury, and dimethylmercury.

Status note (2020-10-28): This dataset has been replaced. Use the following dataset instead: <https://www.bco-dmo.org/dataset/827530>

Methods & Sampling

Samples were collected according to Cutter and Bruland 2012, utilizing a trace metal clean rosette (GT-C rosette). Samples above 20 m were pumped from a hole drilled in the ice far away from the ship, using clean sampling techniques.

Sample bottles were cleaned according to Hammerschmidt et al. 2012.

Dissolved mercury analyses were performed at sea, with the exception of monomethylmercury. The full methodology may be found in the Hg Cook Book, and more detailed method descriptions can be found in Lamborg et al. 2012, Hammerschmidt and Bowman 2012, Bowman et al. 2015, Bowman et al. 2016.

All dissolved species of mercury were measured with a Tekran Model 2500 CVAFS Mercury Detector. Analysis of total mercury (HgT; Fitzgerald and Gill 1979, Bloom and Fitzgerald 1988, Bloom 1989) has a method detection limit of 0.10 pM, and procedural duplicates varied by $11 \pm 5\%$ ($n = 17$).

Elemental mercury (Hg0) was analyzed according to Fitzgerald and Gill (1979), Bloom and Fitzgerald (1988) and Bloom (1989). The method detection limit was 0.04 pM and procedural duplicates varied by $11 \pm 14\%$ ($n = 8$).

Monomethylmercury (MMHg) was analyzed according to Bowman and Hammerschmidt (2011), Tseng et al. (2004) and Munson et al. (2014). The method detection limit was 0.020 pM, and procedural duplicates averaged $11 \pm 8\%$ ($n = 5$).

Dimethylmercury (DMHg) was analyzed according to Bowman and Hammerschmidt (2011), Tseng et al. (2004) and Baya et al. (2013). The method detection limit was 0.012 pM.

Data Processing Description

WOCE Quality Flags:

- 2 - Ok,
- 3 - Questionable,
- 4 - Bad Analysis,
- 5 - Sample lost,
- 6 - Mean of replicates,
- 9 - Sample was not collected for Hg analysis.

"nd" indicates no data.

"bdl" indicates below detection limit.

BCO-DMO Processing:

- separated cruise column into two: cruise_name and cruise_id;
- modified parameter names; applied GEOTRACES names provided by submitter;
- joined the following columns from HLY1502 event log, ODF bottle file, and GT-C bottle file, based on the GEOTRC_SAMPNO: GETORC_EVENTNO, GEOTRC_INSTR, ISO_DATETIME_UTC_START_EVENT, ISO_DATETIME_UTC_END_EVENT, EVENT_LAT, EVENT_LON, BTLNBR, BTLNBR_FLAG_W, BTL_DATE, BTL_TIME, BTL_LAT, BTL_LON, CTDPRS, CTDDEPTH, ODF_CTDPRS, ODF_CTDDEPTH.
- 30-October-2018: removed embargo on dataset.

[[table of contents](#) | [back to top](#)]

Related Publications

Baya, P. A., Hollinsworth, J. L., & Hintelmann, H. (2013). Evaluation and optimization of solid adsorbents for the sampling of gaseous methylated mercury species. *Analytica Chimica Acta*, 786, 61–69.

doi:[10.1016/j.aca.2013.05.019](https://doi.org/10.1016/j.aca.2013.05.019)

Methods

Bloom, N. (1989). Determination of Picogram Levels of Methylmercury by Aqueous Phase Ethylation, Followed by Cryogenic Gas Chromatography with Cold Vapour Atomic Fluorescence Detection. *Canadian Journal of Fisheries and Aquatic Sciences*, 46(7), 1131–1140. doi:[10.1139/f89-147](https://doi.org/10.1139/f89-147)

Methods

Bloom, N., & Fitzgerald, W. F. (1988). Determination of volatile mercury species at the picogram level by low-temperature gas chromatography with cold-vapour atomic fluorescence detection. *Analytica Chimica Acta*, 208, 151-161. doi:10.1016/S0003-2670(00)80743-6 [https://doi.org/10.1016/S0003-2670\(00\)80743-6](https://doi.org/10.1016/S0003-2670(00)80743-6)
Methods

Bowman, K. L., & Hammerschmidt, C. R. (2011). Extraction of monomethylmercury from seawater for low-femtomolar determination. *Limnology and Oceanography: Methods*, 9(4), 121-128. doi:[10.4319/lom.2011.9.121](https://doi.org/10.4319/lom.2011.9.121)
Methods

Bowman, K. L., Hammerschmidt, C. R., Lamborg, C. H., & Swarr, G. (2015). Mercury in the North Atlantic Ocean: The U.S. GEOTRACES zonal and meridional sections. *Deep Sea Research Part II: Topical Studies in Oceanography*, 116, 251-261. doi:[10.1016/j.dsr2.2014.07.004](https://doi.org/10.1016/j.dsr2.2014.07.004)
Methods

Bowman, K. L., Hammerschmidt, C. R., Lamborg, C. H., Swarr, G. J., & Agather, A. M. (2016). Distribution of mercury species across a zonal section of the eastern tropical South Pacific Ocean (U.S. GEOTRACES GP16). *Marine Chemistry*, 186, 156-166. doi:[10.1016/j.marchem.2016.09.005](https://doi.org/10.1016/j.marchem.2016.09.005)
Methods

Cutter, G. A., & Bruland, K. W. (2012). Rapid and noncontaminating sampling system for trace elements in global ocean surveys. *Limnology and Oceanography: Methods*, 10(6), 425-436. doi:[10.4319/lom.2012.10.425](https://doi.org/10.4319/lom.2012.10.425)
Methods

Fitzgerald, W. F., & Gill, G. A. (1979). Subnanogram determination of mercury by two-stage gold amalgamation and gas phase detection applied to atmospheric analysis. *Analytical Chemistry*, 51(11), 1714-1720. doi:[10.1021/ac50047a030](https://doi.org/10.1021/ac50047a030)
Methods

Hammerschmidt, C. R., & Bowman, K. L. (2012). Vertical methylmercury distribution in the subtropical North Pacific Ocean. *Marine Chemistry*, 132-133, 77-82. doi:[10.1016/j.marchem.2012.02.005](https://doi.org/10.1016/j.marchem.2012.02.005)
Methods

Hammerschmidt, C. R., Bowman, K. L., Tabatchnick, M. D., & Lamborg, C. H. (2011). Storage bottle material and cleaning for determination of total mercury in seawater. *Limnology and Oceanography: Methods*, 9(10), 426-431. doi:10.4319/lom.2011.9.426 <https://doi.org/10.4319/lom.2011.9.426>
Methods

Lamborg, C. H., Hammerschmidt, C. R., Gill, G. A., Mason, R. P., & Gichuki, S. (2012). An intercomparison of procedures for the determination of total mercury in seawater and recommendations regarding mercury speciation during GEOTRACES cruises. *Limnology and Oceanography: Methods*, 10(2), 90-100. doi:[10.4319/lom.2012.10.90](https://doi.org/10.4319/lom.2012.10.90)
Methods

Munson, K. M., Babi, D., & Lamborg, C. H. (2014). Determination of monomethylmercury from seawater with ascorbic acid-assisted direct ethylation. *Limnology and Oceanography: Methods*, 12(1), 1-9. doi:[10.4319/lom.2014.12.1](https://doi.org/10.4319/lom.2014.12.1)
Methods

Tseng, C.-M., Hammerschmidt, C. R., & Fitzgerald, W. F. (2004). Determination of Methylmercury in Environmental Matrixes by On-Line Flow Injection and Atomic Fluorescence Spectrometry. *Analytical Chemistry*, 76(23), 7131-7136. doi:[10.1021/ac049118e](https://doi.org/10.1021/ac049118e)
Methods

[[table of contents](#) | [back to top](#)]

Related Datasets

Replaced by New Version

Lamborg, C., Hammerschmidt, C. (2021) **Concentrations of dissolved and particulate mercury species collected on the US GEOTRACES Arctic cruise (HLY1502, GN01) from August to October 2015.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 2) Version Date 2021-02-03 doi:10.26008/1912/bco-dmo.827530.2 [[view at BCO-DMO](#)]

Parameters

Parameter	Description	Units
cruise_name	Cruise identification	unitless
cruise_id	Official cruise ID	unitless
STNNBR	GEOTRACES station number	unitless
Latitude	Latitude of sample collection; provided by the data contributor	decimal degrees north
Longitude	Longitude of sample collection; provided by the data contributor	decimal degrees east
GEOTRC_EVENTNO	GEOTRACES event number; joined from event log	unitless
GEOTRC_INSTR	Sampling instrument; joined from event log	unitless
ISO_DATETIME_UTC_START_EVENT	Date and time, formatted to the ISO 8601 standard, at the start of the sampling event, according to the event log. Format: YYYY-MM-DDTHH:MM:SS[.xx]Z	unitless
ISO_DATETIME_UTC_END_EVENT	Date and time, formatted to the ISO 8601 standard, at the end of the sampling event, according to the event log. Format: YYYY-MM-DDTHH:MM:SS[.xx]Z	unitless
EVENT_LAT	Latitude at the start of the event; north is positive; joined from event log.	decimal degrees
EVENT_LON	Longitude at the start of the event; east is positive; joined from event log.	decimal degrees
GEOTRC_SAMPNO	GEOTRACES sample number	unitless
BTLNBR	Bottle number; joined from bottle file	unitless
BTLNBR_FLAG_W	Bottle number quality flag; follows WOCE conventions. 2 = no problems noted; 3 = leaking; 4 = did not trip correctly; 9 = samples not drawn from this bottle.	unitless
Depth	Depth of sample collection; provided by the data contributor	meters (m)
BTL_DATE	Date (year, month, day) when the bottle was fired; according to the bottle file. Format: yyymmdd	unitless
BTL_TIME	Time (hours and minutes) when the bottle was fired; according to the bottle file. Format: HHMM	unitless
BTL_LAT	Latitude of bottle firing; north is positive.	decimal degrees
BTL_LON	Longitude of bottle firing; east is positive.	decimal degrees
Hg_D_CONC_BOTTLE	Total mercury measurement	picomoles per kilogram (pmol/kg)
Hg_D_CONC_BOTTLE_QF	Total mercury quality flag (WOCE)	unitless
Hg_0_D_CONC_BOTTLE	Elemental mercury measurement	picomoles per kilogram (pmol/kg)

Hg_0_D_CONC_BOTTLE_QF	Elemental mercury quality flag (WOCE)	unitless
Hg_MM_D_CONC_BOTTLE	Monomethylmercury measurement	picomoles per kilogram (pmol/kg)
Hg_MM_D_CONC_BOTTLE_QF	Monomethylmercury quality flag (WOCE)	unitless
Hg_DM_D_CONC_BOTTLE	Dimethylmercury measurement	picomoles per kilogram (pmol/kg)
Hg_DM_D_CONC_BOTTLE_QF	Dimethylmercury quality flag (WOCE)	unitless
CTDPRS	CTD pressure; joined from bottle file	decibars
CTDDEPTH	CTD bottle firing depth; joined from bottle file	meters
ODF_CTDPRS	The ODF software acquisition measurement of pressure; joined from bottle file	decibars
ODF_CTDDEPTH	The ODF software acquisition measurement of depth; joined from bottle file	meters

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Tekran Model 2500 CVAFS Mercury Detector
Generic Instrument Name	Cold Vapor Atomic Fluorescence Spectrophotometer
Dataset-specific Description	Cold Vapor Atomic Fluorescent Spectrometer (CVAFS) quantitatively measures volatile heavy metals, such as mercury. Gaseous mercury atoms are carried to the detector via inert gas, excited by a UV light source, and radiate the absorbed energy. The fluorescence is detected by either a photomultiplier tube or UV photodiode.
Generic Instrument Description	A Cold Vapor Atomic Fluorescent Spectrophotometer (CVAFS) is an instrument used for quantitative determination of volatile heavy metals, such as mercury. CVAFS make use of the characteristic of mercury that allows vapor measurement at room temperature. Mercury atoms in an inert carrier gas are excited by a collimated UV light source at a particular wavelength. As the atoms return to their non-excited state they re-radiate their absorbed energy at the same wavelength. The fluorescence may be detected using a photomultiplier tube or UV photodiode.

Dataset-specific Instrument Name	GT-C rosette
Generic Instrument Name	GO-FLO Teflon Trace Metal Bottle
Generic Instrument Description	GO-FLO Teflon-lined Trace Metal free sampling bottles are used for collecting water samples for trace metal, nutrient and pigment analysis. The GO-FLO sampling bottle is designed specifically to avoid sample contamination at the surface, internal spring contamination, loss of sample on deck (internal seals), and exchange of water from different depths.

Deployments

HLY1502

Website	https://www.bco-dmo.org/deployment/638807
Platform	USCGC Healy
Report	https://datadocs.bco-dmo.org/docs/302/geotraces/GEOTRACES_ARCTIC/data_docs/cruise_reports/healy1502.pdf
Start Date	2015-08-09
End Date	2015-10-12
Description	Arctic transect encompassing Bering and Chukchi Shelves and the Canadian, Makarov and Amundsen sub-basins of the Arctic Ocean. The transect started in the Bering Sea (60°N) and traveled northward across the Bering Shelf, through the Bering Strait and across the Chukchi shelf, then traversing along 170-180°W across the Alpha-Mendelev and Lomonosov Ridges to the North Pole (Amundsen basin, 90°N), and then back southward along ~150°W to terminate on the Chukchi Shelf (72°N). Additional cruise information is available in the GO-SHIP Cruise Report (PDF) and from the Rolling Deck to Repository (R2R): https://www.rvdata.us/search/cruise/HLY1502

Project Information

U.S. Arctic GEOTRACES Study (GN01) (U.S. GEOTRACES Arctic)

Website: <https://www.geotraces.org/>

Coverage: Arctic Ocean; Sailing from Dutch Harbor to Dutch Harbor (GN01)

Description from NSF award abstract:

In pursuit of its goal "to identify processes and quantify fluxes that control the distributions of key trace elements and isotopes in the ocean, and to establish the sensitivity of these distributions to changing environmental conditions", in 2015 the International GEOTRACES Program will embark on several years of research in the Arctic Ocean. In a region where climate warming and general environmental change are occurring at amazing speed, research such as this is important for understanding the current state of Arctic Ocean geochemistry and for developing predictive capability as the regional ecosystem continues to warm and influence global oceanic and climatic conditions. The three investigators funded on this award, will manage a large team of U.S. scientists who will compete through the regular NSF proposal process to contribute their own unique expertise in marine trace metal, isotopic, and carbon cycle geochemistry to the U.S. effort. The three managers will be responsible for arranging and overseeing at-sea technical services such as hydrographic measurements, nutrient analyses, and around-the-clock management of on-deck sampling activities upon which all participants depend, and for organizing all pre- and post-cruise technical support and scientific meetings. The management team will also lead educational outreach activities for the general public in Nome and Barrow, Alaska, to explain the significance of the study to these communities and to learn from residents' insights on observed changes in the marine system. The project itself will provide for the support and training of a number of pre-doctoral students and post-doctoral researchers. Inasmuch as the Arctic Ocean is an epicenter of global climate change, findings of this study are expected to advance present capability to forecast changes in regional and global ecosystem and climate system functioning.

As the United States' contribution to the International GEOTRACES Arctic Ocean initiative, this project will be part of an ongoing multi-national effort to further scientific knowledge about trace elements and isotopes in the world ocean. This U.S. expedition will focus on the western Arctic Ocean in the boreal summer of 2015. The scientific team will consist of the management team funded through this award plus a team of scientists from U.S. academic institutions who will have successfully competed for and received NSF funds for specific science

projects in time to participate in the final stages of cruise planning. The cruise track segments will include the Bering Strait, Chukchi shelf, and the deep Canada Basin. Several stations will be designated as so-called super stations for intense study of atmospheric aerosols, sea ice, and sediment chemistry as well as water-column processes. In total, the set of coordinated international expeditions will involve the deployment of ice-capable research ships from 6 nations (US, Canada, Germany, Sweden, UK, and Russia) across different parts of the Arctic Ocean, and application of state-of-the-art methods to unravel the complex dynamics of trace metals and isotopes that are important as oceanographic and biogeochemical tracers in the sea.

Collaborative Research: GEOTRACES Arctic Section: Mercury Speciation and Cycling in the Arctic Ocean (GEOTRACES Arctic Mercury)

Coverage: Western Arctic Ocean

NSF Award Abstract:

In this project, investigators from Woods Hole Oceanographic Institution and Wright State University participating in the 2015 U.S. GEOTRACES Arctic expedition will measure total mercury and mercury species in seawater, particles, sediments, snow and ice samples to better understand its cycling in the Arctic Ocean. In common with other multinational initiatives in the International GEOTRACES Program, the goals of the U.S. Arctic expedition are to identify processes and quantify fluxes that control the distributions of key trace elements and isotopes in the ocean, and to establish the sensitivity of these distributions to changing environmental conditions. Some trace elements are essential to life, others are known biological toxins, and still others are important because they can be used as tracers of a variety of physical, chemical, and biological processes in the sea. In its methylated form, mercury poses a serious human and ecosystem health threat, and this research will provide important information on the extent and rate of change of mercury in the sensitive Arctic ecosystem. Results from this study will be shared through outreach efforts to middle schools students in the Dayton, Ohio area, and the research will involve training for graduate and undergraduate students.

Mercury is a toxic trace metal that originates from natural and anthropogenic sources, and can enter the oceans through many processes, most importantly atmospheric deposition, riverine discharge, and coastal erosion in the Arctic Ocean. Mercury in the ocean can be transformed into varying species by a variety of abiotic and biotic processes. Its transformation to methylmercury is of primary concern as it can biomagnify in food webs. Wildlife in the Arctic has experienced unprecedented increases of methylmercury in their tissues during the past 200 years. While there has been a great deal of research related to mercury cycling in the Arctic in recent years, there remain large gaps in fundamental understanding, particularly with respect to mercury distributions and speciation. The researchers will 1) measure the concentration of four mercury species in the water column, particles, sediments, snow and sea ice samples, 2) determine the concentration of other related chemical species (thiols and snow/ice bromine), and 3) explore the mercury-related genomics of bacteria in various samples. Results from this work will yield new insights into the extent and rate of change of mercury loadings in the Arctic Ocean.

[[table of contents](#) | [back to top](#)]

Program Information

U.S. GEOTRACES (U.S. GEOTRACES)

Website: <http://www.geotraces.org/>

Coverage: Global

GEOTRACES is a [SCOR](#) sponsored program; and funding for program infrastructure development is provided by the [U.S. National Science Foundation](#).

GEOTRACES gained momentum following a special symposium, S02: Biogeochemical cycling of trace elements

and isotopes in the ocean and applications to constrain contemporary marine processes (GEOSECS II), at a 2003 Goldschmidt meeting convened in Japan. The GEOSECS II acronym referred to the Geochemical Ocean Section Studies To determine full water column distributions of selected trace elements and isotopes, including their concentration, chemical speciation, and physical form, along a sufficient number of sections in each ocean basin to establish the principal relationships between these distributions and with more traditional hydrographic parameters;

- * To evaluate the sources, sinks, and internal cycling of these species and thereby characterize more completely the physical, chemical and biological processes regulating their distributions, and the sensitivity of these processes to global change; and

- * To understand the processes that control the concentrations of geochemical species used for proxies of the past environment, both in the water column and in the substrates that reflect the water column.

GEOTRACES will be global in scope, consisting of ocean sections complemented by regional process studies. Sections and process studies will combine fieldwork, laboratory experiments and modelling. Beyond realizing the scientific objectives identified above, a natural outcome of this work will be to build a community of marine scientists who understand the processes regulating trace element cycles sufficiently well to exploit this knowledge reliably in future interdisciplinary studies.

Expand "Projects" below for information about and data resulting from individual US GEOTRACES research projects.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1434650
NSF Division of Ocean Sciences (NSF OCE)	OCE-1434653
NSF Division of Ocean Sciences (NSF OCE)	OCE-1534315

[[table of contents](#) | [back to top](#)]