

# Fish survey data from five sites in the Galapagos Islands in March of 2018

**Website:** <https://www.bco-dmo.org/dataset/740321>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2018-07-18

## Project

» [The Role of Temperature in Regulating Herbivory and Algal Biomass in Upwelling Systems](#) (Temperature and Herbivory)

Contributors	Affiliation	Role
<a href="#">Bruno, John</a>	University of North Carolina at Chapel Hill (UNC-Chapel Hill)	Principal Investigator, Contact
<a href="#">York, Amber D.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Survey data from March 2018 fish surveys at 5 sites, across the Galapagos.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Parameters](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:0.54438 E:-90.422 S:-1.28307 W:-91.431

**Temporal Extent:** 2018-03-25 - 2018-03-30

## Dataset Description

These data have been submitted to BCO-DMO and are in the process of being served.

## Methods & Sampling

SCUBA surveys were used to quantify the composition and diversity of fishes at five sites. We performed four 50x10x5 m (length, width, height) transects at each site (2 per dive, 1 per diver). This protocol was based on the ecological monitoring protocol used by the Charles Darwin Foundation and the Galapagos National Park. The diver swam along one side of the 50m long transect (the right side first, the left side second) and identified and recorded every fish present in a projected 5x5x5 m<sup>3</sup> volume (width, height, depth forward). At the end, the diver surveyed the other side of the transect. The values for each side were included for each transect in the data, but they should not be considered independent. They could, however, be used as terms nested within transect. Values in each cell are the number of fish of each species observed.

The species names for the species codes in this dataset can be found in the "2018 Galapagos Fish Survey: species list" dataset <https://www.bco-dmo.org/dataset/740929>.

## Data Processing Description

#### BCO-DMO Data Manager Processing Notes:

- \* added a conventional header with dataset name, PI name, version date
- \* modified parameter names to conform with BCO-DMO naming conventions
- \* blank values in this dataset are displayed as "nd" for "no data." nd is the default missing data identifier in the BCO-DMO system.
- \* Added lat/lon values for sites from site list
- \* Bartolomé changed to Bartolome in the data due to character restrictions

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

*Parameters for this dataset have not yet been identified*

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### The Role of Temperature in Regulating Herbivory and Algal Biomass in Upwelling Systems (Temperature and Herbivory)

**Website:** [http://github.com/johnfbruno/Galapagos\\_NSF.git](http://github.com/johnfbruno/Galapagos_NSF.git)

#### *NSF Award Abstract:*

A well-known pattern in coastal marine systems is a positive association between the biomass of primary producers and the occurrence or intensity of upwelling. This is assumed to be caused by the increase in nutrient concentration associated with upwelling, enabling higher primary production and thus greater standing algal biomass. However, upwelling also causes large, rapid declines in water temperature. Because the metabolism of fish and invertebrate herbivores is temperature-dependent, cooler upwelled water could reduce consumer metabolism and grazing intensity. This could in turn lead to increased standing algal biomass. Thus upwelling could influence both bottom-up and top-down control of populations and communities of primary producers. The purpose of this study is to test the hypothesis that grazing intensity and algal biomass are, in part, regulated by temperature via the temperature-dependence of metabolic rates. Broader impacts include the training and retention of minority students through UNC's Course Based Undergraduate Research program, support of undergraduate research, teacher training, and various outreach activities.

The investigators will take advantage of the uniquely strong spatiotemporal variance in water temperature in the Galápagos Islands to compare grazing intensity and primary production across a natural temperature gradient. They will combine field monitoring, statistical modeling, grazing assays, populations-specific metabolic measurements, and in situ herbivore exclusion and nutrient addition to measure the effects of temperature on pattern and process in shallow subtidal communities. The researchers will also test the hypothesis that grazer populations at warmer sites and/or during warmer seasons are less thermally sensitive, potentially due to acclimatization or adaptation. Finally, the investigators will perform a series of mesocosm experiments to measure the effect of near-future temperatures on herbivores, algae, and herbivory. This work could change the way we view upwelling systems, particularly how primary production is regulated and the temperature-dependence of energy transfer across trophic levels.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1737071</a>

[ [table of contents](#) | [back to top](#) ]

