

# Data from an experiment that measured the occurrence of feeding among 4 Prorocentrum minimum strains on the cryptophyte Teleaulax amphoxeia

**Website:** <https://www.bco-dmo.org/dataset/750795>

**Data Type:** experimental

**Version:** 1

**Version Date:** 2018-12-06

## Project

» [Exploring the physiological and ecological basis of mixotrophy in marine food webs](#) (Mixo Foodwebs)

Contributors	Affiliation	Role
<a href="#">Johnson, Matthew D.</a>	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
<a href="#">Rauch, Shannon</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset contains data from an experiment that measured the occurrence of feeding among 4 Prorocentrum minimum strains on the cryptophyte Teleaulax amphoxeia.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Temporal Extent:** 2015-07 - 2015-07

---

## Dataset Description

This dataset contains data from an experiment that measured the occurrence of feeding among 4 Prorocentrum minimum strains on the cryptophyte Teleaulax amphoxeia.

## Methods & Sampling

Prorocentrum minimum culturing: All cultures were maintained routinely in F/2-Si in 32 PSU seawater, at 18C and 14:10 light:dark cycle at 50 uE (u = micro). All cultures were transferred once every two weeks.

At each time point, 2 ml of cells were removed from experimental culture flasks and preserved with gluteraldehyde (1% final concentration) and stored at 4C until used to make microscopy slides. To make slides, 1 ml of preserved sample was filtered onto a black 2 um nucleopore polycarbonate filter, and then mounted on a glass microscope slide with fluorescence grade immersion oil. Slides were then counted using fluorescence microscopy and stored at -20C.

Culturing and experimental methods can be found in Johnson 2015.

## Data Processing Description

BCO-DMO Processing: modified parameter names.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Pmin_strain_feeding.csv</b> (Comma Separated Values (.csv), 1.43 KB) MD5:2b3047ba520be16ec3d7604a67e5364d Primary data file for dataset ID 750795

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Johnson, M. D. (2015). Inducible Mixotrophy in the Dinoflagellate *Prorocentrum minimum*. *Journal of Eukaryotic Microbiology*, 62(4), 431–443. doi:[10.1111/jeu.12198](https://doi.org/10.1111/jeu.12198)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
CULTURE	Prorocentrum minimum culture strain name	unitless
TIME	Sample time	minutes
REP	Replicate for each strain (treatment) n=3	unitless
Pmin_COUNT	Total number of P. minimum cells counted	unitless
OFI	Number of orange fluorescent inclusions (OFIs); OFIs are food vacuoles from ingesting phycoerythrin-containing cryptophyte prey	unitless
Pmin_OFI_COUNT	Total number of P. minimum cells with OFIs	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Instruments

<b>Dataset-specific Instrument Name</b>	
<b>Generic Instrument Name</b>	Fluorescence Microscope
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of fluorescence and phosphorescence instead of, or in addition to, reflection and absorption of visible light. Includes conventional and inverted instruments.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Exploring the physiological and ecological basis of mixotrophy in marine food webs (Mixo Foodwebs)

**Coverage:** laboratory: Woods Hole, Mass. USA

Marine phytoplankton are responsible for about half of global primary production despite being seasonally or chronically nutrient limited. To cope with this, many phytoplankton supplement their nutritional needs through mixotrophy, which involves feeding on bacteria or other algae. These microscopic Venus Fly Traps of the ocean are major players in marine microbial food webs, yet we know so little about when they feed and how their eating is balanced with photosynthesis. This research will shed light on how environmental and cellular factors control mixotrophy, and how mixotrophy and photosynthesis are integrated in the overall metabolism. While understanding the ecological role of mixotrophy in ocean food webs is center to this work, results from this study will also shed light on the evolution of mixotrophy by identifying potential tradeoffs between feeding and photosynthesis.

Mixotrophy refers to species that combine some level of phagotrophy and phototrophy, and represents a diverse array of ecological interactions and cellular and metabolic adaptations. While often perceived as an exception to the norm, mixotrophy is commonplace in marine food webs, affording phytoplankton greater ecological fitness during periods of low or limiting nutrients while stabilizing food webs. Many mixotrophs have a low chlorophyll: carbon ratio, which tends to make them poor phototrophic competitors. In turn, feeding allows these species to achieve maximum growth while in some cases also eliminating their competitors. Other mixotrophs are strong phototrophic competitors, and only feed when severely nutrient limited. This research will determine the cellular and environmental factors that lead to feeding by marine phytoplankton, and how the contrasting metabolisms of heterotrophy and photosynthesis are integrated within a cell. This research will involve laboratory-based experiments on model dinoflagellate and chrysophyte cultures. Using microscopy, physiology, proteomics and metabolomics approaches, this work will test hypotheses about the ultimate causes and consequences of mixotrophy. The major objectives are to determine 1) environmental controls for inducing mixotrophy, 2) the role of prey quality on predator selection, 3) cellular and molecular controls of mixotrophy, and 4) nutrient assimilation and integrated metabolism. Using these various research approaches, this work will produce a comprehensive view of several mixotrophs and provide new insights into cellular, ecological, and evolutionary aspects of mixotrophy. Results from this research will improve our understanding of the physiological and ecological role of mixotrophy in marine phytoplankton, and provide much needed molecular markers for studying this process in both the laboratory and field.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1436169</a>

[ [table of contents](#) | [back to top](#) ]