

# Event log from R/V Sikuliaq SKQ201701S from January to February 2017

**Website:** <https://www.bco-dmo.org/dataset/755088>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2019-01-10

## Project

» [Collaborative Research: A metabolic index to predict the consequences of climate change for midwater ecosystems](#) (Metabolic Index)

Contributors	Affiliation	Role
<a href="#">Wishner, Karen</a>	University of Rhode Island (URI)	Principal Investigator
<a href="#">Roman, Christopher Neil</a>	University of Rhode Island (URI)	Co-Principal Investigator
<a href="#">Seibel, Brad</a>	University of South Florida (USF)	Co-Principal Investigator
<a href="#">Biddle, Mathew</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Event log from R/V Sikuliaq cruise SKQ201701S, which took place from January to February 2017.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Related Datasets](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:22.42 E:-116.5 S:21.22 W:-118.05

**Temporal Extent:** 2017-01-22 - 2017-02-11

---

## Dataset Description

Date, time, and location of sampling operations.

## Data Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- added date\_MST and date.UTC to hold dates reformatted into yyyy-mm-dd.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>eventlog.csv</b> (Comma Separated Values (.csv), 12.40 KB) MD5:097302580ca7cb374c7e31dd2ca04fd8
Primary data file for dataset ID 755088

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Childress, J. J., Barnes, A. T., Quetin, L. B., & Robison, B. H. (1978). Thermally protecting cod ends for the recovery of living deep-sea animals. *Deep Sea Research*, 25(4), 419–422. doi:[10.1016/0146-6291\(78\)90568-4](https://doi.org/10.1016/0146-6291(78)90568-4)  
*Methods*

J. J. Childress, B. A. Seibel, Life at stable low oxygen levels: adaptations of animals to oceanic oxygen minimum layers. *J. Exp. Biol.* 201, 1223–1232 (1998). <http://jeb.biologists.org/content/jexbio/201/8/1223.full.pdf>  
*Methods*

Roman, C., Ullman, D. S., Hebert, D., & Licht, S. (2019). The Wire Flyer Towed Profiling System. *Journal of Atmospheric and Oceanic Technology*, 36(2), 161–182. doi:[10.1175/jtech-d-17-0180.1](https://doi.org/10.1175/jtech-d-17-0180.1)  
*Methods*

Wiebe, P. H., Morton, A. W., Bradley, A. M., Backus, R. H., Craddock, J. E., Barber, V., ... Flierl, G. R. (1985). New development in the MOCNESS, an apparatus for sampling zooplankton and micronekton. *Marine Biology*, 87(3), 313–323. doi:10.1007/bf00397811 <https://doi.org/10.1007/BF00397811>  
*Methods*

Wishner, K. F., Seibel, B. A., Roman, C., Deutsch, C., Outram, D., Shaw, C. T., ... Riley, S. (2018). Ocean deoxygenation and zooplankton: Very small oxygen differences matter. *Science Advances*, 4(12), eaau5180. doi:[10.1126/sciadv.aau5180](https://doi.org/10.1126/sciadv.aau5180)  
*Results*

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### IsRelatedTo

Seibel, B., Roman, C., Wishner, K. (2021) **Respirometry data for pelagic crustaceans, cephalopods, and fish collected on R/V Sikuliaq cruise SKQ201701S from January to February 2017**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-07-22 doi:10.26008/1912/bco-dmo.855732.1 [[view at BCO-DMO](#)]

Wishner, K., Outram, D., Seibel, B., Roman, C. (2021) **Abundances of copepod species in each net from MOCNESS tows in the Eastern Tropical North Pacific collected on four research cruises from 2007-2017**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-07-09 doi:10.26008/1912/bco-dmo.855395.1 [[view at BCO-DMO](#)]

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Event_no	event number	unitless
Gear_Type	gear type	unitless
Cast_no	cast number	unitless
Station	station	unitless
Tow_Type	tow type	unitless
Date_Local_MST	Date Local MST (MMDDYYYY)	unitless
Date_UTC	Date UTC (MMDDYYYY)	unitless
Time_In_Local_MST	Time In Local MST	unitless
Time_Out_Local_MST	Time Out Local MST	unitless
Time_In_UTC	Time In UTC	unitless
Time_Out_UTC	Time Out UTC	unitless
Lat_In	Lat in; North is positive, negative denotes South	decimal degrees
Lon_In	Lon in; East is positive, negative denotes West	decimal degrees
Lat_Out	Lat Out; North is positive, negative denotes South	decimal degrees
Lon_Out	Lon Out; East is positive, negative denotes West	decimal degrees
Day_Night	Day/Night	unitless
Min_Depth	min depth	meters (m)
Max_Depth	max depth	meters (m)
Comments	comments	unitless
date_time_in_UTC	date and time (UTC) in the water following ISO8601 yyyy-mm-ddTHH:MM	unitless
date_time_in_MST	date and time local (MST) in the water following ISO8601 yyyy-mm-ddTHH:MM	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	CTD
<b>Generic Instrument Name</b>	CTD Sea-Bird
<b>Dataset-specific Description</b>	CTD (Sikuliaq equipment, SeaBird rosette)
<b>Generic Instrument Description</b>	Conductivity, Temperature, Depth (CTD) sensor package from SeaBird Electronics, no specific unit identified. This instrument designation is used when specific make and model are not known. See also other SeaBird instruments listed under CTD. More information from Sea-Bird Electronics.

<b>Dataset-specific Instrument Name</b>	MOCNESS
<b>Generic Instrument Name</b>	MOCNESS
<b>Dataset-specific Description</b>	MOCNESS (Wishner: Multiple Opening Closing Net and Environmental Sensing System (Wiebe et al. 1985), 1 m <sup>2</sup> system, 8 or 9 nets, 222 µm mesh, updated software and Sea-Bird SBE911plusCTD system (SIO upgrades), day and night vertically- stratified and horizontally- sequenced tows, Wishner et al. 2018)
<b>Generic Instrument Description</b>	The Multiple Opening/Closing Net and Environmental Sensing System or MOCNESS is a family of net systems based on the Tucker Trawl principle. There are currently 8 different sizes of MOCNESS in existence which are designed for capture of different size ranges of zooplankton and micro-nekton Each system is designated according to the size of the net mouth opening and in two cases, the number of nets it carries. The original MOCNESS (Wiebe et al, 1976) was a redesigned and improved version of a system described by Frost and McCrone (1974).(from MOCNESS manual) This designation is used when the specific type of MOCNESS (number and size of nets) was not specified by the contributing investigator.

<b>Dataset-specific Instrument Name</b>	Wire Flyer
<b>Generic Instrument Name</b>	Sea-Bird SBE 49 FastCAT CTD Sensor
<b>Dataset-specific Description</b>	Wire Flyer (Roman development: oscillating deep towed hydrographic profiler with Sea-Bird 49 FastCAT CTD and Aanderaa 4831F oxygen sensor, Roman et al 2018)
<b>Generic Instrument Description</b>	The SBE 49 FastCAT is a CTD sensor for use in autonomous platforms. It contains a SBE 3P temperature sensor, a SBE 4C conductivity sensor and a strain-gauge pressure sensor as standard. It can operate in autonomus (16 Hz per sec) or polled mode (transmits each sample). The sensor is depth-rated to 350 m (plastic housing) or 7000 m (titanium housing). Accuracy: +/- 0.002 deg C (temperature), +/- 0.0003 S/m (conductivity), 0.1% of full scale range (pressure).

<b>Dataset-specific Instrument Name</b>	Tucker trawl net
<b>Generic Instrument Name</b>	Tucker Trawl
<b>Dataset-specific Description</b>	MTT + MOC (Seibel: large Tucker trawl net (3 m <sup>2</sup> mouth, 100 m long) with MOCNESS software and sensors (Childress and Seibel 1998), large insulated closing cod end (Childress et al. 1978), live animal collections)
<b>Generic Instrument Description</b>	The original Tucker Trawl, a net with a rectangular mouth opening first built in 1951 by G.H. Tucker, was not an opening/closing system, but shortly thereafter it was modified so that it could be opened and closed. The original had a 183 cm by 183 cm flexible rectangular mouth opening 914 cm long net with 1.8 cm stretched mesh for the first 457 cm and 1.3 cm mesh for last 457 cm. 152 cm of coarse plankton or muslin netting lined the end of the net. Tucker designed the net to collect animals associated with the deep scattering layers, principally euphausiids, siphonophores, and midwater fish. (from Wiebe and Benfield, 2003). Currently used Tucker Trawls usually have 1-m <sup>2</sup> openings and can have a single net or multiple nets on the frame.

## Deployments

### SKQ201701S

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/755461">https://www.bco-dmo.org/deployment/755461</a>
<b>Platform</b>	R/V Sikuliaq
<b>Start Date</b>	2017-01-19
<b>End Date</b>	2017-02-15
<b>Description</b>	See additional cruise information from R2R: <a href="https://www.rvdata.us/search/cruise/SKQ201701S">https://www.rvdata.us/search/cruise/SKQ201701S</a>

## Project Information

### **Collaborative Research: A metabolic index to predict the consequences of climate change for midwater ecosystems (Metabolic Index)**

**Coverage:** Eastern Tropical North Pacific

#### *Description from NSF award abstract:*

With climate change, ocean temperatures are expected to increase which in turn will reduce oxygen availability and increase metabolic oxygen demand in marine organisms. The investigators will conduct shipboard physiological experiments for various marine organisms and determine their distributions in relation to environmental conditions within an oxygen minimum zone (OMZ) in the Eastern Pacific Ocean. The goal will be to model and map a Metabolic Index (MI) to predict how vertical and horizontal distributions for these species might change throughout the world's oceans in the future. The MI is defined as the ratio between environmental oxygen supply and temperature-dependent oxygen demand. Oxygen supply includes both the environmental oxygen concentration across a habitat range and the physiological features of organisms that facilitate oxygen uptake, such as gills and circulatory systems. Thus, the MI will integrate measured tolerance and environmental exposure to low oxygen with environmental data. The investigators will measure tolerance to low oxygen, focusing on under-studied organisms, including the effect of temperature and organism size. They will sample along a natural gradient in oxygen content south of the California Current in the Eastern Pacific. The science team and a videographer will develop a blog about deep-sea biology and climate change using web-based and video technologies. Four graduate students will be funded on this project, and in conjunction with a recently developed course in pelagic ecology, several undergraduates will have the opportunity to participate in seagoing research.

This research fills a critical need for a physiology-based metric that can be used to predict changing marine communities as the oceans warm and hypoxic zones expand. Modern OMZs are extensive and characterized by deep-water (300-800 m) oxygen partial pressures lethal to most marine organisms, yet thriving communities exist there. Climate change is predicted to further deplete oxygen. The investigators will model and map a Metabolic Index (MI) for diverse marine species to help predict how in vertical and horizontal distributions of species may change throughout the world's oceans in the future. The MI will derive oxygen supply and demand data from published and planned measurements of the minimum environmental partial pressure of oxygen to which individual species are exposed (based on their distributions in the water column) and the minimum requirements to support routine aerobic metabolic demand (from shipboard respiration measurements of individuals). During research cruises in the Eastern Pacific along a gradient of OMZ intensity, the investigators will conduct shipboard physiological measurements to determine metabolic demand for understudied mesozooplankton and gelatinous taxa and determine the size- and temperature dependence for diverse species for incorporation into the MI. Vertically-stratified net sampling and in situ photography will identify and characterize unique OMZ community features, such as the lower oxycline biomass peak present in some OMZs and the oxygen-dependence of day and night habitat depths for vertically-migrating species. The MI will be mapped using climatological data to both test and generate hypotheses about the response of oceanic communities to climate change. In preliminary analysis, the MI suggests a metabolic constraint at a MI

of ~2 that may act to limit vertical and horizontal habitat ranges.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1459243</a>

[ [table of contents](#) | [back to top](#) ]