

HADES-K CN concentrations and ratios and d13C bacterial biomass signatures in Kermadec Trench during RV/Thompson cruise TN309, May 2014

Website: <https://www.bco-dmo.org/dataset/764208>

Data Type: Cruise Results

Version: 0

Version Date: 2019-04-08

Project

» [Controls on Hadal Megafaunal Community Structure: a Systematic Examination of Pressure, Food Supply, and Topography](#) (HADES)

Contributors	Affiliation	Role
Shank, Timothy M.	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
Drazen, Jeffrey C.	University of Hawaii at Manoa (SOEST)	Co-Principal Investigator
Yancey, Paul	Whitman College	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

This dataset includes C and N concentrations, C:N ratios, and d13C bacterial biomass signatures in Kermadec Trench near New Zealand from sediment cores collected on R/V Thomas G. Thompson cruise TN309, May 2014.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Supplemental Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:-31.9307423 E:-177.290253 S:-35.905159 W:-178.961227

Temporal Extent: 2014-05-02 - 2014-05-08

Dataset Description

This dataset includes C and N concentrations, C:N ratios, and d13C bacterial biomass signatures in Kermadec Trench near New Zealand from sediment push cores collected on R/V Thomas G. Thompson cruise TN309, May 2014.

Data not yet submitted.

Methods & Sampling

Push cores (6.35 cm diameter) of sediment were collected in situ using the manipulator arm of the Hybrid Remotely Operated Vehicle Nereus on the margin of the Kermadec Trench (6013m) and along the main trench

axis (7137 to 9177m) at roughly 1000m increments.

Cores were sectioned at 1cm intervals and sieved for the meiofaunal fraction.

Upon retrieval of sediment cores, the 0-1, 1-2, 2-3, 3-5, 5-10 cm depth horizons were sectioned and stored frozen (-80 °C) until analysis. Food availability, described as the bulk quantities of organic carbon and nitrogen and benthic bacterial biomass, were determined from freeze dried and decalcified sediment samples with a Fisons NA 1500 elemental analyzer using sulphhanillic acid as a standard.

Attempts were made to obtain 3 replicate cores at each depth horizon (every 1000 meters from 6000m to 10,000m. For logistical reasons with vehicle performance (and loss at 10,000), all of the replicates were not able to be collected.

Data Processing Description

BCO-DMO Processing Notes:

[[table of contents](#) | [back to top](#)]

Supplemental Files

File
Sediment sample C and N dry weights vs. depth (6011-7142 meters) filename: image002_core_CN.png (Portable Network Graphics (.png), 66.48 KB) MD5:e94d708d42a722d8b755c833df0bfd4 Sediment sample C and N dry weights vs. depth (6011-7142 meters)
Sediment sample C and N dry weights vs. depth (8081-9177 meters) filename: image004_core_CN.png (Portable Network Graphics (.png), 64.11 KB) MD5:681d6f77705d40a92f18a5f85829a31a Sediment sample C and N dry weights vs. depth (8081-9177 meters)
Sediment sample CN ratios and delta-13C vs. core depth (6011-7142 meters) filename: image011_core_CN_d13C.png (Portable Network Graphics (.png), 65.15 KB) MD5:0367baa16cd239f4ec3fccc000fc39f
Sediment sample CN ratios and delta-13C vs. core depth (8081-9177 meters) filename: image012_core_CN_d13C.png (Portable Network Graphics (.png), 60.89 KB) MD5:226d80a133bb1611f5eced115be2d81f

[[table of contents](#) | [back to top](#)]

Parameters

Parameters for this dataset have not yet been identified

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Fisons NA 1500 elemental analyzer
Generic Instrument Name	Elemental Analyzer
Generic Instrument Description	Instruments that quantify carbon, nitrogen and sometimes other elements by combusting the sample at very high temperature and assaying the resulting gaseous oxides. Usually used for samples including organic material.

Dataset-specific Instrument Name	
Generic Instrument Name	HROV Nereus
Generic Instrument Description	Nereus is an efficient, multi-purpose “hybrid” vehicle that can explore and operate in the crushing pressures of the greatest ocean depths. An unmanned vehicle, Nereus operates in two complementary modes. It can swim freely as an autonomous underwater vehicle (AUV) to survey large areas of the depths, map the seafloor, and give scientists a broad overview. When Nereus locates something interesting, the vehicle’s support team can bring the vehicle back on board the ship and transforms it into a remotely operated vehicle (ROV) tethered to the ship via a micro-thin, fiber-optic cable. Through this tether, Nereus can transmit high-quality, real-time video images and receive commands from skilled pilots on the ship to collect samples or conduct experiments with a manipulator arm. Technical specifications: Weight on land: 2,800 kg Payload capacity: 25 kg Maximum speed: 3 knots Batteries: rechargeable lithium ion, 15 kilowatt hours in two pressure housings Thrusters: 2 fore and aft, 2 vertical, 1 lateral (ROV mode) 2 fore and aft, 1 vertical (AUV mode) Lights: variable output LED array, strobes Manipulator arm: Kraft TeleRobotics 7-function hydraulic manipulator Sonar: scanning sonar, forward look and profile, 675 KHz Sensors: magnetometer, CTD (to measure conductivity, temperature, and depth) Nereus supports a variety of science operations: Push coring, measuring heat flow, geotechnical and geochemical sensing, rock sampling and drilling, biological sampling, water sampling, high resolution acoustic bathymetry, and optical still and video imagery. More information is available from the operator site at URL.

Dataset-specific Instrument Name	
Generic Instrument Name	Push Corer
Dataset-specific Description	Push cores were borrowed from the National Deep Submergence Facility to be able to standardize results. The push cores were 6.35cm diameter).
Generic Instrument Description	Capable of being performed in numerous environments, push coring is just as it sounds. Push coring is simply pushing the core barrel (often an aluminum or polycarbonate tube) into the sediment by hand. A push core is useful in that it causes very little disturbance to the more delicate upper layers of a sub-aqueous sediment. Description obtained from: http://web.whoi.edu/coastal-group/about/how-we-work/field-methods/coring/

[[table of contents](#) | [back to top](#)]

Deployments

TN309

Website	https://www.bco-dmo.org/deployment/536488
Platform	R/V Thomas G. Thompson
Start Date	2014-04-10
End Date	2014-05-20
Description	Original data are available from the NSF R2R data catalog

[[table of contents](#) | [back to top](#)]

Project Information

Controls on Hadal Megafaunal Community Structure: a Systematic Examination of Pressure, Food Supply, and Topography (HADES)

Website: <http://www.whoi.edu/hades/>

Coverage: Kermadec Trench adjacent to New Zealand: approximately 37 12.75 S and 178 51.43 E to 31 51.29 S and 176 49.07 W


Extracted from the NSF award abstract:

Severe technical challenges associated with the extremes of hydrostatic pressure have prevented major advances in hadal ecological studies, and relegated hadal systems to among the most poorly investigated habitats on Earth. Through this project, Hadal Ecosystems Studies (HADES) program, PIs will determine the composition and distribution of hadal species, the role of hadal pressures (piezolyte concentrations, enzyme function under pressure), food supply (distribution of POC with the abundance and biomass of trench organisms, and metabolic rates/energetic demand), and depth/topography (genetic divergence and spatial connectivity of populations) have on impacting deep-ocean community structure. This project will examine these factors using the world's first full-ocean depth hybrid remotely operated vehicle (HROV) in conjunction with the only full-ocean depth imaging lander (Hadal-Lander). This project will provide the first seafloor data and samples in one of the world's best, yet little known trenches- the Kermadec Trench (SW Pacific Ocean).

Megafaunal community structure and the relationship between POC and benthic bacterial biomass will be examined as a function of depth and location by systematic high-definition imaging and sediment/faunal sampling transects from abyssal to full trench depths both along and perpendicular to the trench axis. Population genetic approaches will provide levels of genetic divergence and evolutionarily independent lineages to assess the role of depth and topography in trenches and their adjacent abyssal plain in promoting the formation of species. Physiological constraints will be investigated by examining in-situ respiration of selected fauna and tissue concentrations of such protein stabilizers as trimethylamine oxide (TMAO), and the structural adaptations of macromolecules.

Image of NEREUS Deployment Sites. [click on the image to view a larger version]

HADES-K Deployment sites



[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1131620

[[table of contents](#) | [back to top](#)]