

Gridded in-situ oxygen profiles from glider deployments in the San Pedro Channel, CA in 2013 and 2014

Website: <https://www.bco-dmo.org/dataset/768685>

Data Type: Other Field Results

Version: 2

Version Date: 2019-07-03

Project

» [Collaborative Research: Use of Triple Oxygen Isotopes and O₂/Ar to constrain Net/Gross Oxygen Production during upwelling and non-upwelling periods in a Coastal Setting](#) (UpRISEE O₂ upwelling)

Contributors	Affiliation	Role
Haskell, William	University of California-Santa Barbara (UCSB-MSI)	Principal Investigator
Hammond, Douglas E.	University of Southern California (USC)	Co-Principal Investigator
Prokopenko, Maria	Pomona College (Pomona)	Co-Principal Investigator
Jones, Burton	King Abdullah University of Science and Technology (KAUST)	Scientist, Contact
Levine, Naomi M.	University of Southern California (USC)	Scientist
Teel, Elizabeth N.	University of Southern California (USC)	Scientist, Contact
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Gridded in-situ oxygen profiles from glider deployments in the San Pedro Channel, CA in 2013 and 2014.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:33.72 E:-118.25003 S:33.45003 W:-118.59992

Temporal Extent: 2013-03-12 - 2014-07-07

Dataset Description

Related dataset (collected during same glider deployments):

Gridded in-situ profiles: <https://www.bco-dmo.org/dataset/751128>

In addition to the tabular form of this dataset available by clicking the "Get Data" button on this page, it is also available as a Matlab (*.mat) file containing data matrices and vectors. See the "Supplemental Files" section on this page for the data in this form along with the matlab m-file used to transform the data matrices to a tabular dataset.

Methods & Sampling

Glider deployment: For this study, a slocum glider was deployed between March and July in 2013 and 2014 in the San Pedro Channel, located in the Southern California Bight off the coast of Los Angeles. The glider was deployed on a 28 km cross-channel path between Catalina Island and the Palos Verdes Peninsula and completed a single cross-channel pass every 1.5–2 days (average speed 1 km h⁻¹). Data were collected between ~ 3 and 90 m, with the exception of when the glider crossed the major shipping lanes where the glider was constrained to depths below 20 m to avoid damage or loss from ship traffic. The GMT time stamp indicates the earliest time stamp for data binned within a single profile. Glider deployments were from small boats belonging to the USC Wrigley Marine Sciences Center or from the R/V Yellowfin.

Optode oxygen concentration was corrected for measurement lag and calibrated using Winkler titrations.

Data Processing Description

BCO-DMO Data Manager Processing Notes: Data Version 1: 2018-12-14

- * tabular dataset constructed from original .mat data file
- * tabular dataset imported into the BCO-DMO data system with added header with dataset name, PI name, version date.
- * original matlab file served as a downloadable link on this dataset landing page.
- * blank values in the BCO-DMO data system are displayed as "nd" for "no data." nd is the default missing data identifier in the BCO-DMO system. Blank values in the original .mat file are the matlab default value NaN.

BCO-DMO Data Manager Processing Notes: Data Version 2: 2019-07-03

- * Same processing performed on version 1 as version 1.
- * New version had lat/lon values corrected by the data submitter.
- * Version 2 has column OXYGEN which was called O2 in version 1.

[[table of contents](#) | [back to top](#)]

Data Files

File
GliderO2.csv (Comma Separated Values (.csv), 80.53 MB) MD5:52fc8cd42d676aaa2262c0b9c945d662 Primary data file for dataset ID 768685

[[table of contents](#) | [back to top](#)]

Supplemental Files

File
Gridded in-situ oxygen profiles as matrices (MATLAB .mat) filename: GliderO2.mat (Octet Stream, 23.13 MB) MD5:c37ecd362d8abdcd5cb3776b8f62a56e Matlab (.mat) file containing gridded in-situ oxygen profiles as matrices.
Matlab script to transform gridded O2 profile matrices into a tabular dataset filename: GliderO2_mat2csv.m (Octet Stream, 3.60 KB) MD5:e81ed093b0f313f0e12e80ae4851718d Matlab script to transform data matrices to a tabular dataset. The input MATLAB .mat file can be found here https://www.bco-dmo.org/node/772250 .

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
BINID	Bin identifier	unitless
DATENUM_GMT	Matlab datenum (Greenwich Mean Time)	unitless
ISO_DateTime_UTC	Timestamp (UTC) in standard ISO 8601:2004(E) format yyyy-mm-ddTHH:MM:SSZ	yyyy-MM-dd'THH:mm:ss'Z'
YEAR	Year in format yyyy	unitless
LAT	Latitude	decimal degrees (DD)
LON	Longitude	decimal degrees (DD)
DEPTH	Depth	meters (m)
OXYGEN	Oxygen concentration	millimoles per cubic meter (mmol m ⁻³)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Aanderaa Oxygen Optode 5013w (3835)
Generic Instrument Name	Aanderaa Oxygen Optodes
Generic Instrument Description	Aanderaa Oxygen Optodes are instrument for monitoring oxygen in the environment. For instrument information see the Aanderaa Oxygen Optodes Product Brochure.

Dataset-specific Instrument Name	Sea-Bird flow-through CTD
Generic Instrument Name	CTD Sea-Bird
Dataset-specific Description	Sea-Bird flow-through CTD (used for temperature, salinity, and pressure)
Generic Instrument Description	Conductivity, Temperature, Depth (CTD) sensor package from SeaBird Electronics, no specific unit identified. This instrument designation is used when specific make and model are not known. See also other SeaBird instruments listed under CTD. More information from Sea-Bird Electronics.

[[table of contents](#) | [back to top](#)]

Deployments

UpRISEE_SPOT_13-14

Website	https://www.bco-dmo.org/deployment/684011
Platform	R/V Yellowfin
Start Date	2013-01-16
End Date	2014-06-19
Description	A series of cruises were conducted from January 2013 to June 2014 to the San Pedro Ocean Time-Series (SPOT) station. These cruises were part of a study aimed at characterizing the biological response to upwelling at SPOT: the Upwelling Regime In-Situ Ecosystem Efficiency (Up.R.I.S.E.E.) study.

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Use of Triple Oxygen Isotopes and O₂/Ar to constrain Net/Gross Oxygen Production during upwelling and non-upwelling periods in a Coastal Setting (UpRISEE O₂ upwelling)

Coverage: Northeast Pacific Ocean

The marine biological pump is one of the primary pathways via which anthropogenic carbon dioxide may be sequestered from the atmosphere and exported to the deep ocean as organic carbon. While the link between nutrient supply and high primary productivity in upwelling regions is well established, factors controlling the organic carbon export efficiency of upwelling ecosystems are not well known. Scientists from the University of Southern California and Pomona College plan to determine the factors that control the rates and magnitudes of two components of biological production, Net Community Production (NCP) and Gross Primary Production (GPP), as well as particulate organic carbon export efficiency, at the San Pedro Ocean Time Series, a coastal site in the California Borderland during periods of minimal and high upwelling velocity over a 2-year span. At this site, past and ongoing observations of hydrography and carbon rain will provide an historical context for interpreting results and mechanisms at work.

Rates of NCP and GPP will be quantified at different upwelling intensity, using dissolved oxygen to argon (O₂/Ar) ratios and the oxygen triple isotope composition of dissolved oxygen (O₂). The export of organic carbon will be established using ²³⁴Th (thorium) profiles in the water column, coupled with floating sediment trap deployments, and the development of a carbon isotope balance for the water column. Upwelling will be characterized using non-steady state budgets for atmospheric ⁷Be (beryllium) input and its depth-integrated decay, as well as estimating rates based on remote measurements of wind stress curl and budgets for dissolved inorganic carbon and silicon. Application of the O₂/Ar ratio and the oxygen triple isotope approach will require depth-integrated profiles of these tracers to evaluate the impact of upwelling on mixed layer inputs and use of non-steady state models during seasonal transitions in upwelling. The comprehensive data set to be obtained should provide insights into the organic carbon export efficiency under variable upwelling regimes and help to relate the satellite-based measurements of chlorophyll to the organic carbon export of these highly productive ecosystems.

Broader Impacts: One graduate and one undergraduate student from the University of Southern California and two undergraduate students from Pomona College would be supported and trained as part of this project.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1260296
NSF Division of Ocean Sciences (NSF OCE)	OCE-1260692

[[table of contents](#) | [back to top](#)]