

# Net tow metaproteome of Trichodesmium species mapped to a Trichodesmium metagenome plus cyanoGEBA species genomes in units of normalized peptide spectral counts from samples collected in the Atlantic and Pacific Ocean between 2000 and 2018

**Website:** <https://www.bco-dmo.org/dataset/787168>

**Data Type:** Cruise Results

**Version:** 1

**Version Date:** 2020-04-30

## Project

- » [Collaborative Research: Iron and phosphorus balanced limitation of nitrogen fixation in the oligotrophic ocean \(TriCoLim\)](#)
- » [New technology for high resolution analysis of proteins and other organic materials produced by marine microorganisms](#) (MM Proteins and Organics Tech)
- » [Marine Microbial Investigator Award: Investigator Mak Saito](#) (MM Saito)
- » [Collaborative Research: Evolutionary, biochemical and biogeochemical responses of marine cyanobacteria to warming and iron limitation interactions](#) (Cyanobacteria Warming Responses)

## Program

- » [Marine Microbiology Initiative](#) (MMI)

Contributors	Affiliation	Role
<a href="#">Saito, Mak A.</a>	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
<a href="#">Held, Noelle</a>	Woods Hole Oceanographic Institution (WHOI)	Student
<a href="#">York, Amber D.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Net tow metaproteome of Trichodesmium species mapped to a Trichodesmium metagenome plus cyanoGEBA species genomes, analyzed by 2D LC-MS/MS in units of normalized peptide spectral counts. Samples were collected in North Atlantic surface waters, at station BATS (Bermuda Atlantic Time-series Study), and station ALOHA (A Long-Term Oligotrophic Habitat Assessment) between 2000 and 2018.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:23.22 E:-21.59 S:0.17 W:-158

**Temporal Extent:** 2000-07-27 - 2018-03-18

## Dataset Description

Net tow metaproteome of Trichodesmium species mapped to a Trichodesmium metagenome plus cyanoGEBAs species genomes, analyzed by 2D LC-MS/MS in units of normalized peptide spectral counts. Samples were collected in North Atlantic surface waters, at station BATS (Bermuda Atlantic Time-series Study), and station ALOHA (A Long-Term Oligotrophic Habitat Assessment) between 2000 and 2018. Related datasets: Trichodesmium field metaproteomes - sequence fasta: <https://www.bco-dmo.org/dataset/787181> Trichodesmium field metaproteomes - protein spectral counts: <https://www.bco-dmo.org/dataset/787147> Trichodesmium sample provenance: <https://www.bco-dmo.org/dataset/787093> - Sample provenance file, which includes sample locations, filter sizes

## Methods & Sampling

A 200um plankton net was deployed to ~20m depth, then recovered to just below surface, repeating five times. Trichodesmium colonies were hand picked into 0.2um filtered surface seawater, rinsed twice in 0.2um filtered surface seawater, and decanted onto a 0.2-4um Supor filter (indicated in sample provenance table, see dataset <https://www.bco-dmo.org/dataset/787093>).

Proteins were extracted and trypsin digested in-gel following Saito et al., 2014 (Science). Peptides were analyzed by LC-MS/MS in data discovery mode on a Thermo Orbitrap Fusion.

## Data Processing Description

Thermo proteome discoverer 2.2 was used to search the data with the SEQUEST algorithm. Statistical validation was performed in Scaffold (Proteome Software) at the 1% protein and peptide false discovery rate (FDR) levels. This file represents the peptide identifications made as normalized spectral counts. Peptides may be assigned to more than one protein.

A publicly available Trichodesmium metagenome was used in this search (IMG ID 2821474806) as well as the contents of the CyanoGEBAs project (Shih et al, 2013).

BCO-DMO Data Manager Processing Notes:

- \* prefix "20" added to year values to change format from yy-mm-dd to ISO 8601 format yyyy-mm-dd
- \* Time values "00009" changed to 09:00 after checking the sample log.
- \* Delimiter for multiple IDs in the other\_protein\_ids column changed from comma to semicolon.
- \* Rows with missing date,time, cruise, station, and location information missing were removed (filtered to remove all rows with time value "00nan" )
- \* Commas removed from number values in columns median\_retention\_time, total\_precursor\_intensity,TIC so they could be correctly typed as numbers not strings.

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>Trichodesmium field metaproteomes - peptide spectral counts</b> filename: peptides.csv (Comma Separated Values (.csv), 43.72 MB) MD5:bd1a6d484f598281daff452c63f69d6b Net tow metaproteome of Trichodesmium species mapped to a Trichodesmium metagenome plus cyanoGEBAs species genomes in units of normalized peptide spectral counts from samples collected in the Atlantic and Pacific Ocean between 2000 and 2018.  Parameters in this comma delimited file are included in the Parameter section of the Dataset Landing Page "Trichodesmium field metaproteomes - peptide spectral counts" <a href="https://www.bco-dmo.org/dataset/787168">https://www.bco-dmo.org/dataset/787168</a> .

[ [table of contents](#) | [back to top](#) ]

## Related Publications

Saito, M. A., McIlvin, M. R., Moran, D. M., Goepfert, T. J., DiTullio, G. R., Post, A. F., & Lamborg, C. H. (2014). Multiple nutrient stresses at intersecting Pacific Ocean biomes detected by protein biomarkers. *Science*, 345(6201), 1173–1177. <https://doi.org/10.1126/science.1256450>  
*Methods*

Shih, P. M., Wu, D., Latifi, A., Axen, S. D., Fewer, D. P., Talla, E., ... Kerfeld, C. A. (2012). Improving the coverage of the cyanobacterial phylum using diversity-driven genome sequencing. *Proceedings of the National Academy of Sciences*, 110(3), 1053–1058. doi:[10.1073/pnas.1217107110](https://doi.org/10.1073/pnas.1217107110)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
protein_id	Unique protein peptide belongs to	unitless
protein_molecular_weight_kDa	Molecular weight of the sample (no data, column included to comply with Ocean Protein Portal template)	kilo-Daltons (kDa)
peptide_sequence	Peptide genomic sequence	unitless
best_peptide_id_probability	Probability (0-1) the peptide belongs to this protein.	unitless
best_sequest_DCn_score	delta Cn score	unitless
best_sequest_Xcorr_score	Xcorr score	unitless
plus_2H_spectra_count	Peptide in the 2+ state spectral count	unitless
plus_3H_spectra_count	Peptide in the 3+ state spectral count	unitless
plus_4H_spectra_count	Peptide in the 4+ state spectral count	unitless
spectral_count_sum	Sum of spectral counts (+2, +3, +4 ions)	unitless
median_retention_time	Median retention time of peptide	seconds (s)
total_precursor_intensity	Total precursor intensity at the MS1 stage of the peptide	unitless
TIC	Total ion current (TIC) value of peptide	unitless
peptide_start_index	Start location of peptide within protein sequence	unitless
peptide_stop_index	Stop location of peptide within protein sequence	unitless
other_protein_ids	Additional proteins the peptide may belong to. Multiple identifiers delimited with a semicolon.	unitless
sample_id	Unique sample location ID	unitless
cruise_id	Cruise name	unitless
station_id	Station name	unitless
cruise_station	Station identifier where sample was taken	unitless
time_h-m-s	Time (Local ship time) of sampling in format HH:MM (various time zones, see sample provenance dataset <a href="http://lod.bco-dmo.org/id/dataset/787093">http://lod.bco-dmo.org/id/dataset/787093</a> )	unitless
date_y-m-d	Date (Local ship time) in format yyyy-mm-dd (various time zones, see sample provenance dataset <a href="http://lod.bco-dmo.org/id/dataset/787093">http://lod.bco-dmo.org/id/dataset/787093</a> )	unitless
maximum_filter_size_microns	Minimum size of the collection filter	microns (?m)
minimum_filter_size_microns	Maximum size of the collection filter	microns (?m)
latitude_dd	Latitude (South is negative)	decimal degrees (DD)
longitude_dd	Longitude (West is negative)	decimal degrees (DD)
depth_m	Sample depth	meters (m)
sample_type	Sample type "Trichodesmium net tow"	unitless

## Instruments

<b>Dataset-specific Instrument Name</b>	Thermo Orbitrap Fusion mass spectrometer
<b>Generic Instrument Name</b>	Mass Spectrometer
<b>Generic Instrument Description</b>	General term for instruments used to measure the mass-to-charge ratio of ions; generally used to find the composition of a sample by generating a mass spectrum representing the masses of sample components.

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### AT39-05

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/765978">https://www.bco-dmo.org/deployment/765978</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2018-02-11
<b>End Date</b>	2018-03-14
<b>Description</b>	For study of iron and phosphorus balanced limitation of nitrogen fixation in the oligotrophic ocean.

### JC150

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/786689">https://www.bco-dmo.org/deployment/786689</a>
<b>Platform</b>	RRS James Cook
<b>Start Date</b>	2017-06-26
<b>End Date</b>	2017-08-12

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### **Collaborative Research: Iron and phosphorus balanced limitation of nitrogen fixation in the oligotrophic ocean (TriCoLim)**

**Coverage:** Tropical Atlantic

NSF abstract:

Marine cyanobacteria are able to use or "fix" atmospheric nitrogen gas, and so supply much of the essential nutrient nitrogen that supports open ocean food chains. Oceanographers have usually thought that the growth of these nitrogen-fixing cyanobacteria is limited at any particular time and place by the supply of either iron, or of phosphorus. Preliminary experiments have shown, though, that these nitrogen fixers instead grow best when both iron and phosphorus are scarce at the same time. In this project, the researchers will use cellular indicators that are specific for iron and phosphorus limitation to determine how important this type of "balanced limitation" of nitrogen-fixing cyanobacteria is in controlling the productivity of ocean food chains in

the tropical Atlantic Ocean. Two graduate students will be trained at the University of Southern California (USC) and Woods Hole Oceanographic Institution, as well as a postdoctoral researcher at USC. Educational outreach efforts will take place at a Los Angeles inner city high school with a student body that is over 98% Hispanic and African-American, and with underrepresented undergraduates in the USC Global Environmental Microbiology course. In addition, two Research Experiences for Undergraduates students will be supervised for summer research projects to help them learn about science career options.

The researchers will investigate the biological and biogeochemical consequences of this unique balanced iron/phosphorus-limited phenotype, using both laboratory and fieldwork approaches. During the first year of this project, the nitrogen-fixing cyanobacteria will be cultured under iron and/or phosphorus limitation, followed by application of proteomics and transcriptomics to identify genes that are potential diagnostic biomarkers for iron/phosphorus balanced limitation. Preliminary work has already identified one promising candidate biomarker in one cyanobacterium, an EzrA protein domain that appears to be associated with the cell size decreases seen specifically under balanced limitation, and the researchers have identified numerous other potential candidates for similar biomarkers. During the second year, these new co-limitation biomarkers and others previously validated for iron limitation (IsiB) and phosphorus limitation (SphX) will be used to investigate balanced limitation during a research cruise transecting from relatively high-iron, low-phosphorus North Atlantic waters, to the relatively high-phosphorus, low-iron South Atlantic. This fieldwork component will survey nitrogen fixing cyanobacteria populations across this natural iron/phosphorus gradient for genetic, proteomic, and physiological indicators of balanced limitation, as well as testing their responses to iron and phosphorus manipulations in shipboard incubation experiments. The third year will be devoted to sample analysis, and publications exploring the responses of oceanic nitrogen fixers to simultaneous limitation by both iron and phosphorus.

### **New technology for high resolution analysis of proteins and other organic materials produced by marine microorganisms (MM Proteins and Organics Tech)**

**Website:** <https://www.moore.org/grant-detail?grantId=GBMF3934>

In support of acquiring a high resolution mass spectrometer that incorporates the latest technologies for analyzing proteins and other organic materials.

### **Marine Microbial Investigator Award: Investigator Mak Saito (MM Saito)**

In support of obtaining deeper knowledge of major biogeochemically relevant proteins to inform a mechanistic understanding of global marine biogeochemical cycles.

### **Collaborative Research: Evolutionary, biochemical and biogeochemical responses of marine cyanobacteria to warming and iron limitation interactions (Cyanobacteria Warming Responses)**

NSF abstract:

The oceans absorb much of the heat generated by human activities, and this warming of the surface ocean has consequences for important groups of marine organisms. Marine cyanobacteria are one such key group of organisms, since they supply much of the essential carbon and nitrogen that supports nearly all the rest of the marine food web. Currently, the growth of cyanobacteria is mostly constrained by scarce supplies of the micronutrient element iron, but they are also very sensitive to the ongoing increases in seawater temperature. Preliminary results suggest that warming could partly mitigate the negative effects of iron limitation on marine cyanobacteria. This project examines in depth how these interactions between warming and iron limitation will affect the future ocean carbon and nitrogen cycles, using laboratory culture experiments showing how cyanobacteria respond to simultaneously changing temperature and iron supplies. Both short-term response studies and long-term evolutionary experiments testing for adaptation use a comprehensive set of molecular

biology tools targeting genes to proteins. The final goal is to apply the results of these experiments to improve quantitative models predicting how the ocean's carbon and nitrogen cycles, biological productivity, and living resources will respond to a warming future climate. Two graduate students, a postdoc and 3-4 underrepresented undergraduate researchers are supported, and the investigators also mentor summer science interns from largely Hispanic local high schools.

The physiology, biochemistry and biogeography of nitrogen-fixing cyanobacteria and unicellular picocyanobacteria are strongly influenced by temperature, subjecting them to intense selective pressure as the modern ocean steadily warms up. These groups have likewise been rigorously selected under chronic iron (Fe) scarcity, and the availability of this crucial micronutrient is also changing with a shifting climate. This project examines short-term acclimation and long-term evolutionary responses of Fe-stressed marine cyanobacteria to a warmer environment. Preliminary data show that Iron Use Efficiencies (IUE, mols N fixed.hr<sup>-1</sup> mol cellular Fe<sup>-1</sup>) of Fe-limited *Trichodesmium* increase 4 to 5-fold with a 5°C temperature increase, allowing the cells to much more efficiently leverage scarce available Fe supplies to grow and fix nitrogen. This means that warming can to a large degree mitigate the negative effects of Fe limitation on *Trichodesmium*, resulting in a modelled 22% increase in global nitrogen fixation by 2100 in a warmer climate. This project aims to uncover the cellular biochemical mechanisms involved in this Fe-limitation/thermal IUE effect in a four-year experimental evolution study of the diazotrophs *Trichodesmium* and *Crocospaera* and the picocyanobacteria *Synechococcus* and *Prochlorococcus*, under a multi-variate selection matrix of temperature and Fe availability. The objectives are to 1) Assess the long-term adaptive responses of fitness, IUE and physiology to Fe limitation and warming interactions in these four major cyanobacterial groups; 2) Determine the molecular and biochemical mechanisms behind the surprising Fe/warming interactive effect on IUE using genomics, transcriptomics and quantitative proteomics coupled with 'metalloproteomics' determinations of Fe content in critical proteins; 3) Compare and contrast acclimation and adaptation responses to Fe limitation and warming in key cyanobacteria taxa, and 4) Integrate results using a published biogeochemical modeling approach to assess global consequences for marine productivity and nitrogen fixation. This project offers a mechanistic and predictive understanding of adaptation to Fe and warming co-stressors in a rapidly changing future ocean environment for some of the most important photoautotrophic functional groups in the ocean.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[ [table of contents](#) | [back to top](#) ]

---

## **Program Information**

### **Marine Microbiology Initiative (MMI)**

**Website:** <https://www.moore.org/initiative-strategy-detail?initiativeld=marine-microbiology-initiative>

A Gordon and Betty Moore Foundation Program.

Forging a new paradigm in marine microbial ecology:

Microbes in the ocean produce half of the oxygen on the planet and remove vast amounts of carbon dioxide, a greenhouse gas, from the atmosphere. Yet, we have known surprisingly little about these microscopic organisms. As we discover answers to some long-standing puzzles about the roles that marine microorganisms play in supporting the ocean's food webs and driving global elemental cycles, we realized that we still need to learn much more about what these organisms do and how they do it—including how they evolved and contribute to our ocean's health and productivity.

The Marine Microbiology Initiative seeks to gain a comprehensive understanding of marine microbial communities, including their diversity, functions and behaviors; their ecological roles; and their origins and evolution. Our focus has been to enable researchers to uncover the principles that govern the interactions among microbes and that govern microbially mediated nutrient flow in the sea. To address these opportunities, we support leaders in the field through investigator awards, multidisciplinary team research projects, and efforts to create resources of broad use to the research community. We also support development of new instrumentation, tools, technologies and genetic approaches.

Through the efforts of many scientists from around the world, the initiative has been catalyzing new science through advances in methods and technology, and to reduce interdisciplinary barriers slowing progress. With our support, researchers are quantifying nutrient pools in the ocean, deciphering the genetic and biochemical bases of microbial metabolism, and understanding how microbes interact with one another. The initiative has five grant portfolios:

Individual investigator awards for current and emerging leaders in the field.

Multidisciplinary projects that support collaboration across disciplines.

New instrumentation, tools and technology that enable scientists to ask new questions in ways previously not possible.

Community resource efforts that fund the creation and sharing of data and the development of tools, methods and infrastructure of widespread utility.

Projects that advance genetic tools to enable development of experimental model systems in marine microbial ecology.

We also bring together scientists to discuss timely subjects and to facilitate scientific exchange.

Our path to marine microbial ecology was a confluence of new technology that could accelerate science and an opportunity to support a field that was not well funded relative to potential impact. Around the time we began this work in 2004, the life sciences were entering a new era of DNA sequencing and genomics, expanding possibilities for scientific research – including the nascent field of marine microbial ecology. Through conversations with pioneers inside and outside the field, an opportunity was identified: to apply these new sequencing tools to advance knowledge of marine microbial communities and reveal how they support and influence ocean systems.

After many years of success, we will wind down this effort and close the initiative in 2021. We will have invested more than \$250 million over 17 years to deepen understanding of the diversity, ecological activities and evolution of marine microbial communities. Thanks to the work of hundreds of scientists and others involved with the initiative, the goals have been achieved and the field has been profoundly enriched; it is now positioned to address new scientific questions using innovative technologies and methods.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)	<a href="#">GBMF3934</a>
<a href="#">Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)</a>	<a href="#">GBMF3782</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1657766</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1850719</a>

[ [table of contents](#) | [back to top](#) ]